AGRICULTURAL CHEMICALS

In This Issue:

Posticide Dévelopment Costs Evaluated

NACA to Annual Sering Lake Meeting

Fertilizer Prespect

Fungicide Residues

Fortifixor Sofety Meeting in October

Control Officials to Washington Meeting

NFA to Hold Florida

Ohio Pesticide Tour

National Shade Tres Conference Meets



deliver the final kill ... with



Late-season spraying-the last measure of defense against insect damage-calls for materials of high toxicity and sure-fire effectiveness.

Whether you need Pyrethrum and Rotenone on fall vegetable crops-or TEPP (Tetraethyl Pyrophosphate) or PARATHION on cotton-you can get this high toxicity and effectiveness with POWCO BRAND insecticides.

But whatever your needs . . . in any season . . . look to POWELL.

John Powell & Co., Inc. ONE PARK AVENUE, NEW YORK 16, N. Y.

TOXAPHENE BHC CHICRDANE UNDANE POTENONE SABADINA FYRETHRUM & PYRING FIFERONYL BUTCHIDE ANTU TEPP ALDRIN 2, 4 D & 2, 4, 5-T



METHODS and RESULTS

Many different methods of promoting Grasslands Farming are being pursued. All promote greater productivity and profit for our farms.

The program shows tangible results in millions of acres of improved pasture land and greatly increased farm income.

To the agencies who initiated and are successfully spreading the gospel of "Green Pastures" we pledge our continued support.

POTASH COMPANY OF AMERICA Carlsbad, New Mexico

GENERAL SALES OFFICE . . . 1625 Eye Street, N. W., Washington, D. C. MIDWESTERN SALES OFFICE . . . First National Bank Bldg., Peorio, III. SOUTHERN SALES OFFICE . . . Candler Building, Atlanta, Ga.



will improve

production product profit

The user of Attaclay knows this to be the case.

For—during six seasons of severe appraisal—he's seen

Attaclay do the extraordinary as well as the ordinary jobs
demanded of a carrier. Results:

He's improved production. Attaclay accepts higher percentages of liquid and low melting point toxicants, yet the product remains lump-free, dry, free-flowing.

He's improved product. Attaclay gives across-the-board flowability to a high degree—throughout the plant from grinder to packager, after months of storage, in the customer's hands.

He's improved profits. Attaclay drives unit costs down by stepping up plant capacities 30-50%. The increased production it makes possible is a real boon in the rush season. It meets the most exacting formula requirements. It is the trade-accepted standard for use with all of the popular poisons, including such newcomers as aldrin, dieldrin and dilan.

It is logical, therefore, that a large majority of experienced agricultural chemical producers should be—as they are—so unqualified in their praise of Attaclay. Its continuing leadership is convincing evidence.

If you are not yet acquainted with Attaclay, may we work with you?

ATTAPULGUS CLAY COMPANY

Dop's P. 210 West Washington Squere, Phile. S. Po.

AGRICULTURAL CHEMICALS



A Monthly Magazine For the Trade

LAWRENCE A. LONG
Editor

JOHN A. NICHOLSON Advertising Manager

Editorial Advisory Board . . .

DR. ALVIN J. COX Palo Alto, Calif.

LEA S. HITCHNER Washington, D. C.

Dr. S. E. A. McCallan Yonkers, N. Y.

Dr. CHARLES E. PALM Ithaca, N. Y.

DR. COLIN W. WHITTAKER Washington, D. C.

THIS MONTH'S COVER

Perfectly-formed fruit free from damage by insects or plant disease is made possible by timely and accurate application of modern agricultural chemicals. Here a new-type sprayer is at work in apple orchard. Photo by F. E. Myers & Bro. Co., Ashland, Ohio.

VOL. 7 No. 9 SEPTEMBER 1952

In This Issue:

Editorials	2)
Guest Editorial	31
Cost of Marketing New Products	32
The Story of Heptachlor	35
NAC Association to Spring Lake	38
Fertilizer Prospects for Years Ahead	40
Dithiocarbamate Fungicide Residues	45
Fertilizer Safety Meeting Planned	50
Control Officials to Meet	52
NFA to Florida Meeting	55
Supplying Food for the "Fifth Plate"	57
The Listening Post	61
By Paul R. Miller & Kelvin Dorward	70
Suppliers Bulletins	
Technical Briefs	-
Ohio Pesticide Institute Tour	
Shade Tree Conference Meets	
Industry News	
Classified Advertising	144
Advertisers' Index	145
Industry Meeting Calendar	145
Tale Ends	146

AGRICULTURAL CHEMICALS

Subscription Rales: One year, United States, \$3.00; Canada and Pan American countries, \$4.00; Foreign, \$7.00. Published monthly on the 15th by Industry Publications, Inc. Wayne E. Dorland, President; Ira P. MacNair, Secretary-Treasurer. Publication office, 123 Market Place, Baltimore 2, Md. Advertising and editorial office 175 Fifth Ave., New York 10, New York — Chicago Office, 338 N. Michigan Blvd. Advertising rates made known on application. Closing date for copy—15th of the month previous to date of issue.

Entered as second-sines matter November 4, 1949, at the Post Office at Baltimore, Md., under the Act of March 8, 1879.



A WHOLE FARM THRIVES ON NOURISHMENT

Kittens or cabbages . . . sheep or alfalfa . . . growth on a farm is a relentless process.

This never-ending struggle for survival and growth is a tremendous drain on the rich plant-food elements within the soil. No matter how deviously, every living thing must look to the soil for its basic nourishment.

Nature often cannot replenish these vital soil ingredients, and fertilizers containing FOTASH are used. Sunshine State Potash, a product of New Mexico, helps make such fertilizers more than a mere soil nutrient. It strengthens the crops . . . aids in effectively resisting disease and drought.



UNITED STATES POTASH COMPANY, Incorporated, 30 Rockefeller Plaza, New York 20, N. Y.



PONTANEOUS DISPERSIO

TOXIMUL

a balanced anionic emulsifier for agricultural insecticides

TOXIMUL is the family name of Ninol Laboratories' new series of Emulsifiers, designed to improve performance and reduce costs of biocidal spray concentrates. Various members of the series have been developed for use with different types of insecticides such as Toxaphene, Chlordane, DDT, etc.

TOXIMUL 100, for example, features spontaneous dispersion when used in Toxaphene emulsion concentrates. When added to water of any hardness, concentrates containing TOXIMUL 100 form stable emulsions with a minimum of agitation. Such concentrates are therefore ideal for general agricultural use, and particularly where agitation conditions are poor . . . as in aeroplane sprayers, knapsack tanks and cattle dips.



NINOL LABORATORIES

1719 S. CLINTON • CHICAGO 16 • PHONE CHESAPEAKE 3-9625

In Canada Chemical Developments of Canada Ltd., 420 Lagauchetiere Street W., Montreal 1, Quebec

ND THIS COUPON TODAY! MINOL LABORATORIES, DEPT. A 1719 S. CLINTON ST., CHICAGO 16, ILL. Please send me sample and further in.

formation about TOXIMUL emulsifiers. Our O Toxaphene Chlordono [lether) 0 DDT

TITLE COMPANI

ZONI STATE





Your dry dust concentrate will give top field performance—even though dense let-down agents have been added—when you use Celite.

You'll find that the presence of a small percentage of Celite—because of its great bulk per unit of weight, and the irregular shape of the individual particles—fluffs up the final dust and helps neutralize the effect of heavy extenders. Better dispersion in the field is thereby assured . . . at very little cost.

Celite offers other advantages, too:

As a Grinding Aid: Celite aids in grinding of both solid and semi-solid low melting point poisons and makes a more potent dry dust concentrate.

As an Absorption Medium: Celite's high liquid absorption capacity allows a higher percentage of liquid poison to be absorbed while still maintaining a "dry" dust. This gives you a highly concentrated product, and lower packaging and shipping costs. The use of Celite as a diluent and grinding aid can well be a profitable one for your operation. For additional information, write to Johns-Manville, Box 60, New York 16, N. Y.

Properties of CELITE

FINENESS: Approximately 100% through 325 mesh DENSITY (Vibrated): 11 pounds per cubic foot BUK: Celtie bulks much higher than most diluents ABSORPTION: 200% of its weight of water 300% of its weight of kerosene pH VALUE: Below 7.0

pH VALUE: Below 7.0 INERTNESS: Compatible with insecticide and fungicide

poisons
SUSPENSION: Excellent in both air and water

COMPOSITION: Celite is amorphous diatomaceous silica (SiO:)

Celite is Johns-Manville's registered trade mark for its diatomacoous silice products.





Johns-Manville CELITE

DILUENTS AND GRINDING AIDS



The addition of new, modern and complete Multiwall Bag manufacturing facilities at our new Los Angeles plant, brings to West Coast Multiwall Bag users, Fulton's experience and know-how, and an unbroken record of dependable service in fulfilling every customer commitment on time.

Call on Fulton for all your Multiwall Bag requirements — pasted or sewn bottom, open mouth or valve, in every size!

Fulton
BAG & COTTON MILLS

LOS ANGELES, 4901 Se. Boyle * Affente * St. Louis * Belles

Benver * New Orleans * Minneapelis * Kansas City, Kans. * New York City, 347 Medison Avenu

in insecticidal, fungicidal and herbicidal sprays

to really make a killing

there's nothing like Nacconol

as the wetting agent

to wettable DDT BHC, 2,4-D; Chlordane, Sulfur and similar products and mixtures. Results are instantly observable in increased coverage and wetting which means a better kill.

Flake, powder or bead form-all rapidly soluble in cold water at any pH. For samples, prices and suggestions on

RECTOR STREET, NEW YORK 6. N.Y. . BOwling Green

Baston 14, Wass., 150 Causeway St. Prevolence 3, # 1, 15 Westmonster St. Phriadalphia G. Pa., 200 284 S. Front St. San Francisco S, Cal., 517 Howard St. Partland S, Bre., 730 West Burnside St. Chicago 54, III., The Worchandise Wart

CAptal 7-0499 SEctor 1-2008 L0mbard 3-6382 SUtter 1-7597 Stigeria: 7-3387 Charlette, 1 N.C., 281-283 West First St. Charlette 3 9721

Butmand 19, Va., 8 North Fidth St. **Bichmani 7 1939** Columbus, Ga., Columbus Interstate Bidg Greensbara, M.C., Jefferson Standard Bidg. CReensbare 2-2518 Chattaneego 2, Tenn., James Building Bilanto 2, Sa., 254 E. Paces Ferry Bg. CYPIess 2821 Hew Orleans 18, La., 714 Carandalet Building Toronto 2, Canada, 137-143 Wellington St. W.





Columbia-Southern has just completed extensive tests on Hi-Sil, a finely divided hydrated silica, and has proved this product excellent for formulating 75% DDT dusts and wettable powders.

Highly concentrated DDT powders offer many advantages, the three principal being reduced shipping costs, especially on long hauls, reduced packaging costs, and its application in the rapidly developing field of concentrate sprays.

Columbia-Southern's Hi-Sil was originally

developed as a reinforcing pigment for rubber products where it has gained wide recognition and acceptance. Since Hi-Sil is a very finely divided hydrated silica, having uniform particle size, high absorbency, and good fluidity, it was decided to put Hi-Sil to exhaustive tests to evaluate its usefulness as a carrier for DDT and other agricultural chemicals.

This research shows Hi-Sil as the most promising diluent to support higher DDT loadings. As a result, the major part of the DDT powder made in the United States is now produced using Columbia-Southern Hi-Sil.

INVESTIGATE TODAY!

A paper has been published by the Columbia-Southern Technical Department based on this study which discusses the findings in detail. You are invited to write for it today. Address your inquiries for this paper or for any other information or working samples to our Pittsburgh address.

COLUMBIA - SOUTHERN CHEMICAL CORPORATION

SUBSIDIARY OF PITTSBURGH PLATE GLASS COMPANY FIFTH AVE. AT BELLEFIELD • PITTSBURGH 13. PA.



DISTRICT OFFICES: BOSTON • CHARLOTTE • CHICAGO • CINCINNATI • CLEVELAND • DALLAS • HOUSTON • MINNEAPOLIS • NEW ORLEANS • NEW YORK • PHILADELPHIA • PITTSBURGH • ST. LOUIS • SAN FRANCISCO

High Grad Auriate of Potash!



DUVAL SULPHUR and POTASH COMPANY Modern New Plant and Refinery at Carlsbad, New Mexico

Address all communications to: ASHCRAFT-WILKINSON CO.

Exclusive Distributors ATLANTA, GEORGIA Cable Address ASHCRAFT

. CHARLESTON. S. C. . TAMPA. FLA. . COLUMBUS, OHIO





FOR FLY Control

for SAFETY'S SAKE use EMULSIFIABLE SULFOXIDE-PYREXCEL 1-10

(PYRETHRINS PLUS SULFOXIDE)



Emulsifiable Sulfoxide-Pyrexcel 1-10 in proper dilution is safe for direct application to dairy cattle and other livestock.


In the Dairy Barn
In the Milk Handling and
Milk Processing Rooms

In fact, wherever a safe insecticide is required.

FOR INFORMATION WRITE TO

S. B. PENICK & COMPANY

COTTON FOLIAGE TAKES FLIGHT



Niagarathal-DF Spray

A Real "Down-To-Earth" Cotton Defoliant

Niagarathal-DF Spray is a concentrated liquid containing Endothal*, considered to be the first organic defoliant. When properly applied Niagarathal-DF Spray produces a physiological action on cotton plants which approaches the "natural" defoliating action of the plant.

Niagarathal-DF Spray dilutes readily in water and requires no agitation. Being a liquid there is no nozzle clogging or abrasion. And Niagarathal-DF Spray is safe to apply in any suitable equipment since it presents no fire hazard.

Thoroughly tested in the cotton belt, Niagarathal-DF Spray has proved to be a thoroughly effective defoliant that has a wider timing range than similar products. Growers report increased returns on acres defoliated with Niagarathal-DF Spray. Write for literature.

*Trade Mark—Sharples Chemicals Inc., Subsidiary of Penna Solt Mfg. Co.

QUICK FACTS ABOUT COTTON DEFOLIATION

- Permits earlier harvests. Boll rot is prevented or greatly reduced. Fibre and seed spoilage are retarded.
- Defaliated cotton is easier to pick.
- Machine harvesting can be done more efficiently. Better cotton grades are obtainable.
- Defoliation is an aid to cotton insect control. Damage by aphids and late leaf worm is prevented. Boll weevil populations are reduced.

CHEMICAL DIVISION

FOOD MACHINERY AND CHEMICAL CORPORATION

MIDDLEPORT, N. Y. Richmond, Calif., Jacksonville, Tampo, Pompano, Fla., New Orleans, La:, Ayer, Mass., Greenville, Miss., Harlingen and Pecos, Tex., Yakima, Wash., Subsidiary: Pine Bluff Chem-ical Co., Pine Bluff, Ark. Canadian Associate: NIAGARA BRAND SPRAY CO., LTD., Burlington, Ont.



The ALL-PURPOSE Unit

· For Producing

HIGH CONCENTRATE INSECTICIDE POWDERS

RAYMOND

WHIZZER EQUIPPED

ROLLER MILL


This is a wide utility grinding mill which may be used for economically handling all of your insecticide powder production . . . whether it involves regular field strength, high concentrates or sulphur.

The sturdy construction of this modern Raymond Mill makes it capable of continuous 24-hour operation. It is a slow speed, cool running unit, and it does not require artificial cooling of the material. It has the additional advantages of automatic handling of the material and dust-free operation.

Whatever your capacity requirements or dust formulating problem may be, there is a Raymond Mill for handling your job economically to meet today's maximum specifications. For further details write for Insecticide Bulletin No. 68.



RAYMOND LABORATORY MILL grinding small samples of material



RAYMOND ROLLER MILL showing Double Whizzer Unit

COMBUSTION ENGINEERING - SUPERHEATER, INC.

1314 NORTH BRANCH ST. CHICAGO 22. ILLINOIS

Kaymond PULVERIZER DIVISION

SALES OFFICES IN PRINCIPAL CITIES looking for

IMP DVEMENTS?

EMULSIFIER L

When you start looking for ways to improve your formulations of insecticides and herbicides, investigate Monsanto Emulsifiers L and H.

a mobile anhydrous liquid, is a carefully balanced blend of surface-active agents developed to give quickforming, stable emulsions of: Chlordan; DDT; 2,4-D esters; 2,4,5-T esters; Aldrin; Dieldrin; trichlorobenzene; Lindane; parathion; tetraethyl pyrophosphate.

EMULSIFIER H


also an easy-to-handle, mobile liquid, is built specifically for preparing emulsifiable concentrates of pentachlorophenol.

EMULSIFIER



Write for a free copy of the Monsanto leaflet, "Emulsifiers for the Agricultural Industry," giving 19 sample formulations employing Emulsifier L. For quotations and other information on Monsanto Emulsifiers, contact the nearest Monsanto Sales Office or MONSANTO CHEMICAL COMPANY, Organic Chemicals Division, 1700 South Second Street, St. Louis 4, Missouri.

DISTRICT SALES OFFICES: Birmingham, Boston, Charlotte, Chicago, Cincinnati, Cleveland, Detroit, Houston, Los Angeles, New York, Philadelphia, Portland, Ore., San Francisco, Seattle, Twin Cities. In Canada, Monsanto Canada Limited, Montreal.



This man may have been good with a wheelbarrow and hand shovel, but his productive capacity is several times greater as a "PAYLOADER" pilot. He's happier and so is his boss, because they both make more money.

"PAYLOADER" tractor-shovels are helping fertilizer and chemical plants slash production costs. They actually pay for themselves in a few months doing the many jobs listed here . . . doing them faster and cheaper — and release manpower for more productive work.

These special tractor-shovels are available in seven sizes, from 12 cu. ft. to 1½ cu. yd. to master your jobs both indoors and outdoors. Get full facts now on cost-cutting, time-saving, production-boosting "PAYLOADER". The Frank G. Hough Co., Sunnyside Ave., Libertyville, Illinois.

FOR MANY JOBS. Unload box cars — dig and carry fertilizers, chemicals and all loose materials — feed conveyors, hoppers, baggers, elevators — unload trucks — load box cars — carry bagged materials — clean up aisles, gangways — maintain private roadways — stockpile and carry coal — handle ashes — remove snow — spot cars — lift, haul, push, pull.



PAYLOADER



end "Industrial Handling" is full of profitable ideas developed by "PAYLOADER" users. A request on your letterhood gets you on the regular moiting list without cast.



75% DDT WETTABLE POWDER

when you want it promptly



Check with Michigan Chemical for your continuous, fast supply of DDT. You can be sure of exceptionally high purity, uniform production, and prompt service.

Strategically located in the midwest agricultural area, Michigan Chemical is a basic producer of DDT for Agricultural use.

This, together with favorable transportation facilities, assures you a most satisfactory source. Just as a trial, place your next order with Michigan Chemical. You'll get results that surpass promises.






MICHIGAN CHEMICAL CORPORATION . St. Louis, Michigan

SALES OFFICES: 75 East Wacker Drive Chicago 1, Illinois 230 Park Avenue New York 17, New York

BASIC MANUFACTURERS OF INDUSTRIAL PHARMACRUTICAL AND AGRICULTURAL CHEMICALS



Emulsions last longer — disperse faster — with Armour's Etho-chemicals

Armour's oil-soluble, water-dispersible Etho-chemicals make any emulsification easier and faster—and longer-lasting. For instance, Ethomeen S/12 and Ethomeen S/15, mildly cationic chemicals which are not affected by water hardness, combine to form an excellent emulsifier for 2,4-D isopropyl ester, even in extremely low concentrations. Another excellent emulsifier is Ethofat 142/20, a non-ionic chemical for use with kerosene or xylol as a solvent. Chlordane can be emulsified directly into water with this chemical, without a solvent.

Write today for complete information about these and Armour's other emulsifiers, including formulas, methods of use, and prices.



ARMOUR CHEMICAL DIVISION

Armour and Company, 1355 W. 31st St., Chicago 9, Ill.



For proof of acceptance, we have referred in the past to users of DILUEX whose annual consumption amounted to trainloads. Whatever statement was made in such terms a year ago would be an understatement now. On merit, the demand for DILUEX has multiplied.

In impregnating liquid toxicants-mill-

ing DDT, BHC, or other organics—conditioning blended dusts, DILUEX gives the greatest assurance of quality in the finished product.

Production facilities have been enlarged to keep pace with all anticipated requirements. Inquiries and orders will receive prompt attention.

FLORIDIN COMPANY

. ADSORBENTS ... DESICCANTS . . . DILUENTS

Dept. M., 220 Liberty Street, Warren, Pa.



LION ANHYDROUS AMMONIA—For formulation. A uniformly high-quality basic product. Nitrogen content, 82.25%.

LION AQUA AMMONIA—For formulation or acid oxidation. Ammonia content about 30%. Other grades to suit you.

LION NITROGEN FERTILIZER SOLUTIONS—For formulation.

Three types to suit varying weather and manufacturing conditions.

LION AMMONIUM NITRATE FERTILIZER—For direct application or formulation. Improved spherical pellets. Guaranteed 33.5% nitrogen.

LION SULPHATE OF AMMONIA—For direct application or formulation. Large free-flowing crystals. Guaranteed nitrogen content, 21%.



Serving Southern States Lion provides special technical assistance for fertilizer manufacturers. Write us if you have a formulation problem.

LION OIL COMPANY CHEMICAL DIVISION, EL DORADO, ARK.



Now-for peat moss and humus: paper package that won't rot

made possible by coating of Du Pont ALATHON*

Coating of "Alathon" preserves paper from attack by moisture and micro-organisms

At last, producers of peat moss and humus can be sure their packaged product will stay packaged... in multi-wall bags coated on the inside with Du Pont "Alathon" polythene resin. This is a typical example of how "Alathon" can provide better packaging for so many different chemical products.

Before this bag lined with "Alathon" was developed, all conventional packages then being used sooner or later disintegrated because of moisture and micro-organisms in the peat moss.

The coating of "Alathon" inside this multi-wall bag cannot be harmed by micro-organisms—it slows vapor transmission—and the bag isstronger drop-breakage tests proved it) than all other bags tested! What's more, it protects the peat moss or humus by keeping air out, moisture sealed in

Coating of "Alathon" on packaging materials provide a unique combination of properties, offering new ways to meet many packaging needs in the agricultural chemicals industry. "Alathon" is tough, strong and scuff-resistant—chemically inert—tasteless, odorless, non-toxic—resists vegetable oils, acids, alkalis—stays flexible at 70°F. below zero!

... Write for free booklet describing the properties and uses of "Alathon" in the packaging field. We'll gladly put you in touch with sources of supply for packaging materials coated with "Alathon."

E. I. du Pont de Nemours & Co. (Inc.) Polychemicals Dept. 9E, Du Pont Building Wilmington 98, Delaware



1000 pounds of 10-10-10 plowed down nearly doubles corn and potato yields



NORTHUMBERLAND, PA.



• Until 1948, Leon Epler of Northumberland, Pa., was getting corn yields of 40 to 50 bushels per acre by using approximately 250 pounds of row fertilizer. But that year he began plowing down 1000 pounds of 10-10-10 in addition to the row fertilizer. His yield jumped to 70-75 bushels per acre.

He climaxed this outstanding production in 1950 with a plot that yielded 122.1 bushels per acre in the Five Acre Corn Club contest sponsored by the Pennsylvania Crop Improvement Association. The special fertilizing program used on this plot included 10-10-10.

Plowing under 800-1000 pounds of 10-10-10 gave Mr. Epler equally outstanding results on potatoes. His average yield jumped from 300-350 bushels per acre to 500-550 bushels. Improved spraying and cultural methods also played a role in this increase.

Bigger yields for farmers mean better business for you

 On all types of crops, high-nitrogen complete fertilizers are paying for themselves over and over through higher quantity and better quality yields

Every bumper crop produced by high-nitrogen fertilizers increases the demand for these products. And now is the time for you to cash in on this demand by putting heavy promotional effort behind the high-nitrogen fertilizers in your line.

When you use U·S·S Ammonium Sulphate in your mixed fertilizers, you'll find the selling job

easier. Farmers know the name "United States Steel" stands for top quality. Many of them have used U-S-S Ammonium Sulphate as a straight-nitrogen material... they know it's dry and freerunning... they know it won't set up in storage and that it handles well in distributing equipment.

For complete details on U·S·S Ammonium Sulphate contact our nearest Coal Chemical Sales Office or write directly to United States Steel Company, 525 William Penn Place, Pittsburgh 30, Pennsylvania.

U·S·S AMMONIUM SULPHATE



2-1421

UNITED STATES STEEL



You can <u>depend</u> on *Pittsburgh* Parathion to deliver peak quality every time!

You can rely on Pittsburgh Parathion to give you top performance because it's Quality-Controlled from coal to finished product at our own agricultural chemical plant. That's why Pittsburgh Parathion is a better answer to the problem of controlling most principal insect pests that attack fruit, vegetable and forage crops.

We'll gladly send you information bulletins giving the complete story on Pittsburgh Quality-Controlled Parathion. Write today!

PITTSBURGH AGRICULTURAL CHEMICAL COMPANY

7601 EMPIRE STATE BUILDING, NEW YORK 1, NEW YORK

OFFICES: Saw Francisco * Deuter * Dallas * St. Louis * Chicago * New York * Charlotte * Toronto (P.O. Box 111, Station K) * Montreal (1100 Craig St., East)

Standard for Quality
ORGANIC INSECTICIDES: Benzene Hexachloride, Toxaphene, Dichloro Diphenyl Trichloroethane, Aldrin, Dieldrin, Chlordane.

ORGANIC PHOSPHATE INSECTICIDES: Parathion Wethoble
Powders, Parathion Liquid Concentrate, Metacide. WEED

RILLERS: 2,4-D Acid, 2,4-D Amine Concentrates, 2,4-D Ester
Formulated Concentrates, 2,4-D Sodium Solt Monohydrate,
2,4,5-T Formulations. PUNGICIDES, SEED DISINFECTANTS,
COTTON SPRAYS AND DUSTS, AND OTHER SPECIAL AGRICULTURAL CHEMICALS. CULTURAL CHEMICALS.


COAL CHEMICALS . AGRICULTURAL CHEMICALS . PROTECTIVE COATINGS . PLASTICIZERS . ACTIVATED CARBON . CORE . CEMENT . PIG IRON



FOR PROMPT SERVICE FROM

VO STRATEGICALLY LOCATED
PLANTS





Hammond "personalized service" is more than a slogan. Call our nearest representative.



HAMMOND BAG & PAPER CO.

General Offices: Wellsburg, W. Va. Plants in Wellsburg, W. Va. and Pine Bluff, Ark.





Naval Stores Department

HERCULES POWDER COMPANY

970 Market Street, Wilmington 99, Delaware

MAKERS OF TECHNICAL TOXAPHENE FOR AGRICULTURAL INSECTICIDES



THE EDITOR COMMENTS



GOOD many reasons are being set forth as to the cause of the current situation in the insecticide trade. Why did the past season's business

go sour? Was it widespread over-production? Was it the lack of insect infestations where heavy populations had been anticipated? Or is the industry somewhat at fault in failing to carry out an educational program on the use of insecticides as a *preventive* measure?

Probably all of these factors, and more, entered into the picture, but in the opinion of self-critical industry spokesmen, some manufacturers have not yet learned how to regulate their new potential for tremendous production. Output of most technical insecticidal chemicals has followed a straight uphill pattern for the past 3 or 4 years, with the exception of the ancestor of all the newer organics, DDT, which has experienced valleys and peaks during the past several seasons.

Looking back to the beginning of the 1951-52 season, inventories were regarded as being relatively low so far as manufacturers and distributors were concerned, although there appears to be no particular agreement about amounts on hand. But the reaction to this condition seemed to be unamimous... full production! As one industry observer described it, "The spigots were turned high, wide and handsome," with the results of "fantastic" production of material, with no apparent reason. Production of DDT is reported to have reached 130 million pounds per year, with BHC output even higher!

With this heavy production and no major insect infestation to consume great quantities of pesticides, it is easy to see how the industry could paint itself into a corner. We recall some talk in the March-April period that the rate of output was too high at that time... but still the production continued until just about all levels of the distribution setup were practically suffocating by August.

Yet, the element of chance is ever present. Suppose production were to be cut down and inventories allowed to be reduced, and then a widespread need for pesticides should arise sud-renly! The industry must be prepared for such an eventuality, but at the same time should not over-produce! How to determine the happy medium between superabundance and too little is a mighty difficult problem in this business.

We can't presume to have a ready-made answer to this basic problem, but we do feel confident that this year's experience will act as a guide to scheduling production in future seasons.



ITH its announcement of a new Speakers' Bureau, the California Fertilizer Association has taken a step forward toward a good public

relations program which should bear fruit in that area. Service and women's clubs, garden clubs and other local organizations have been invited to ask for speakers to address their meetings, and the CFA promises to have a qualified person available on short notice.

This step seems important to us mainly because it is in groups of this character that the "organiculturists" get in their best licks against the use of commercial fertilizer materials. One has only to talk five minutes to a club lady who has been subjected to anti-fertilizer propaganda, to see how deeply ingrained is their feeling. We are glad to see the CFA acting to offset this, and hope that its efforts will be successful.



HE way in which medical reports can be twisted out of their original context and meaning is illustrated once more in the case of a Hercules

Powder Company report on toxaphene toxicity. Reporters, either not reading the medical report carefully, or else purposely exaggerating its sensational parts, came up with some rather frightening interpretations. The threat of such "scare" copy isn't over yet, although it has slowed down of late. Education of the user seems still to be the best defense against it.


BARDEN CLAY

The systemating low-cost pesticide diluent

If you use diluents, you will want a copy of this new Barden Clay bulletin. You will find in it tables and descriptions of the unique properties of this low-cost pesticide diluent. Write for your copy, today

J. M. HUBER CORPORATION
100 PARK AVENUE, NEW YORK 17, N. Y.






It's time for Long-Range Planning

Arthur W. Mohr

President. California Spray-Chemical Corp. Richmond. California President, National Agricultural Chemicals Assn.



What happened this season in the manufacture, processing and distribution of pesticides illustrates the peculiarities of our industry - peculiarities that point up the necessity for planning that should extend, not from one season to the next, not for 2, 3, 5 or 10 years but from 15, 20 or 25 years and longer.

The 1952 season did not develop into a severe situation in regard to insect infestation. In the South, where the biggest chunk of our production is used for the protection of cotton, insect infestations did not materialize due to weather conditions adverse to their development. Other crop pests such as fruit insects and diseases, vegetable insects and diseases, weeds and other pests of field crops helped to drain off the huge and varied supply of pest control chemicals available.

In the main, and across the board, this was not a good year for the pesticides industry. It is true that some manufacturers and distributors, with a particular chemical in good demand



ARTHUR W. MOHR NACA President

for a unique pest, had a highly successful year; but this was far from the typical situation.

Producing at the rate of about 11/4 billion pounds of technical pesticidal chemicals annually, the industry was in full production prior to the beginning of the crop season. Distribution of the technical chemicals was carried out rapidly and formulators soon were stocked to capacity. To match the agricultural goals of the U.S. Department of Agriculture, the industry expanded plant facilities and built new plants to meet an expected demand for pesticides. In cooperation with the U. S. Department of Agriculture and Land Grant Colleges, the industry attempted to get dealers and growers to buy products early. But the early buying program broke down in the face of an adequate supply of pest control chemicals reported to be available for the current season. This season, it was possible for farmers to purchase enough pesticides to protect their crops without jeopardizing distribution facilities.

Due to the importance of agricultural chemicals in meeting the production goals of the Department of Agriculture, the pesticides industry was able to obtain sufficient raw materials

(Turn to Page 131)

Economics of Developing

Agricultural Chemicals

N the United States, highly organized industrial research laboratories, with their company's raw materials available, are a prolific source of new chemicals. Company biologists, accustomed to screening techniques, rapidly sort out those new chemicals which appear promising for agricultural uses. Then follows widespread field work, usually in experiment stations and the United States Department of Agriculture. It is the purpose of this paper to estimate the average cost involved in bringing to the market a new agricultural chemical through this sequence. The term "agricultural chemicals" is used here for convenience to refer to chemicals other than fertilizers which are used on the farm. Most of the emphasis will be placed on insecticides, fungicides and herbicides.

Four Testing Methods

THERE are at least four ways by which chemicals to be tested for agricultural uses may be selected:

 All chemicals available to the company, or made from other research projects, may be screened as agricultural chemicals. A company's basic products or their intermediates in, for instance, synthetic fibers, might thus be tested.

 Chemicals may be made specifically for use as agricultural chemicals based on accumulated knowledge of active chemical groupings. Thus it was logical that several companies investigated the entire series of the dithiocarbamates and their metallic salts once these compounds had been shown to be fungicidally active.

3. Compounds may be synthesized which are closely related to, or which duplicate naturally occurring materials. The synthesis of allethrin by the U.S.D.A. group was based on a knowledge of the chemistry of the naturally-occurring pyrethrin.

4. A study of physiology of the organism to be controlled may reveal that a given enzyme poison or antimetabolite may be indicated. For example, warfarin, the anti-vitamin K rodenticide, was selected on the basis that it was an effective anti-coagulant which could be readily antidoted by Vitamin K if accidentally ingested by humans or domestic animals.

In general, the cost, per compound, is least where materials are

Table 1.
Cost of Various Fungicidal Screening

Tests (after McCallan-194	6)
Slide germination	\$10
Greenhouse Early Blight	15
Greenhouse Late Blight	15
Phytotoxicity on tomatoes and	
beans	10
Pea Damping off	20
Mildew Resistance	10
Total	\$80

already on the shelf, and most where a physiological and biochemical study must precede the synthesis of the desired compound. However, the percentage of effective compounds is likely to be least in those generally available, increase appreciably where they have been specifically designed for biological activity and be still higher where they are synthetic analogues of some naturally-occurring products. The method of physiologic study is fraught with uncertainties since the metabolic processes are so complex. Actually, the case of warfarin is the only one that comes to mind where this approach has worked, though only infrequent efforts have been made along this line.

Most companies supply their chemicals through one or the other, or more usually both, of the first two approaches mentioned. For the purpose of attempting to arrive at some figures of the cost of agricultural chemicals let us assume that the company is making specifically for agricultural purposes each compound that is to be screened. In any calculation it is only fair that each compound bear its share of all expenses pertinent to operating a chemical laboratory. On such a basis, each initial synthesis may cost as little as \$25, if it is one of an homologous series of simple compounds. On the other hand, the cost may be as much as, or more than, \$1,000 if it is a new composition of matter made by

Dr. R. H. Wellman

Carbide & Carbon Chemicals Co.
Division of
Union Carbide & Carbon Corp.
New York

a synthesis with which the chemist is unfamiliar. The average cost might be between \$100 and \$200 per compound.

Compounds thus made are then transmitted to the biologists for screening. Dr. S. E. A. McCallan at Boyce Thompson Insitute in 1946 for his own information made some estimates on the costs of laboratory screening work with fungicides exclusive of administrative or overhead costs. These estimates appear in Table I. and apply only where many compounds are regularly being tested.

T would seem reasonable that costs have doubled since 1946, so that a similar screening program today would cost \$160 per compound. In light of our own experience, these figures seem reasonable, though a group set-up for this purpose and operating continuously might be able to complete initial fungicide screening for \$100 per compound. We believe the above tests are the minimum necessary to evaluate properly compounds as fungicides in screening tests and we subject all our compounds to about this number of tests before selections are made.

Insecticidal and herbicidal screening are equally complex and hence can be assumed to cost \$100 each per compound; though if all three are run together, certain duplications can be avoided such as phytotoxicity tests so that a figure of \$250 seems reasonable to screen a compound as an insecticide, fungicide and herbicide. It may seem unnecessary to screen materials that are specifically synthesized for biological activity for all three purposes.

Surely one should be able to tell in advance whether a compound is to be an herbicide, fungicide or insecticide. Maybe one should, but it is doubtful if any of those associated with such screening programs would rely on their judgment beyond being willing to assume that the next higher or lower member of a homologous series does not have to be tested if the parent compound was inactive. By using such an a-priori judgment screen, then the cost overall could come down to \$200 because as much as 1/5 the time the compound would be presumed to be inactive and would not be tested (thus running through a series of homologues as fungicides they would not all have to be tested as insecticides).

Compounds that survive screening will go through further laboratory or greenhouse testing in order to compare them as precisely as possible with the standards available. Also their stability, compatabilities and formulations must be investigated. The cost of such further work will vary greatly depending on their end use but since the work now becomes quantitative and usually must be such that it is suitable for statistical evaluation, it is rare that this work will be less than three times as expensive as the total screening cost and it may

be easily as much as ten times. An average might be five times or \$1,000.

Chemicals that survive this further work are customarily passed on to the toxicologist for initial appraisal. His work at this stage might consist of acute oral toxicity, skin absorption, eye and skin irritation tests at a cost of approximately \$400 per chemical, though if vapor hazards were involved, another \$100 would have to be added for range finding tests on this phase of the hazards.

Field Tests Necessary

I NITIAL field tests are now warranted. The fees charged by Dr. G. R. Townsend in Florida for initial field plot work are perhaps typical. He charges approximately \$15 per plot. The chemical to be tested must be compared with probably two standard materials. In this stage, the chemical may be examined in three different formulations. At least three concentrations for each and the necessary standards; also four replications will be needed. The bill for the test as outlined would be about 900.

The initial field tests, when they look promising, are followed usually by patent applications, development of analytical techniques for residue analyses and further work on its toxicology. Then must follow large scale field tests in several different areas on all purposed applications over at least a two-year period. It will be necessary also to conduct two-year feeding tests on rats to gain knowledge of the chronic hazards from a toxicological standpoint, to

Table II.

Cost of Development of an Agricultural Chemical
(after Persing—1951)

	Range		
Attorney's Fees	\$ 500 - \$ 5,000		
Field Studies (all charges)	50,000 50,000		
Toxicological Studies	20,000 — 30,000		
Analytical Methods	10,000 — 10,000		
Production Research and			
Pilot Plant Construction	75,000 - 250,000		
Administration	10,000 — 10,000		
Total	\$165,500 \$355,000		

conduct tests for residue and also for presence of the compound within the edible portions of the crops treated.

During this time, the samples available have been turned out laboriously first in large scale laboratory work, later in a developmental unit and perhaps by now, in a pilot plan designed for this chemical. Engineering design of a full scale plant must be well along so that, coupled with a survey of the market, the economics of making and selling the compound can be evaluated thoroughly.

Dr. C. O. Persing, Stauffer Chemical Co., gave some figures in a recent talk before the Association of Economic Control Officials on cost of this phase of development.

These estimates seem entirely reasonable and could be averaged at about \$250,000. As this stage develops, the experiment stations play an increasingly important part because of their unexcelled field testing facilities and because their obligation to serve their farmers makes it imper-

ative that they find for the farmer the best solution of his problems. I believe industry and state and federal workers are to be congratulated on the wholeheartedness and effectiveness of this cooperative endeavor.

Agricultural Government workers will now want to test the new chemical on their own, fitting it into their own programs, using it in many different ways and for many different purposes. Such a program never ends but is usually particularly heavy during the five years after the compound reaches the stage described above. Of course during a major portion of this period the compound is being sold, at least in a developmental way, but the purpose of this paper is to attempt to assess the actual experimental costs which may be assessed to such a material. If the compound is at all generally promising it will be tested by at least thirty experiment stations. They will also be examining other materials.

Perhaps a typical case is one

work its fair share of experiment station costs, including the physical plant and administration, it is obvious that the cost per compound per year can scarcely be less than \$1,000. This must be multiplied by thirty stations and five years to reach a cost of \$150,000. The company's contribution to this work is, customarily, to supply the compound free and to have a man who serves the purpose of liason between the company and the experiment station worker. If each of the stations uses 10 lbs. of material per year, a total of 1,500 lbs. has been supplied. In these amounts, the chemical delivered to the station will cost the company between \$2.00 and \$20.00 a pound with \$5.00 per pound being a possible average value for a total \$7,500. The man's time, plus his expenses, may be assessed against two or more products. By reason solely of the relatively small number of compounds reaching this stage, it is unlikely that the average will be more than two compounds per man. Half a man's time and travel expense for five years might be \$30,000.

where a research worker and an as-

sistant spend half-time investigating

ten materials. Assessing against this

In order to assess the cost per successful compound, it is necessary to estimate the number of failures and the stage at which failure is recognized. A few years ago there was a note in N. A. C. A. News that one company had screened 50,000 compounds as agricultural chemicals. In view of the number of companies interested, it seems entirely probable that at least 100,000 compounds had been screened by 1946. There has inevitably been a large amount of duplication but this has not affected the cost of the work done. A survey of the number of compounds selling in amounts more than 100,000 pounds a year which had been introduced in the last decade, found 35 such compounds. Perhaps a few were missed but if this experience is typical, it can be assumed that approximately 50 out of every 100,000 compounds examined, or one out of every 2000 compounds, may be expected to suc-

(Turn to Page 137)

Table III.

Costs Chargeable Against a Successful Agricultu	ral Ch	emical
Synthesis	_ 5	150
Initial Screening	1000	200
		350
1 in 30 succeed	oet	x 30
		10,500
Further Laboratory & Greenhouse work	***	1,000
		11,500
1 in 10 succeed		x 10
		15,000
Initial Field Tests		900
1 in 3 succeed		x 3
	2	47,700
Company development cost	. 2	50,000
	3	97,700
1 in 2 succeed		× 2
	1,1	95,400
Company liason with Experiment Station cost		37,500
	1,2	32,900
Cost of State and Federal Work	. 1	50,000
cost of successful chemical	. \$1,3	82,000

Properties and Effectiveness of

HEPTACHLOR

EPTACHLOR is one of a series of synthetic chlorinated hydrocarbons developed by the Velsicol Corporation to meet the ever-expanding need for materials to control agricultural pests. Early test results indicated the promise and value of this compound for the control of a wide variety of insect pests.

Following its initial screening evaluation, heptachlor was supplied to research workers in the United States Department of Agriculture Experiment Stations, University and State Experiment Stations and other field and research laboratories. These programs progressed concurrently, duing which time Velsicol's heptachlor formulations were tested and proven under actual field and commercial conditions.

From the results obtained, these formulations were found particularly effective in the control of several of the insect pests destructive to cotton. In addition to the insecticidal efficiency of heptachlor, tests made on cotton seed oil and cotton seed meal, taken from heptachlor-treated plots, showed no evidence of translocation.

Based upon this extensive and thorough testing program, the Velsicol product was recommended and given label approval for use in the control of several of the cotton insect pests.

Description and Properties

HEPTACHLOR is the common name which has been assigned to the compound 1,4,5,6,7,8,8-hepta-chloro-3a,4,7,7a-tetrahydro-4, 7-meth-anoindene

In the pure form, heptachlor is a white crystalline substance with a melting point of 95°.96° C. The total chlorine content of pure heptachlor is 66.5%.

Technical heptachlor is the term designated for the product as commerically produced. It consists of 67% heptachlor and 33% related compounds. This composition provides a convenient basis for the purpose of calculating formulations. To formulate on the basis of weight of actual heptachlor, it is only necessary to use one and one-half pounds of technical heptachlor for each pound of actual heptachlor required in the formulation.

Compatibility

FROM laboratory and field test results, heptachlor has been found compatible with most of the following commonly used insecticides and fungicides:

DDT
Chlordane
Dusting sulfur
Benxene hexachloride
Toxaphene
Dinitro compounds
Dithocarbamates
Quinones
Bordeaux Mixture

Types of Formulations

HEPTACHLOR can be applied in many formulations and has been used in both large and small scale tests as dusts containing from

TABLE I

Typical Properties of Technical Heptachlor

Physical Properties

Form @ Room Temperature	Soft, waxy solid
Melting Point Range	115°·165° F. (46.1°·73.9° C.)
Density @ 90° C.	
Rate of change of Density with Temperature	(0.002/degrees C. in the Range of 40°-95°C.)
Viscosity @ 90° C.	50-75 Centistokes
Rate of change of Viscosity with Temperature	(2 centistokes/degree C. in the range of 40°-60° C.)
Color	
Specific Heat	
Vapor Pressure	0,0003 mm @ 25° C.

Solubility

Xylene 40%		by	weight	
Methylated	Naphthalenes	40%	by	weight
Water		Insolu	able	

1 to 10%, as a 25% wettable powder as base oil solutions containing up to 10 percent and as emulsifiable concentrates containing 2 pounds of heptachlor per gallon, 2 pounds of heptachlor plus 2 pounds of DDT per gallon and 1 pound of heptachlor plus 2 pounds of DDT per gallon. The recommended formulations for cotton insect control are: 1. Heptachlor 2-1/2% Dust. 2. Heptachlor 2-1/2% — DDT 5% Dust.

3. Heptachlor 2-1/2% — DDT 5% Sulfur 40% Dust.

4. Heptachlor Emulsifiable Concentrate containing 2 lbs. of actual heptachlor per gallon.

 Heptachlor Emulsifiable Concentrate containing 1 lb. heptachlor and 2 lbs. DDT per gallon. Insecticidal Uses

ONTROLLING insects as both a contact and stomach insecticide, heptachlor affects both sucking and chewing insects. It acts as a stomach poison when used as a foliage treatment for leaf eating insects, such as grasshoppers, alfalfa weevils, and crickets. As a contact insecticide, it is effective against a large number of

Pests Controlled by Heptachlor and Suggested Concentrations

Insect Pest	Scientific Name	Dosages Required for Control
Agromoyzid leaf miner	Agromyza sp.	35-1 lb. actual heptachlor per acre showed promise in control.
Alfalfa snout beetle	. saconyan of	5 lbs. actual per acre showed only .43% plant injury in 2 weeks.
Alfalfa Weevil	Hypera postica (Gyll.)	2-16 to 4 oz. actual per acre has given control. It may be
		applied as an oil or water type spray.
Ambrosia Beetle	Scolytus sp.	A 5% oil solution applied to logs stacked at the mill or in forest gave good control of the ambrosia beetle.
Ant	Formicidae	Add 2-3 teaspoonsful of 25% wettable powder to 1 quart of water and pour in entrance to nest. Also 1 tablespoonful of 2-1/1/16 dust may be applied at the nest entrance.
Argentine Ant	Iridomyrmex humilis Mayr.	8 Ibs. of 25% wettable powder in 100 gallons of water has given good control when applied to the trunks or litter under trees. The emulsifiable concentrate gave 6 months residual pro- tection while wettable powder gave 4 months.
Black Vine Weevil	Brachythinus sp.	Soil applications of 5 to 10 lbs, per acre have given control Also 40 to 50 lbs, of 2-1,5% dust applied to alfalfa per acre gave control of adults.
Blow Fly	Calliphora sp.	.4 to .5 lb. per acre gave control.
Boll Weevil	Anthonomus grandis Boheman	Apply 5 to 8 lbs. of 2-5/2/4 dust or .15 lb. heptachlor as a spray when cotton is small. When full grown, apply 3/4 lb. of actual heptachlor per acre.
Cabbage Maggot	Hylemyia brassicae (Bouche)	2-1/4% dust applied to roots and stems before planting gave control. Also plants dipped in a water containing 1 lb. in 50 gallons of water gave control.
		1 lb. actual per 50 gallons of water may be poured along the row at planting time. 4-6 lbs. actual per acre may be applied to the soil and worked in top few inches.
Carpet Beetle	Attagenus sp.	Dilutions of 1 to 1000 have given control as well as giving good residual protection.
Chinch Bug	Blissue leucopterus (Say)	14-1 lb. actual heptachlor per acre has given control.
Clover Root Borer	Hylastinus obscurus (Marsham)	1-1/2 to 1-3/4 lb. actual heptachlor per acre will give very effective control.
Corn Borer (European)	Pyrausta nubilalis (Hubner)	2-3 lbs. of 25% wettable powder per 100 gallons of water has given promising control results.
Corn Earworm	Heliothis obsoleta (Fab.)	0.5 to 0.75% heptachlor in white mineral oil has given good results. 2 or 3 applications at the time the silks appear may be necessary.
Cern Rootworm	Diabrotica sp.	½ to 3 lbs. per acre will give control depending upon type of application, type of soil, and crop protected.
Cotton Thrap	Theips sp.	Apply .08 lb. per acre as a spray or 5 to 8 lbs. of 2-1/2% dust per acre when cotton is small. As the cotton becomes larger, increase the amount of the material used per application.
Cowpea Curculio	Chalcodermus aeneus Boheman	20-35 lbs. of 2-3/5% dust per acre will give control. 2 or 3 applications may be necessary.
Cricket .	Gryllus sp.	4 oz. per acre has given good control where the crickets are attacking crops in the field.
Cucumber Beetle	Diabrotica sp.	20-30 lbs, of 2-35% dust has given control.
Cutworm	Noctuidas	1 lb. of actual heptachlor per acre.
European Chafer	Hoplia sp.	1/2 to 2 lbs, actual heptachlor per acre applied as a soil application.
Eye Gnat	Hippelates pusio (Loew)	2-2-1/2 lbs. actual heptachlor added to the top few inches of the soil.
Flea Beetle	Epitrix sp.	1-1/2 lbs. per acre as soil application shows promise in the control of flea beetle larvae.
Flea	Ctenocephalus sp.	Concentrations as low as .005% have given control.
Garden Wehworm	Loxastege similalis	Apply 0.35 lb. heptachlor per acre. Either sprays or dusts may be used.

insects such as cotton flea hoppers, tarnished plant bugs and rapid plant bugs. Many insects are controlled by susceptibility to both modes of action.

During the past several seasons of testing, heptachlor gave effective control of most of the cotton insect pests and it was especially effective against the cotton boll weevil. Depending upon geographical location, and degree of infestation, 3 to 7 applications per season provide adequate protection to growing cotton. The high percentage of kill of boil weevil in the squares prevented a build-up in the population throughout the season.

Heptachlor has label registration for use in the control of several cotton insects. Insects included in the registration, and the amount of actual heptachlor which is recommended per acre are as follows: cotton boll weevil, cotton flea hopper, tarnished plant bug, and rapid plant bug, for all of which an application of from 0.17 lb. to 0.5 lb. actual heptachlor per acre is recommended; thrips, for which the recommendation is 0.08 lb. per acre;

(Turn to Page 125)

Grasshopper	Locustidae	2 to 4 oz. actual heptachlor per acre as a dust or spray. For bait, add 2½ oz. of heptachlor with 100 lbs. of bait and apply at the rate of 10 lbs. per acre.
Housefly	Musca domestica (Linne)	.05 to 0.1% concentrations are very effective in the control of houseflies036 lb. per acre gave 75% control of houseflies.
Human Lice	Pediculus sp.	0.1% concentration gave complete control.
Japanese Beetle	Popillia Japonica (Newman)	I to 5 lbs, of actual heptachlor per acre applied to the soil has given control. The dosage depends upon the type of soil,
Leaf Miner	Agromyzidae	3 to 5 lbs. of heptachlor per acre have given satisfactory control results.
Lygus Bug	Lygus sp.	1/2 lb. heptachlor in 30 gallons of water per acre gave good control of lygus bugs in alfalfa.
Mormon Cricket	Anabrus simplex Haldeman	3 oz. per acre give good control.
Mosquito	Culicidae	Heptachlor at 1 part in 40,000,000 gave control of larvae. .05 lb. — .075 lb. per acre gave excellent control. 0.1 lb. in oil gave control of salt marsh mosquito. 1.6 — 2 oz. of heptachlor per acre gave good control of adult mosquitoes.
Narcissus Bulb Fly	Merodon equestris	1 lb. heptachlor in 1 or 2% oil solution gave good control when 3 or 4 applications were made.
Onion Maggot	Hylemyia antiqua (Meigen)	1/4 to 1/2 lbs. heptachlor per acre.
Onion Thrip	Thrips tabaci Lindeman	A spray containing ½ lb. heptachlor per acre gives good con- trol. Wettable powder or emulsifiable concentrates may be ap- plied. Also 10-15 lb. of 2-½% dust per acre may be used.
Plum Curculio	Conotrachelus nenuphar (Herbst.)	3-4 lbs. of 25% wettable powder per 100 gallons of water have given control. 2 or 3 applications may be necessary.
Rapid Plant Bug	Adelphocoris rapidus (Say)	Apply ¼ lb. actual heptachlor as a spray or 2-1/2% dust per acre per application.
Roach	Blattidae	Concentrations of 0.1% have given 100% kill within 48 hours.
Screwworm	Cochliomyia sp.	A salve containing 0.01% concentration heptachlor gave good control.
Seed Corn Maggot	Hylemyia cilicrura (Rondani)	2 oz. of 25% wettable powder per 100 lbs, of seed have given good control.
Serpentine Leaf Miner	Agromyza pusilla (Meigen)	35 lbs. of 1-1/2% dust per acre gave excellent control.
Spittle Bug Sugar Beet Root Maggot	Aphrophora sp. Tetanops aldrichi	2 oz. to 3 oz. per acre applied to alfalfa have given good control.
Togar beet Non Maggor	recarolis alunchi	4 oz. of heptachlor per 100 lbs. of seed gives good control. Also 2 lbs. actual per acre applied to the soil has given good control.
Sweet Clover Weevil	Sitona cylindricollis (Fabr.)	4 applications of 0.44 lb. heptachlor per acre gave good control.
Tarnished Plant Bug	Lygus pratensis (Linn)	14-1/2 lb. heptachlor per acre has given good control of tarnished plant bugs in alfalfa.
Tobacco Flea Beetle	Epitrix parvula (Fabr.)	A 5% dust applied to the plant bed prior to transplanting gave very good control.
Tomato Fruitworm	Heliothis sp.	40 lbs. of 2-1/2% dust in early season and 60 lbs. of 2-1/2% dust in late season per acre have given satisfactory control. Also apply
Tuber Flea Beetle	Epitrix sp.	1 lb. per 100 gallons of water at weekly intervals. Soil application of 10 lbs. per acre gave 100% control. No
Turnip Maggot	Hylemyia sp.	lower dosages were used for this pest. 6.3 lbs. of actual heptachlor per acre as a soil treatment gave fair control of this pest.
Western Harvester Ant	Pogonomyrmex Occidentalis (Cresson)	l to 2 tablespoonsful of 25% wettable powder added to the entrance of the nest gives complete control.
White Grub (June Beetle)	Phyllophaga sp.	4 to 10 lbs. per acre gave good control. Three years control has been obtained by adding 10 lbs. heptachlor per acre to the soil.
Wireworm	Elateridae	Apply 1 to 2 lbs. per acre as broadcast treatment and work into the top few inches of the soil. Spray sugar cane cuttings with 1 lb. in 100 gallons of water at planting time. 4 oz. per 100 lbs. of sugar beet seed using 800 lbs. of seed per acre. This uses 32 oz. of heptachlor

per acre.



ARTHUR W. MOHR N. A. C. President

Past, Present and Future of Agricultural Pesticides Trade Viewed at 19th Fall Meeting of

Natl. Agricultural

PREVIEWING the past and contemplating the future in a wellrounded program, the National Agricultural Chemicals Association was to hold its nineteenth annual fall meeting at the Essex & Sussex Hotel, Spring Lake, N. J., September 3-5.

Problems attending the exportation of pesticides were to be viewed by a panel of experts, led by Wallace S. Moreland, Rutgers University, New Brunswick, N. J. This panel was expected to point out the assets and liabilities involved in exporting and to discuss the economics involved.

In addition to Mr. Moreland, the others to appear on the panel, included John H. S. Barr, vice-president, Pennsalt International Corp.; Earl R. Beckner, chief, Chemicals and Semi-Manufactured Products Branch, Manufactured Products Staff, Office of International Materials Policy, U. S. State Department; P. H. Groggins, Chief, Agricultural Chemicals Section, NPA, U. S. Dept. of Commerce; and Dr. Ralph Stewart, Director, Agricultural Products Division, Office of International Trade, U. S. Department of Commerce.

The panel was to mark the finale of the convention, as a climax to three days of serious discussion.

Earlier speakers were scheduled to present papers on both the broad view of agricultural production and the specific materials needed to bring about greater crop yields. Problems of toxicity were also to be considered. Dr. Clair R. Spealman, chief of the Safety Projects Branch, C. A. A., Washington, D. C., was to talk on "Medical Problems in Airplane Spraying and Dusting"; W. A. Minor, assistant to the Secretary of Agriculture, "Food for the Fifth

NACA's 1952

WEDNESDAY, SEPTEMBER 3

Session in charge of Paul Mayfield, vice-president

10:10 a.m. President's Address

Arthur W. Mohr, NAC president

"Medical Problems in Airplane Dusting and Spraying"

Dr. Clair R Spealman, chief, Safety Projects Branch, Civil Aeronautics, Administration, Washington

"Food for the Fifth Plate"

W. A. Minor, Assistant to the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D. C.

11:10 a.m.
"Investment in Progress"

Dr. G. D. Humphreys, president, University of Wyoming

"Agriculture's Role in the American Story"
Dr. Paul R. Sanders, Editor, THE SOUTHERN PLANTER

6:00 p.m. Reception for Members & Guests

Reception for Members & Guests Ladies Invited (Dress optional)

9:30 p.m. Meeting of NACA Board of Directors

THURSDAY, SEPTEMBER 4 10:00 a.m.

Meeting Called to Order Arthur W. Mohr presiding

10:10 a.m.
"Summary of Pesticide Requirements for 1952-53"
Dr. H. H. Shepard, Staff Specialist, Office of Materials & Facilities.
PMA, U. S. Department of Agriculture

10:30 a.m. Announcements

"Agriculture and Technological Improvements"

Oriz V. Wells, Chief, Bureau of Agricultural Economics, U. S.
Dept. of Agriculture, Washington, D. C.

Chem. Assn



LEA S. HITCHNER Exec. Secretary & Treasurer



PAUL MAYFIELD N. A. C. Vice-President

Plate"; Dr. G. D. Humphreys, president of the University of Wyoming "Investment in Progress" and Dr.

Paul R. Sanders, Editor, "The Southern Planter", Richmond, Va., will present his paper.

Fall Program

10:00 a.m

"Conservation Programs and Pesticides"

W. W. Dykstra, Assistant to the Chief, Branch of Predator & Rodent Control, Fish & Wildlife Service, Department of Interior

11:30 a.m. "Merchandising"

Gene Flack, Sales Counsel and Director of Advertising, Sunshine Biscuits, Inc.

7:00 p.m. Banquet for NACA Members & Guests

Ladies Invited (Dress optional)
(The annual golf tournament is held Thursday afternoon. Chairman of this event is John Rodda. Prizes will be awarded.)

FRIDAY, SEPTEMBER 5 Meeting Called to Order

Arthur W. Mohr President

10:05 a.m. Annual Report

Lea S. Hitchner, executive secretary and treasurer, NAC Association

10:20 a.m. "The Bureau of Entomology and Plant Quarantine in 1952" Avery S. Hoyt, chief, B.E.P.Q., U. S. Dept. of Agriculture, Washington, D. C.

> 10:40 a. m. "Meeting the Insecticide Residue Problem"

George C. Decker, Head, Section of Economic Entomology, State Natural History Survey Div., College of Agriculture, University of Ill.

11:00 a.m. Panel on Exports

Wallace S. Moreland, Rutgers University, New Brunswick, N. I.,

presiding. Other members of panel:

John H. S. Barr, Vice-president, Pennsalt International Corporation

Earl R. Beckner, Chief, Chemicals and Semi-manufactured Products

Branch, Manufactured Products Staff, Office of International Materials

Policy, U. S. Department of State

Phillip H. Groggins, Chief, Agricultural Chemicals Section, Chemicals Division, National Production Authority, U. S. Department of Commerce Dr. Raiph Stewart, Director, Agricultural Products Division, Office of International Trade, U. S. Department of Commerce

Pesticide Requirements

PROJECTING known facts into the next season to estimate pesticide requirements is the assignment given Dr. H. H. Shepard, staff specialist, Office of Materials & Facilities, Production and Marketing Administration, U. S. Dept. of Agriculture.


"Agriculture and Technological Improvements", was to be viewed by Oris V. Wells, Chief, Bureau of Agricultural Economics, U. S. Dept. of Agriculture, Washington.

W. W. Dykstra, assistant to the Chief, Branch of Predator & Rodent Control, Fish & Wildlife Service, U. S. Dept. of Interior, was to point out the connections between insecticides, fungicides and rodenticides and conservation. His paper, "Conservation Programs and Pesticides," was scheduled for Thursday.

A talk on "Merchandising" was on Thursday's agenda for presentation by Gene Flack, sales counsel and director of Advertising. Sunshine Biscuit Co.

The annual NAC banquet is scheduled for Thursday evening. No formal speeches were on the program for that time, according to the advance program.

Golf was to be the order of the day for Thursday afternoon, time of the annual tournament. Prizes were to be awarded following the completion of the contests.**



NTERTILIZER, applied scientifically and used with other fruitful farming practices, is a cornerstone of the welfare of the nation. It is the one single method above all others that will permit farmers to meet our bigger future needs. Without more of it, the job cannot be done." This was one of the significant statements contained in a paper "Fertilizer Resources and Requirements of the United States." prepared by the U.S.D.A., the TVA and the U.S. Department of Interior, in the President's Materials Policy Commission report for 1952.

The following is a condensation of the report:

Looking ahead beyond the immediate years and into the future 25 years, the United States has the prospect of providing food and fiber for about 40 million more people than the 150 million figure of 1950.

Should the years through 1975 be peaceful, the U. S. will need much more food and fiber than today because of its growing population. Each year brings from 2 to 2½ million more persons to feed and clothe. By 1975, it is probable that our population will have reached 190 million.

On the other hand, if the next quarter-century is marked with international turmoil, farm production requirements will be even greater.

How will this tremendous production be achieved? New lands for cultivation no longer exist . . . The transition from horses to tractors on farms has provided more than 70 million acres in past years, but the change-over is now practically complete and further acres from this source are relatively negligible. This means that the bulk of future farm production — for the next 25 years or the next 100 — must come from land already under cultivation. Thus, fertilizers become the key to this production.

Fertilizer Ups Production

JUDGING by the amount of feod and clothing required to support the 1950 population of 150 million, it will take about 40 percent more farm production to support 190 million Americans in 1975 at a high standard of living. Under the best conditions, in order to produce enough to meet the national need, farmers will need by 1975, more than 2½ times the 4 million tons of the primary plant nutrients (nitrogen, phosphate, and potash) contained in



the 18.5 million tons of commercial fertilizers which farmers used in the crop year 1950.

The use of fertilizers for supplying plant nutrients to increase farm crop yields produces results. When used under suitable conditions and in conjunction with other desirable practices — provided moisture and other climatic conditions are favorable — fertilizer can bring about a greater increase in production of all crops than any other single influence. It is the key to the tremendous food and fiber production required in the future.

Most crops in all areas of the United States do not get enough needed nutrients from the soil and other natural sources to permit them to grow into their full productive power. Depending upon the type of ground in which they are planted, crops usually lack one or more of

the three primary plant nutrients: nitrogen, phosphate, and potash (N, P₂O₅, and K₂O, respectively). Many also lack secondary nutrients, such as sulfur, calcium, and magnesium and so-called trace elements, like boron, copper, manganese, and zinc. Fertilizing the soil with these nutrients in scientifically-determined proportions provides crops with well-balanced diets.

In order to produce maximum results, the use of fertilizer must be supported by use of other beneficial farm production practices, such as rotation planting, contour plowing, and improved seed along with adequate manpower and machinery and pest control. But the best combination of other practices will not provide the necessary farm production unless enough fertilizer is supplied.

The historical pattern of increased per-acre yields indicates that increased fertilization of crops and improvement in other practices are likely to take place more or less together. If past relationships between increases in farm output and increases in the use of fertilizer are projected toward a further increase of 40 percent in farm output, farmers would use more than 2½ times as much tonnage of plant nutrients as they do at present. It should be noted that a large portion of an increase in fertilizer use will go to maintain present levels of yield alone.

In recent years, farmers have increased their use of commercial plant nutrients at an annual rate of about 10 percent. If this rate is projected forward 20 years, the quantities used by that time would be four times those used now. Even with no improvement in other practices, such a level of use might result in nearly a 75 percent increase in farm output. Even though the response were that great, if it develops that a healthy economic balance would be reached with a 40 percent increase in output, it appears unlikely that the use of fertilizer will reach that level.

Effects on Certain Crops

OUR major crops, each in selected states, corn, cotton, vegetables, and hay and pastures - received nearly 60 percent of the total farm use of the three primary plant nutrients in 1950. They are used as the basis for estimating the total quantities of plant nutrients required and the total production that would result from application of the most profitable rates. Theoretically, if the factors involved in increasing yields were improved commensurate to maximum practicable fertilizer use, current total farm production could be trebled. Obviously, such results are unattainable as national totals, although a few of the more progressive farmers may attain comparable results in special instances.

Corn — a key crop in expanding meat production — is the outstanding example of how farm output can, through fertilizer, be pushed practicably up to the level of future food and fiber requirements. About onefourth of all commercially produced

plant nutrients used on all crops in the United States is applied to the corn crop. And about 40 percent of that is used in the Corn Belt and Lake States. Preliminary estimates indicate that Iowa farmers could practically boost corn production per acre from the current average of 49 bushels to 60 bushels by doubling the amount of plant nutrients they now use and by improving other farming practices about 50 percent. They could increase production to an average of 75 bushels an acre by doubling once again their plant nutrient application and improving their good farming practices by another 50 per-

A similar situation is possible in cotton which uses about 10 percent of all fertilizer applied. Estimates thus far developed show that the average yield of seed cotton in North Carolina, for example, could be increased from the 1949 average of 729 pounds per acre to 1,250 pounds. This would be possible by increasing the amount of plant nutrients used on the crop from the 1949 average of 112 pounds to about 150 pounds together with about a 50 percent improvement in other practices. Another 200 or 300 pounds could be added to the yield if fertilizer application were further increased to approximate more nearly the 50 percent increase in other practices.

About 11 percent of all plant nutrients are applied on hay and pasture in the United States. Estimated response of legume-grass hay and pasture to fertilizer in New York State indicates that the current average yield is little more than one-third of the yield that could be obtained. The yield of these crops could be increased by going half way toward the most profitable rates of application and making about a 50 percent increase in other practices. This would mean an application of about 100 pounds of plant nutrients per acre. Production of vegetable crops which also receive about 11 percent of all primary plant nutrients used, could be increased about a fourth with more fertilizer.

It is emphasized that the foregoing estimates of over-all use of plant nutrients and resulting production at different rates of application are preliminary and subject to substantial revision as continuing research dictates. The large increases in production that would result from much greater use of fertilizer underscores the need for encouragement of balanced production programs. Over a period of 25 years improvements will surely be made in other known practices and more important unforeseen developments will no doubt occur. Therefore, the quantities of fertilizer needed to support estimated production levels 25 years hence will depend not only on the extent of further adoption of other known practices, but also on new developments in farm technology.

Table I

Commercial fertilizer and primary plant nutrients consumed in the United States and Territories in selected years: 1900-1950

			Plant Available	intent Tarel	centent		
Calendar year	Fertilizers (N)		Nitragen phosphoric Potash		Quantity	Proportion	
	1,000 short tons	1,000 abort tons	1,000 short tons	1,000 ekert tone	1,000 short tens	Percent	
1900	2,730	62	246	86	394	14.4	
1910	5,547	146	499	211	856	15.4	
1920	7,296	228	660	258	1,146	15.7	
1930	8,425	377	793	354	1,524	18.1	
1940	8,656	419	912	435	1,766	20.4	
1950	19,758	1,126	2,071	1,215	4,412	22.3	

Source: U. S. Department of Agriculture.

Fertilizer Use Increases

FERTILIZING to encourage bigger and better crops dates back before the earliest colonial settlers arrived. The colonists adopted this method of bettering crops along with other local sources of fertilizer such as animal and plant residues, lime, and marl. In the nineteenth century the foundation was laid for the United States commercial fertilizer industry upon then current fundamental advances made in the science of plant nutrition. The production of commercial fertilizer in the United States grew from about 32,000 short tons in 1859 to nearly 3 million tons in 1899, and on to a new record high of about 19.8 million tons in 1950. This tremendous amount of commercial fertilizer, in addition to manure and lime and other soil-improving materials, played a major role in the near-record 1951 all-crop production volume

During the last 50 years, the manufacturing processes and the characteristics and types of fertilizer materials and mixtures underwent important changes. Natural organic materials, were superseded largely by chemical products, and liquid fertilizers came into use. The primary plant-nutrients in the fertilizer materials and mixtures used in the country have doubled in the case of certain materials. Marked improvement was made in the physical condition of the materials, in mixtures, and in their packaging.

The United States has become largely self-sufficient in the supply of the three primary plant nutrients, nitrogen, phosphate, and potash. The country now leads in production of nitrogen and phosphate fertilizers. Production of potash is exceeded only by that of Germany.

The 1950 consumption of fertilizer was 128 percent higher than in 1940 and over 7 times the consumption in 1900, while the total use of the 3 primary nutrients in 1950 was 150 percent more than in 1940 and over 11 times the use at the beginning of the century (see table I).

The record 1950 use of commercial fertilizer was the twelfth consecutive year of increase. The production of nitrogen and potash each more than doubled in the last 7 years, while production of phosphate doubled in the last 10 years. Even so, supplies were short of demand in most of the years. Great progress also has been made in the use of soil-liming materials. The total consumption of liming materials in 1950 was 26.5 million tons, or more than 8 times that of 1935. (See table II.)

A large tonnage of liming materials is required to maintain a satisfactory status of lands that have been limed. Holding existing gains and making further improvement of the lime status of agricultural soils in this country would require an annual application of nearly 80 million tons of liming materials. But this level will not be attained without extraordinary educational crop production campaigns.

At the turn of the century, little or nothing was known of the plant requirements for the trace elements, but the need for applying small quantities of these elements to various crops in many parts of the country is now being recognized.

The greatly accelerated increase in consumption of fertilizer and plant nutrients in recent years has been influenced by many factors. These include: (1) the urgent need for additional food and fiber to meet domestic and export needs, (2) the greatly improved economic position of the American farmer, (3) the continued decline in the native fertility of large areas of the nation's soils, and (4) a more general recognition of the potentialities of fertilizer in lowering production costs and increasing cropyields.

Table II

Consumption of agricultural liming materials on farms of the continental United States in selected years: 1930-50

Calendar year: 1930	1,000 short tons 3,498
1935	3,292
1940	13,434
1945	22,357
1950	26,536

Sources: National Lime Association and

Table III

Distribution of commercial-fertilizer consumption in the continental

United States by principal crops 1929, 1942, 1949, 1950

	Percent	of total fertilizer con	sumption
Сгер	1929	1942	19501
Corn	20.7	22.0	24.8
Cotton	28.0	14.6	9.0
Small grains	13.8	14.1	18.4
Tobacco	6.8	5.3	4.1
Potatoes	8.5	7.1	4.2
Hay and pasture	2.1	12.9	12.4
Vegetables ²	6.7	10.4	7.6
Fruits	4.3	6.7	4.2
Other crops	9.1	6.9	15.3

'Your ended June 30.

Sources: National Fertilizer Association and U. S. Department of Agriculture.

While it is not possible to make trustworthy estimates of the extent of the gap between fertilizer supply and quantities farmers would buy at prevailing prices, it is known that annual demand was not met during the last 10 years. Major reasons for failure to meet demand include: (1) lack of sufficient manufacturing and processing facilities and the difficulty of expanding existing facilities, (2) necessity for shipping large quantities of plant nutrients to allies and to occupied countries, and (3) shortages of some chemicals essential for fertilizer manufacture.

The greater proportion of the total annual consumption of commercial fertilizer in the United States is in the form of mixtures — ranging from 67 to 70 percent of the total during the last 8 years. Of the primary nutrients, more than 90 percent of the potash, about 70 percent of the phosphate, and 50 to 60 percent of the nitrogen applied as commercial fertilizer are used in mixtures.

Use Varies in Sections

As one would expect, the consumption of fertilizer varies considerably among the different regions of the country. For many years the South Atlantic and Gulf States, which generally are low in native soil fertility, have accounted for a large portion of the annual consumption. Consumption has greatly increased, however, in other regions, notably the North Central States where soil fer-

tility originally was much higher than in the southerly regions.

In 1933-34 the South Atlantic region used 50 percent of the fertilizer nitrogen used in the United States, while the North Central region used only 4 percent. In 1949-50, however, the respective figures were 29 and 15 percent. The growing need for nitrogen in the North Central region is indicated by the fact that the consumption of this nutrient in 1949-50 was 14 times the consumption in 1933-34. The proportionate increase in the Western region was even larger.

The South Atlantic region also has yielded to the North Central region as the leader in potash consumption. The consumption in the latter region in 1949-50 was about 12.5 times that of 1933-34.

For many years fertilizers were used chiefly on such cash crops as cotton, tobacco, potatoes, and vegetables. The need for a better balance in the production of food, feed, and fiber has been a major factor in alternating the distribution of fertilizer consumption by crops. (See table III.)

The long-range shift in production is toward more grassland farming, a development supported by the United States Department of Agriculture and the Land Grant Colleges. The continuing grassland program is designed to make better use of soil resources and to improve their management so that each acre may contribute more fully to the total

Agricultural Limestone Institute.

agricultural production while building the soil for future output. Future trends with reference to grassland farming will have an influence on total and regional use of fertilizer, and on the relative quantities of the different plant nutrients needed.

Fertilizer Expenditures

OMPARATIVE responses in yields and relative costs of the different combinations determine the farmer's final choice, provided he is well-informed on these matters. While much progress has been made, too many farmers are still not aware of advantages to be gained from increased use of fertilizer.

In 1949, farmers spent nearly 700 million dollars for fertilizer; 3.9 percent of the total agricultural production expense of farm operators in that year. In the past 38 years farmers spent for fertilizer an average proportion of 3.4 percent — ranging from 2.5 percent to 4.8 percent—of their total annual production expense.

The 700 million dollars farmers spent for fertilizer in 1949 was 4.4 times as much as they spent in 1911. During the same period, their total production expenses increased 4.9 times.

The proportion of the farmer's income spent for fertilizer varies greatly. It is generally much higher in the older fertilizer consuming States than in the newer areas of use, however.

Fertilizer Materials

FUTURE demands on raw materials for fertilizer production are so great that we must seek better methods of using our known resources and probe for additional sources as well.

The only fertilizer raw material that is inexhaustible is nitrogen, which exists in the atmosphere. While other sources of nitrogen for fertilizer exist — the fixation of nitrogen synthetically from the atmosphere is the most important source, furnishing more than 65 percent of the annual consumption. This source of nitrogen is limited only by available production facilities required in fixation. Facilities were inadequate to meet 1952 demand, but the long-term trend looks optimistic.

The United States produces nearly half of the world output of phosphate rock, and the reserve is exceeded only by that of French Morocco. In the calendar year 1949, phosphate rock supplied more than 98 percent of the 1,884,000 short tons of the phosphate fertilizer used by farmers.

The present economically minable reserve of phosphate rock in the United States totals about 4 billion long tons; potentially minable reserves total at least 9 billion tons.

By virtue of the development of extensive potash deposits in New Mexico, this country has been substantially self-sufficient in supplies of this essential plant nutrient since 1940. The domestic use of potash is predominantly for fertilizer; 93 percent is used for fertilizer, 7 percent for chemicals.

The Nation's proved reserves of potash economically recoverable under present conditions are located in the New Mexico deposits and in the brines of Searles Lake in California and Salduro Marsh in Utah. While current estimates of minable potash indicate reserves of more than 200 million tons, future fertilizer demands for this material require a broad search for and development of potash resources and research on utilization of lower grade materials. The United States holds large resources of certain secondary fertilizer materials.

Facilities Must Expand

EXPANSION of present manufacturing and processing facilities is vital to make plant nutrients available at increasing rates. Fuller use must be made of existing facilities. In the case of the primary nutrients, generally the capacity for producing these has not kept pace fully with the demand.

A major problem is to increase nutrient-producing capacity to meet present demand and to insure continued expansion as needed in the future.

Despite progress in manufacturing processes and techniques, there is room for great improvement. Research and development in this field must be expanded and accelerated.

Future efforts should be directed toward production of a larger portion of the nation's fertilizer nitrogen requirements in the form of highanalysis, solid products, such as ammonium phosphate and especially urea, the most concentrated of the solid nitrogen fertilizer materials.

Lack of adequate supplies of concentrated phosphates is a major obstacle to expanded production of high-analysis fertilizers.

More than 90 percent of the available phosphates used as fertilizer is produced by processes involving treatment of phosphate rock with sulfuric acid. While plentiful supplies of sulfur have been available in recent years, a current serious shortage must be overcome to support increased phosphate fertilizer production. This can be done by discovery of new sources of sulfur, but perhaps more practically by developing known methods of producing phosphate fertilizers without the use of sulfuric acid, or at least with less of it. Use of nitric acid for treatment of phosphate rock should be developed as one of these methods.

Future increases in the use of fertilizer can be mostly independent of foreign sources of supply. A continuing increase in the domestic manufacture of plant nutrients will help guarantee the nation plentiful food and fiber in the future.

Out of the total domestic fertilizer nitrogen supply of 1,387,00 short tons in 1950-51, domestic sources supplied 1,098,000 tons, while imports totaled 289,000 tons. During the same period, the United States exported 102,000 tons. Expansion in nitrogen fertilizer production, especially by the synthetic ammonia process can provide the United States with required supplies.

The United States traditionally exports much more phosphate rock than it imports. In 1950, exports totaled 1,832,000 long tons, compared with imports of 87,000 tons. During that year, the United States used about 8,509,000 tons of

(Turn to Page 141)

Residues of

Dithiocarbamate Fungicides

-on food crops

DURING the 1930's an interesting pattern of biological activity was revealed through investigations with derivatives of dithiocarbamic acid by Tisdale and Flenner⁸. Since then, the fungicidal properties of the metal salts of dithiocarbamic acid have achieved outstanding importance, and there are in wide use four derivatives with the twin advantages of control of a great variety of plant diseases and low phytotoxicity. These compounds are:

Ferric dimethyldithiocarbamate (assigned generic name: ferbam)

$$\begin{array}{c|c}
CH_3 & N - C - S \\
CH_3 & "
\end{array}$$
Fe

Zinc dimethyldithiocarbamate (assigned generic name: ziram)

Disodium ethylenebisdithiocarbamate (assigned generic name: nabam)

$$\begin{array}{c|c} CH_2 - N - C - S \\ & H & ") \\ CH_2 - N - C - S \\ & H & ") \\ \end{array}$$
 Na₂

Zinc ethylenebisdithiocarbamate (assigned generic name: zineb)

$$\begin{array}{c|c}
CH_2 - N - C - S \\
 & H & " \\
S \\
CH_2 - N - C - S \\
 & H & " \\
S \\
\end{array}$$
Zn

All the compounds listed, except nabam, are essentially water-insoluble, and are used as dusts or sprayed as wettable powders. The soluble dithiocarbamates, including nabam, are quite phytotoxic. Nabam is not applied as such to crops, but is combined with zinc sulfate to form a tank-mix precipitate of the insoluble zinc salt or zincb.

Recently, a new member of the dithiocarbamate fungicide family has reached the market. This product is:

C. J. Krister

E. I. duPont de Nemours & Co., Inc. Grasselli Chemicals Dept. Wilmington. Delaware Manganese ethylenebisdithiocarbamate (abbreviation: MnEBD)

$$\begin{array}{c|c}
CH_2 - N - C - S \\
H & ") \\
CH_2 - N - C - S \\
H & ")
\end{array}$$
Mn

Like zineb, the manganese salt is only slightly soluble in water. It is available as "Manzate" fungicide.

Uses for Manzate

I NITIAL recommendations for "Manzate" fungicide are to control early blight and late blight of potatoes and the five major fruit and foliage diseases of tomatoes. Additional research is being conducted on other promising applications, including diseases of celery, carrots, cucurbits, cabbage, onions, spinach, apples, peaches and raspberries.

The extensive commercial applications of the older dithiocarbamate fungicides are evidenced by many hundreds of favorable references during the Food and Drug Administration's Residue Tolerance Hearings⁸ in 1950. Their usefulness was outlined in part by Tisdale⁵ as follows:

"Ferham is especially useful in the field of fruit disease control. For the control of apple rust and quince rust it has no equal. Under many

^{*}Trademark of E. I. du Pont de Nemours & Co. (Inc.) for its fungicide based on manganese ethylenebisdithiocarbamate.

Table 1

Dithiocarbamate Residue	Results	Classifie	d	Ву	Range	
Range in Parts per Million		Percent	of	All	Residue	Resul

pe in Parts per Million 0—1 p.p.m.		Percent of All Residue Resu 64.9%
1-5	*	23.4
5-10		8.2
Over 10		3.9

Table 2

Effect	of Number	of Treatments	On Dithiocarbo	amate Residue	Levels
Active Ingredient	Crop	Location	Elapsed Time* (Days)	No. of Treatments	Residues (p.p.m.)
Zineb	Cabbage	Ohio	21	6	1.2, 0.9
			21	1	1.8, 1.5, 0.6
Zinch	Tomatoes	Ohio	21	6	0.15 0.15

				0	0.17. 0.
					0.22
			21	1	0.13, 0.
Zineb	Tomatoes	Florida	8	19	0.7
			5	8	0.6

*Between final application and harvest

conditions it is a superior product for the control of apple scab, especially on the fruit. Pear scab is very effectively controlled with ferbam. Ferbam is considered a superior product for the control of brown rot and leaf spot of sweet cherries. (It) is the preferred treatment for several important diseases of small fruits.

"Ziram has found its greatest uses in the field of vegetable crop disease control. Outstanding uses are the control of tomato anthracnose and the early blight of tomato and potato. It is effective for the control of such cucurbit diseases as anthracnose and downy mildew.

"The outstanding use of zineb and the nabam-zinc sulfate combination is for the control of late blight of potatoes and tomatoes. It also controls early blight and other leaf infections of these crops. The early and late blights of celery also are effectively controlled. Other diseases controlled with zineb and nabam are downy mildew and white rust of spinach, rust of beans and onion purple blotch."

Rate of Application

THE rate and frequency of application of these fungicides varies with the crop, the plant disease, the product, and local conditions. Several applications are usually made per season at intervals of five to fourteen days.

Concentrations generally vary between 1 and 3 pounds of wettable powder formulation (65 to 76% active ingredient) per 100 gallons of spray mixture, or 5 to 15% dusts. Nabam is normally used at a rate of 2 quarts (19% nabam) per 100 gallons of spray along with 3/4 lb. of monohydrated zinc sulfate or 1 lb. of flake zinc sulfate.

Sprays are applied to tree fruits at gallonage rates sufficient for adequate coverage and to truck crops at 100-200 gallons per acre, while dusts are normally used at 30 to 50 pounds per acre. Resulting rates of application of active ingredients, using truck crops as an example, are in the order of 1.5 to 7.5 pounds per acre for a single treatment.

Residue Data

RESEARCH has been conducted by the du Pont Company and other agencies to determine crop residues of ferbam, ziram and zineb¹, as well as residue data for the newer MnEBD.

Results of residue studies on eight fresh fruits, eight fresh vegetables and three canned foods indicate that: (1) initial deposits are relatively low; (2) residues weather rapidly; (3) final residues are relatively independent of the number of treatments or rate of application; (4) residues are reduced by washing or processing; and (5) harvest-time residues are generally very low, even where commercial spray schedules involving repeated applications were employed.

Ninety-six percent of all results are less than 10 p.p.m. (parts of the dithiocarbamate fungicide per million parts by weight of treated fruit or vegetable), as shown in Table 1. Almost 260 individual results, excluding replicated determinations, have been obtained.

The number of treatments evidently has little effect on the harvest residue levels. This is shown in the, Table 2 above cited from a paper by Lowen⁴.

The evidence that final residue is relatively independent of the number of treatments or rate of application is apparently due to the rather rapid degradation of dithiocarbamate deposits by ordinary processes of weathering. This reduction in residue level is demonstrated in Figure 1. It may be noted that all residues in excess of 5 p.p.m. appeared on samples taken within seven days following the last application.

The type of surface and the ratio of surface area to mass of the fruit or vegetable are factors influencing the magnitude of residues. As expected, residues are lower on smooth-surfaced, bulky fruits and vegetables than on fruits and vegetables which are fuzzy, leafy or small in size. For example, fresh green beans and celery leaves have exhibited the highest residual deposits in these studies.

The maximum residue found on tomatoes treated with "Manzate" fungicide is 1.3 p.p.m. MnEBD. For peppers - smooth surfaced, but less bulky than tomatoes - residues immediately following applications have varied between 3.2 and 5.2 p.p.m. MnEBD, but these levels declined to less than 1 p.p.m. within a few days. The relatively high residues on celery are reduced by stripping, trimming and washing. Pascal celery showed 8.0 p.p.m. MnEBD and Golden variety 6.3 p.p.m. upon stripping and trimming; both varieties had been sampled one week after the last of 12-15 treatments with "Manzate" fungicide.

Washing these same celery samples by conventional commercial procedures reduced the residue to 1.75 p.p.m. for the Pascal and 2.80 p.p.m. for the Golden variety. To date, "Manzate" fungicide has demonstrated notably promising control of Cercospora and Septoria blights of celery and tests are continuing.

Considerable residue work on "Manzate" fungicide has been conducted with tomatoes, since this product appears to be an excellent single fungicide for control of the major foliage and fruit diseases of tomatoearly blight and late blight, anthracnose, gray leaf spot (stemphylium), and Septoria leaf spot. The results of these studies shown in Table 3, demonstrate the low residue levels even for samples taken immediately after treatment.

Washing experiments on several crops indicate that dithiocarbamate fungicide residues are reduced, but not eliminated by this operation. Higher residues were more greatly reduced percentagewise than were lower levels. Spot-checks on canned fruits and vegetables to ascertain the effect of processing and canning revealed no residue in excess of 0.4 p.p.m.

A selected list of residue data for the five dithiocarbamate fungicides on ten different crops is presented in Table 4 in order to portray actual harvest-time residues resulting from conventional protective fungicide treatments. In all cases, the data are for samples subjected to more than

one application, at rates used according to recommended grower practice. Intervals between the last application and sampling or harvest also depended upon conventional procedure, with a full month interval in the case of cherries, while ranges in tomato and pepper results represent the differences between samples at 6-7 days and those at 0 days.

For the important uses of dithiocarbamate fungicides listed in Table 4, it is indicated that crop residues of these materials at harvest are low in magnitude.

Sampling Procedures

THE attempt was generally made to obtain at least a 5-lb. random sample of the treated crop (or a random composite of replicated treatments); also an untreated sample for use as a control, or in any special analytical studies required for that crop.

These samples were usually shipped in paper grocery bags as the inner containers along with empty bags for use as blanks in the determination. Shipments by air express and refrigeration on receipt of the sample

Table 3
Residue Levels, "Manzate" Fungicide on Tomatoes

Interval Between Last Appln. & Sampling	Interval Between Sampling and Analysis	Desage lbs. per 100 Gals.	Rate of Appln. Gal. per Acre	No. of Apples.	Residue Level as P.P.M. of MnEBD	Location
15 min.	2 days	2	150	1	1.3	Texas
30 min.	7 days	2	150	1	0.5	Florida
1 hour	4 days	2	150	1	0.6	Florida
1 hour	4 days	2	250	1	0.8	Florida
3 hours	4 days	2	200	1	1.05	Florida
7 days	4 days	2	200	1	0.09	Florida
6 days	2 days	11/2	150	7	0.07	Delaware
5 days	5 days	11/2	170	16	0.3	Florida
"0 days"	5 days	11/2	170	17	0.4	Florida
7 days	7 days	11/2	170	19	0.08	Florida
"0 days"	7 days	11/2	170	20	0.4	Florida

Table 4
Harvest-Time Crop Residues of Dithiocarbamate Fungicide

Crop	Fungicide	Residue in P.P.M.
Apples	Ferbam	0.05
Cherries	Ferbam	3.6 (Fresh)
		0.09 (Canned)
Grapes	Ferbam	0.25
Cucumbers	Ziram	0.45
Celery	Zineb (from Nabam)	1.45 (unwashed)
		0.72 (washed)
Potatoes	Zineb (from Nabam)	0.0
Tomatoes	Zineb (from Nabam)	0.7 to 2.1 (unwashed)
		0.15 to 0.6 (washed)
Cabbage	Zineb	0.14 (head wt. basis)
		1.05 (leaf wt basis)
Cucumbers	Zineb	0.1
Celery	MnEBD	1.75 to 2.80 (stripped,
		trimmed & washed)
Peppers	MnEBD	0.16 to 5.2 (unwashed)
		0.0 to 0.09 (washed)
Squash	MnEBD	0.5 (unwashed)
		0.07 (washed)
Tomatoes	MnEBD	0.07 to 1.3

have been the rule. Although it has not always been possible, the goal has been to avoid excessive hold-up of samples before analysis, through scheduling of shipments and availability of analytical manpower. Results of this concerted effort are evident in Table 3, wherein the maximum delay between sampling in the field analysis in the laboratory for MnEBD residues on tomatoes has been 7 days.

Different techniques of handling samples for analysis were found desirable under varying circumstances. For foods with smooth surfaces and low surface-volume ratios, such as apples, the residue was removed from the sample with a solvent, and the residue then isolated for analysis by evaporation of the solvent. It was found necessary to macerate leafy vegetables such as cabbage and spin-ach into a homogeneous pulp and conduct the analysis directly in this medium. In certain other instances, especially with zineb and MnEBD (because of limited solubility in suitable solvents), the residue was removed from the sample by a water wash containing a small amount of wetting agent; the washings were then analyzed directly.


The method described by Lowen's was primarily used in analyzing the treated crops for ferbam, ziram, zineb, (zineb is the active principle of nabam since nabam is used with zinc sulfate) and MnEBD. In this method, which is a modification

of the Dickinson-Viles² procedure, the active ingredients decompose to carbon disulfide upon treatment with acid and heating. The carbon disulfide is absorbed in a copper acetate—dicthylamine reagent and the resulting copper diethyl dithiocarbamate is determined colorimetrically.

Acknowledgment. Grateful acknowledgment is made to the many state and federal investigators who cooperated in providing samples, and to W. K. Lowen, H. M. Baker, field agricultural research and development personnel, and D. A. Carlson, all of du Pont, for analytical work, provision of samples, and compilation of residue data.

Figure 1

THE EFFECT OF TIME INTERVAL BETWEEN LAST APPLICATION & SAMPLING ON DITHIOCARBAMATE FUNGICIDE RESIDUE LEVELS



* PERCENTAGES OF THE 226 RESULTS WHICH WERE CLASSIFIABLE AS TO TIME INTERVAL.

Literature Cited

- Anon., "Residues of Dithiocarbamate Fungicides on Food Crops," Mimeo., E. I. du Pont de Nemours & Co., Inc., July 1, 1951.
- (2) Anon., Am. Phytopath. Soc., "The Present Status of Chemicals Used to Control Diseases of Edible Fruits and Vegetables in the United States," U. S. Food & Drug Admin., Tolerance Hearings, FDC-57, 1950, Exhibit 787.
- (3) Dragt, G., "Method of Analysis for Fungicides in Crop Residues (Dickinson-Viles' Method)", U. S. Food & Drug Admin. Tolerance Hearings, FDC-57, 1950, Exhibit 1012, Part E.
- (4) Lowen, W. K., Anal. Chem., 23, 1846-50 (1951), Dec.
- (5) Tisdale, W. H., "Derivatives of Dithiocarbamic Acid as Fungicides," U. S. Food & Drug Admin. Tolerance Hearings, FDC-57, 1930, R-4105.
- (6) Tisdale, W. H. & A. L. Flenner, Ind. & Eng. Chem., 34, 501-2 (1942) April.

Of particular importance at present, is study of toxic residues on food crops. How to determine the amount of fungicial residue is viewed as fundamental

AGRICULTURAL CHEMICALS

Cary, Hart, Vernon Upped in Food Machinery Corp. Shifts

THE Board of Directors of Food Machinery & Chemical Corporation has announced important changes in the firm's corporate management. J. B. Cary, San Jose, California, formerly executive vice-president, was elected chairman of the Executive Committee of the corporation and will share with the president, Paul L. Davies, responsibilities of overall supervision.

In 1947 Mr. Cary, then president of Niagara Chemical Division, Middleport, N. Y., was made executive vice-president of Food Machinery and Chemical Corporation and moved to San Jose. At that time, Ernest Hart assumed the duties of president of Niagara, which position he has held to this date. Mr. Hart was elected executive vice-president of the corporation and will be responsible for the supervision of operations of its chemical divisions. These include Niagara Chemical Division, Middleport, New York; Westvaco Chemical Division, New York City: Ohio-Apex Division, Nitro, West Virginia; and Buffalo Electro-Chemical Company of Buffalo, New York. Mr. Hart will make his headquarters in the new Chrysler Building East in New York

City. However, he will continue to maintain his residence in Medina, New York- Both Mr. Cary and Mr. Hart are past presidents of the National Agicultural Chemicals Association.

Benjamin C. Carter, a former vice-president and controller of the corporation, was elected executive vice-president in charge of the mechanical divisions of Food Machinery and Chemical Corporation and will continue his headquarters in San Jose, California.

Succeeding Mr. Hart as president of Niagara Chemical Division is Jackson V. Vernon, who has been Division vice-president and sales manager since 1947. Mr. Vernon is a graduate of Mississippi State College. After serving a short time with the Mississippi State Plant Board and the U. S. Department of Entomology, he joined the Niagara sales organization in 1923 as assistant territory manager, operating in the Memphis, Tennessee area. In 1925, he was transferred to the midwest area to develop the company's business in that section on fruits and vegetables. In 1933, he became district sales man-





IOSEPH B. CARY
New Chairman of Food Machinery &
Chemical Corp. Executive Committee



ERNEST HART

Named executive vice-president Food

Machinery & Chemical Corp.

B. EARLE VOSTEEN
Controller & vice-president,
Niagara Chemical Div.



SEPTEMBER, 1952

STUART H. BEAR
Vice-president and sales manager
Niagara Chemical Div.



JACKSON V. VERNON

Named president of Niagara Chemical
Division, succeeding Mr. Hart



Entire industry invited to participate in October

Fertilizer Safety

CTOBER 22-23 are the dates set aside for the second annual meeting of the Fertilizer Section of the National Safety Congress to be held at the Sheraton Hotel, Chicago, Illinois. According to the advance schedule, a program of unusual interest has been planned, with an urgent invitation extended to the entire fertilizer industry.

General chairman J. S. Fields, safety director, Phillips Chemical Co., Bartlesville, Okla. is to open the fertilizer safety section of the congress with a talk outlining the general objectives of the body, and a report on progress made during the past year.

The subject of multiple shot blasting in fertilizer storage will be discussed by Mark Withey, explosives expert, Trojan Powder Co. He is expected to answer questions from the floor regarding the safe practices of blasting and to explain techniques involved in the "delayed action" charges now being used in fertilizer plants to reduce hazards from powerful single-shot blasts.

J. L. Rosenstein, head of the Department of Psychology, Miami University, will discuss that phase of safety, describing the employees' mental processes and attitudes which enter the safety picture. His title will be, "Why Safety."

How the proper designing of

fertilizer plants may aid fire prevention will be described by G. G. Blair, fire prevention engineer of Ebasco Services, Inc. Following his talk, the meeting will be thrown open for questions and further discussion-

A practical demonstration of how multiple shot blasting aids the manufacturer in breaking up safely a hardened pile of material, is scheduled for Saturday morning, October 23. Mr. Withey will set off an actual blast at the Chicago Heights plant of International Minerals & Chemical Corporation. The spectators will be taken to the scene in special buses, according to current plans.

In the afternoon, Mr. Fields will preside at an election of officers, followed by a talk by John E. Smith, safety director, Spencer Chemical Co., Kansas City, Mo., on "1953 Goals for the Fertilizer Section."

Having returned to the hotel after witnessing the multiple shot blasting at Chicago Heights, the

JOHN SMITH
To speak on 1953 goals for fertilizer safety



AGRICULTURAL CHEMICALS

Conference Sheraton Hotel, Chicago. Safety leaders anticipate crowd larger than that of 1951.

Fertilizer Safety Section of the National Safety Congress plans big two-day event October 22 & 23 with headquarters at the Sheraton Hotel, Chicago. Safety leaders anticipate crowd larger than that of 1951.

crowd will discuss with Mr. Withey the methods he uses and further questions regarding his technique.

E. O. Burroughs, Jr., manager of the Insurance Department of F. S. Royster Guano Co., Norfolk, Va., will speak on "Housekeeping in Fertilizer Plants," pointing out how necessary it is to maintain neatness and orderliness in a plant if maximum safety is to be realized.

The hazards encountered from

gas and dust in fertilizer plants will be discussed by Herbert Walworth, Lumberman Mutual Casualty Co. How to control both dust and gas in a plant will be the topic of his talk Saturday afternoon.

Thomas J. Clarke, Personnel Manager, GLF Exchange, Ithaca, N. Y., will talk on the importance of safety meetings in a plant, under the general subject of "How to Conduct a Safety Meeting." Mr. Blair is scheduled to present a talk on "Fire Prevention in Fertilizer Plants Through Maintenance" as the final item on the afternoon's program.

Last year's program in Chicago attracted several times the number of persons expected to be in attendance and the program committee for 1952 is anticipating a large crowd. Representatives of every fertilizer plant are urged to be present.

A. B. PETTIT
Behind-the-scenes booster for
fertilizer plant safety



SEPTEMBER, 1952

JACK FIELDS
General chairman of fertilizer safety section



E. O. BURROUGHS, JR.

To explain importance of good housekeeping practices



A. O. A. C., Feed, Fertilizer and Economic Poisons

Control Officials

To hold annual meetings in Washington, Sept. 29-Oct. 4

OUR important groups of agricultural chemical control of ficials will hold a week-long series of meetings at the Shoreham Hotel, Washington, D. C., September 29 through October 4. The participating groups are, the Association of Economic Poisons Control Officials; the Association of Official Agricultural Chemists; Association of American Feed Control Officials; and the American Association of Fertilizer Control Officials.

AOAC Leads Off

THE AOAC will hold a three-day session beginning September 29, according to K. L. Milstead, secretary-treasurer of the Association. Details of the program were not complete at press time, but Mr. Milstead said that papers will be pre-

sented on the application of flame photometry and on the determination of potassium in fertilizers and sodium in foods. Dr. C. H. Kunsman and his colleagues, Western Regional Research Laboratory, U. S. Department of Agriculture, will also present a paper, "Moisture Determination of Foods by Hydrogen Nuclei Magnetic Resonance."

Henry A. Lepper, AOAC president, will present his address on Monday afternoon, September 29; and the next afternoon, Dr. L. A. Maynard, Cornell University School of Nutrition, will speak on "The Import of Recent Advances in Nutrition on Food and Feed Control Problems."

The AOAC banquet will be held Tuesday evening, September 30, with an address by C. W. Dunn, New York City food attorney, on "Some Observations on the Food and Drug Law."

Feed Control Officials

R EPORTS of Committees comprise the bulk of the two-day (October 1 € 2) meeting of the Feed Control Officials. Dr. L. E. Bopst, secretary-treasurer of the Association will present his annual report on October 1, followed by a business session presided over by president F. W. Quackenbush, Purdue University, Lafayette, Ind.

Dr. Quackenbush will call on committee chairmen and investigators to give their reports, with a total of thirty-four scheduled to appear with various kinds of information gathered during the past year.

According to Dr. Bopst, Tuesday's program was not complete at press time, but some of the speakers had confirmed invitations to appear. Dr. C. R. Grau, Division of Poultry Husbandry, University of California, Berkeley, will speak on "Some Aspects of Protein Nutrition;" and C. W. Sievert, consultant to the feed industry, Chicago, Ill., will present a paper, "The Feed Consultant and the Control Official."

Others tentatively named on the advance program included a representative of the National Fisheries Institute, Washington, D. C. and Ray Bowden, Grain & Feed Dealers National Association, Washington, D. C.

Tuesday afternoon's session will be devoted to reports of the executive, credentials, auditing and resolutions committees, elections of officers; and a session for feed control officials only, wherein matters pertaining to new definitions and regulations will be discussed.

Fertilizer Group

HE annual meeting of the Association of American Fertilizer Control Officials is scheduled to be held at the Shoreham Hotel, Washington, D. C., October 3, according to Dr. Bruce D. Cloaninger, Clemson, S. C., secretary-treasurer of the Association. The meeting will be held in connection with Conventions of the Association of Economic Poison Control Officials: the American Association of Feed Control Officials: and the Association of Official Agricultural Chemists, all of which are also meeting at the Shoreham during the early days of October.

Dr. J. F. Fudge. College Station, Texas, president of the A.A.F.-C.O., will address the Association during the morning program, followed by Dr. Russell Coleman, president, the National Fertilizer Association, Washington, D. C.

Paul T. Truitt, president, American Plant Food Council, Washington, D. C., will speak on "Prospective Promotions for Plant Food" and Dr. Edwin C. Kapusta, chemical engineer, National Fertilizer Association, Washington, D. C., will discuss "Some Manufacturing Problems and New Developments in Fertilizer Technology."

"Polyelectrolyte Soil Amendments" will be the subject of a talk by Alien B. Lemmon, chief, Bureau of Chemistry, Sacramento, California, who is to follow Dr. Kapusta on the program.

To complete the morning's program, Dr. H. J. Fisher, New Haven, Conn., will present the report of the States Relations Committee.

Members of the executive committee will gather for a luncheon session at noon and in the afternoon. Rodney C. Berry and M. B. Rowe, both of the State Department of Agriculture, Richmond, Va., will discuss the distribution of bulk fertilizers. They will be followed by Stacy B. Randle, New Brunswick, N. J., who will report on the model state fertilizer bill. G. W. Michael, Ottawa, Ontario, Canada, will present the report of the executive committee.

The following will also present reports: M. H. Snyder, Charleston, Va.; M. P. Etheridge, State College, Miss.; J. W. Kuzmeski, Amherst, Mass.; Dr. Fudge; R. W. Ludwick, State College, N. M.; W. B. Griem, Madison, Wis.; J. B. Smith, Kingston, R. I.; Rodney C. Berry, Richmond, Va.; Gordon Hart, Tallahassee, Fla.; E. W. Constable, Raleigh, N. C.; John L. Monaghan, Topeka, Kans.; Bruce Poundstone, Lexington, Ky.; F. W. Quackenbush, Lafayette, Ind.; G. H. Laramie, Concord, N. H.; Allen B. Lemmon; E. A. Epps, Jr.,

Baton Rouge, La.; and M. B. Rowe, Richmond, Va.

Reports of the auditing, resolutions and nominating committees, plus election of officers, will conclude the formal portion of the program.

A presentation in the form of six plaques is planned to be made to the past presidents of the Association, according to Dr. Cloaninger.

AEPCO Ends Meeting

COMPLETING the week's series, the A.E.P.C.O. plans an all-day session at the Shoreham. According to Dr. A. B. Heagy, College Park, Md., secretary-treasurer of the group, a round table discussion is to cover details of regulatory activity, to benefit members unable to attend the meeting of the executive committee earlier. Members of the panel had not been selected at press time.

Dr. E. W. Constable, State Chemist, Raleigh, N. C., president of the A.E.P.C.O., will address the assembly on Saturday morning. Others slated to appear on the program include Dr. H. L. Haller, assistant chief, Bureau of Entomology and Plant Quarantine, U. S. Department of Agriculture, Washington, on "Newer Materials as Pesticides."

M. F. Crass, Jr., secretary of the Manufacturing Chemists' Association, will speak on "The Labeling of Pesticides" and Dr. A. J. Lehman, chief, Division of Pharmacology, Food and Drug Administration, Washington, D. C., will discuss "Toxicity of Newer Pesticide Materials."

Four groups of control officials to consider problems of law enforcement in every state. New pesticides, new uses for fertilizers and labeling of soil conditioners all pose important problems.

Crush 9t! Grind 9t Separate 9t





Do It Better,



WILLIAMS


HAMMER MILLS





HELIX-SEAL HAMMER MILLS

For cleaner, safer dustless grinding of dry materials—or non-clogging, accurate grinding of wet, sticky or greasy materials—to finenesses of 100 to 325 mesh. Combines the advantages of a hammer mill with closed-circuit screw-type feed. No separators, fans or cyclones required. Available with steam or water jackets. Low investment—inexpensive to install and operate.



Hammer Mill!

ROLLER

WITH AIR SEPARATION

For accurate, uniform pulverizing or blending. Instantly adjustable to finenesses down to 400 mesh, even micron sizes. Dries and grinds simultaneously. Automatic, self-adjusting feed. Many other exclusive engineering and construction features. Williams Roller Mills have set new standards for product quality, output and economy.

OTHER WILLIAMS EQUIPMENT

IMPACT and DRIER MILLS
AIR SEPARATORS
VIBRATING SCREENS
COMPLETE "Packaged" PLANTS For
Installation In Existing Buildings

At Your Service WILLIAMS TESTING LABORATORY

Consult Williams on any grinding, crushing, shredding or separation problem, no matter what the material. The most complete testing and research facilities are available without cost or obligation. Simply furnish enough raw material for a test run, and a sample or description of the finished product desired.



NFA

Plans November Meeting at Roney Plaza Hotel, Miami Beach Big Attendance is Expected

THE National Fertilizer Association will hold its annual fall convention at the Roney Plaza Hotel, Miami Beach, Fla., November 19 through 21, the NFA has announced.

The board of directors will meet on November 19 and the regular convention program will begin on the morning of November 20. A banquet will be held on the evening of November 21 at the conclusion of the formal proceedings.

Dr. Russell Coleman, NFA president, said that representatives from Federal and State agricultural agencies, and a representative from the American Bankers Association will address the convention. In addition, an outstanding national political leader will discuss matters relating to the fertilizer industry and the national economy.

Since registration is heavy, it is probable that the Roney Plaza Hotel will not be able to acommodate all those who desire to attend the meeting, the Association has indicated. In order to provide sufficient accommodations for those attending the convention, arrangements have

been made with the close by Surf Side and Traymore Hotels to provide for the overflow. Reservations at all hotels will be handled by the Roney Plaza.

Since the business meetings will be morning affairs, ample time is provided for convention visitors, if they wish, to take advantage of golfing and fishing facilities available in this resort area. Also, those who wish to extend their travel, may make arrangements to visit Nassau in the Bahama Islands or Havana, Cuba, either before or after the convention.

Below: Air view of Roney Plaza Hotel, scene of NFA's 1952 Fall Meeting



FORMULATORS!

Only ORTHO Lindane* does all these jobs - so well!



- and gives you all these extras:

GUARANTEED GAMMA

- minimum pure gamma isomer 100%. ORTHO Lindane assures you of true Lindane quality actually higher than minimum Government requirements for pure Lindane.

EASY FORMULATIONS

-easily handled-easily formulated as a spray or dust. ORTHO Lindane particles are dry, free-flowing. Easily ground to micro-size.

STABILITY

-chemically stable.

MANUFACTURING "KNOW-HOW"

-made exclusively by the original manufacturers of Lindane in the U.S.A.

*ORTHO Lindane, a real profit-maker, is a truly amazing insecticide offering high potency, rapid action, and residual control. Kills more than 200 varieties of insects by contact, vapor action, and stomach poison.

always - you profit with ORTHO

For complete information, "Story of Lindane," write:

CALIFORNIA SPRAY-CHEMICAL Corp.

Portland, Ore. Sacramento, Calif. San Jose, Calif. Fresno, Calif. Whittier, Colif.

Caldwell, Idaho Maryland Heights, Mo. Oklahoma City, Okla. Orlando, Fig.

Shreveport, La. Fennville, Mich. Elizabeth, N.J. Medina, N.Y. Goldsboro, N.C.

Home Office: Richmond, Calif. . 25 Additional Branch Offices

World leader in scientific pest control

Increased Farm Production to Fill "Fifth Plate" Depends on Use of

CHEMICALS



THE title of this paper, "Food for the Fifth Plate", may require a certain amount of explaining so that everyone will know what it means.

To me, the fifth plate is a symbol — a symbol that puts a lot of hard-to-comprehend farm facts about our future on an understandable basis. Through this "fifth plate" idea, we hope the man in the street and the housewife will come to understand certain problems of the farmer, who is expected to keep this nation well-clothed and well fed.

Essentially, the fifth plate story is this: If we keep on populating our nation at the rate we are going now—more than two million increase a year — by 1975 we're going to have at least 25 percent more people to feed and clothe. There will be five people for every four that are here today, making necessary a fifth plate on the table where four places are set now. And from our farms will have to come the extra food to heap that fifth plate full, three times a day.

Practically speaking, the extra food must come from the same land now in use. Those acres will have to be made more productive. A better job will have to be done in preventing loss and waste of products. Farmers will have to spend more if they are to produce more and will therefore be subject to greater financial risks unless the farm programs and good farming reduce the risks.

It will take a lot of doing to make sure of 15 billion more eggs a year, an extra 20 million hogs, more than 10 billion extra quarts of milk a year — about 30 million extra quarts a day — just to keep folks eating as well as we are now eating. Of course, we're not thinking just of 1975, but of every day between now and then — and of the years beyond.

I believe that the manufacturers, distributors, and sellers of agricultural chemicals have at least as much to offer as any other industry serving the farm. The country is going to depend on these to help fill that fifth plate.

Frankly, I don't doubt for a minute that the fifth plate can be filled. We have a handful of good weapons to work with to do the job: research and education; power and machines; conservation, price support, credit, and electrification programs; agricultural chemicals; and, most important, good farmers!

One needs only to know that the NAC organization is not quite 20 years old, to understand the influence that the industry has had on our record agricultural output the last 10 or 12 years. When the National Agricultural Chemicals Association was organized in 1933, there were only 12 chemical manufacturers represented. Now, there are more than 10 times that many.

In 1933, there were about 100 million pounds of agricultural chemicals — not including fertilizers — manufactured, while today the figure is more like one billion pounds. That is spectacular progress.

Progress Widespread

IT is significant that similar spectacular progress has been made in total agricultural production in the United States. Since 1933, when the pesticide industry began to take active interest in the chemical needs of the farmers, farm production has increased by 50 percent. This was done primarily, by increasing the crop production from each acre 48 percent during those intervening years. Consider too, that since 1933 more than 7 million people have left the farm.

D

W. A. Minor

Assistant to the Secretary of Agriculture, U. S. Department of Agriculture, Washington 25, D. C.

Talk by W. A. Minor, Assistant to the Secretary of Agriculture, at the Nineteenth Annual Meeting of the National Agricultural Chemicals Association, Spring Lake, New Jersey, September 3, 1952. Available ...

HANDBOOK

of

AGRICULTURAL PEST CONTROL

by Stanley 4. Bailey

and

Loslio M. Smith

Professor of Entomology, U. of California

Associate Professor of Entomology, U. of California

192 Pages

A PRACTICAL handbook of agricultural pest control, designed for use by the custom sprayer, the pest control operator, farm advisor, agricultural chemical salesman and field worker is now ready. This handbook covers the agricultural chemicals (insecticides, fungicides, herbicides, plant hormones, nutrient sprays, defoliants, etc), their rates of application, useful formulas, as well as chapters on fumigation, spray machines, toxicology, dusts and dusting, aircraft, and mosquito control. Use the coupon below for your copy now!

CONTENTS

The Commercial Agricultural Chemicals
Physical and Chemical Properties
Compatibility
Containers
Furnigants and Furnigation
Toxicology and Residues
Spray Oils and their Properties
Spraying Machines
Rates of Delivery of Spray Machines

Rates of Application for Sprays
Dusts and Dusting
Aircraft
Mosquito Control
Hazards
Miscellaneous Topics
Tables and Formulas
Terms and Symbols

Send Check with Order

TEAR OFF AND MAIL

INDUSTRY PUBLICATIONS, INC.

175 Fifth Avenue

New York 10, N. Y.

Enclosed is our check for \$3.25 (Foreign and Canada \$3.75) covering a copy of HANDBOOK of AGRICULTURAL PEST CONTROL. It is understood that we may return the book within ten days for a full refund.

Company

ADDRESS

BY

Production per man hour has gone up 75 percent, so that today only about 16 percent of our entire population is producing the bulk of the food we eat and fiber for the clothes we wear.

Yet we have by no means fulfilled our farm production potential. We have increased production 50 percent in less than 20 years with the help of a chemical industry that, agriculturally speaking, has barely gotten its feet wet. From here on, production increases will probably come harder. We will need to make greater use of all those factors that will increase production — and the chemicals are important among those factors.

Fundamentally, it seems to me that there are two jobs to be done in the next 20 or so years, if we are to fill that fifth plate.

We must increase our output of food, feed, and fiber from each acre and each man. And, we must better utilize our output by protecting that which we raise.

We've all been giving a lot of attention to that first need; not so much to the second. We've concentrated pretty heavily in the past 10 years on higher yielding crop varieties, grown and harvested with power and machinery that replaced the hired hands. Advances in those fields are by no means complete, but they are much further along — at least in the farmers' understanding of them—than are the protective measures, that save what we raise.

The importance of agricultual chemicals in filling these needs is not in doubt. Agricultural chemicals can do a lot of good in increasing per acre production with less labor; they can do the same in protecting this production.

Chemicals Top List

ROUGHLY, fertilizers and the new soil conditioners should be put on the list of agricultural chemicals that have, and can help increase per acre crop production. Chemical fertilizers have been pretty well sold to the American farmer — both in tons and as an idea that they can increase his crop yields. The fact that farmers have doubled their use of fertilizer in the last seven years bears

that out. Today, 60 percent of America's farmers are using 21 million tons of fertilizer. The farmer may look on these fertilizers he uses strictly in terms of better yields, but they are also allowing him to use his machines, fuel, and power more efficiently, and they are conserving and improving his land.

Yet, we still have 40 percent of our farmers not using fertilizers. And think, too, of the gains that will be made in crop production as the farmers now using fertilizers learn to use them better!

The knowledge of how and when to use fertilizers and in what amounts to get a top return in crop yields has yet to be exploited. The Department of Agriculture and the Land Grant Colleges are developing a fertilizer utilization program that should aid in getting fertilizer used so as to increase production of needed crops, lower costs and increase farm income.

I can't be so positive about soil conditioners because we haven't seen them in use for 20 or more years as we have fertilizers. But the indications are that they can help us make use of much of our hard-to-farm land. They can reduce the amount of runoff from our clay soils; make planting easier, encourage seed germination, and allow the roots of important crops to get a better foot-hold in the soil.

Protecting what we produce is a wide-open field for agricultural chemicals. I know that many of you are kept busy manufacturing and selling a few specific chemicals — perhaps only a single insecticide, so it's worth while, I believe, to take a brief look at the pesticide industry as a whole — to check and see how widely these chemicals are used on the farm today.

We have our herbicides that prevent weeds from stealing valuable plant-building moisture and food away from the crops. There are the fungicides, that can prevent losses to disease; defoliants that allow greater harvest of some crops; hormones that give us bigger fruits, and possibly meatier animals. We have rodenticides, that keep the rats and mice from stealing and destroying our stored foods; fumigants that protect our stored foods, protect foods in transport, and control some soil-infesting pests. And, of course, we have insecticides, which often make the difference between a crop and no crop.

Herbicides that replace the cultivator and the hoe have been a Godsend to the corn and small grain farmers who last year treated more than 30 million acres. The experimental use of herbicides in the potentially great corn country of the South has tripled corn yields. In cotton, their use reduced the cost of weed control 31 percent below the cost of mechanical weeding. The use of herbicides to remove sagebrush from Southwestern ranges has increased beef production from these acres as much as 50 percent.

The hormone sprays delay the blossoming of fruit trees and thus avoid the damage of a cold snap. Trees can be prevented from shedding their fruit at too early a date. These chemicals can be used to thin a fruit tree so that the remaining fruits are larger and of better quality.

Defoliants are being used widely in the South to hasten the maturity of the cotton and to make mechanical harvesting an easier job. Defoliants reduce losses to diseases and insects as well. On potatoes, defoliants reduce the damage from late blight disease. They make harvesting easier, and provide a more rugged, bruise-resistant spud.

Rats and mice destroy 4 percent of our grain every year. The Fish and Wildlife Service figures that there is about a rat apiece for each of us in the U. S. — about 150 million of them — taking a \$2 billion toll annually. Rodenticides certainly can help us fill our fifth plate.

Plant Disease Control

USE of agricultural chemicals to prevent fungus disease attacks on our crops has been stepped up considerably with the development of the newer organic compounds. Fungicides are serving us well. They prevent the disease, apple scab, that in

(Turn to Page 121)

SERVE YOUR AGRICULTURE BEST

Sherwin



Williams

Insecticides

Fungicides

Spray Conditioners

- S. W. Arsenate of Lead
- S. W. Arsenate of Calcium
- 5. W. Aphamite 15% Parathion)
- S. W. Benzene Hexachlorida Emul.
- 5. W. Karbam Black
- S. W. Mulsoid Sulfur (Micro-Fine)

- S. W. DDT 50% W. P. S. W. DDT 25% Emul.
- S. W. Dormant Spray Oil
- S. W. Dimite
 - S. W. Karbam White
- S. W. Sulfix Sulfur

THE SHERWIN WILLIAMS CO.

AGRICULTURAL CHEMICAL DIVISION

100 PARK AVENUE, NEW YORK, N. Y.

SOLD BY

MANUFACTURER'S AGENT

SEACOAST LABORATORIES, INC.

156 PERRY STREET, NEW YORK 14, N. Y.

MAKERS OF



UNIFORMLY BLENDED SPATULA TESTED DUSTS

ROTENONE DDT PARATHION DDD

LINDANE CHLORDANE CPR CALCIUM ARSENATE METHOXYCHLOR

COPPER **FERBAM** ZIRAM DITHANE

DEPENDABLE SPRAY MATERIALS

DDT 50% W. P. DDT 25% EMUL. LINDANE 20% EMUL.

CHLORDANE 50% W. P. LINDANE 25% W. P.

MOSQUITO LARVICIDE FOR SURFACE SPRAYING AND FOGGING

The Listening Post

Fungicides Tested on Fruit Fire Blight

This department, which reviews current plant disease and insect control problems, is a regular monthly feature of AGRICULTURAL CHEMICALS. The comments on current plant disease problems are based on observations submitted by collaborators of the Plant Disease Survey Bureau of Plant Industry, Soils, and Agricultural Engineering, U. S. Department of Agriculture, Beltsville, Md.



By Paul R. Miller

IRE blight caused by the bacterium Erwinia amylovora has been one of the principal limiting factors in the production of apples and pears in Colorado, according to a report by W. D. Thomas, Jr., and W. J. Henderson, of the Agricultural Experimental Station and Agricultural Extension Service in Colorado. It has been conservatively estimated that this disease causes an annual loss of about 10 percent of the entire apple and pear crops in the state, or about 157,000 bushels of apples and 19,900 bushels of pears.

Until 1947, only Bordeaux mixture had been used as a spray in Colorado to combat fire blight. Spray tests for the control of powdery mildew in 1946 in western Colorado, however, indicated that trees sprayed with "Dithane D-14" in the cluster bud and calyx stage of blooming had far fewer spurs infected with fire blight than did the unsprayed checks.

Spray tests for the control of fire blight were established in Fremont and Mesa Counties in 1947. The original experiments were designed to investigate the efficacy of carbamate fungicides in controlling this disease on apples and pears.

In 1947 plots were established on pears in Mesa County and on apples in Fremont County. "Dithane D-14" and Zerlate" were used in the former plot and "Dithane D-14" only was used in the latter. Two quarts of "Dithane" were mixed with one pound of zinc sulfate and one-half pound hydrated lime per 100 gallons

of water. "Triton B-1956" (one ounce) and 50% DDT (two pounds) were added to all treatments. Check plots were sprayed with 50% DDT only. Applications were timed as indicated in Table 1.

The treatments were applied to three-tree blocks in four randomized replications. An unsprayed buffer row and an unsprayed three-tree buffer block was left between each sprayed row and block to prevent overlapping of spray materials. Observations were made by counting the total number of infected spurs

on the center tree of each non-sprayed check and sprayed block.

The results shown in Table 1 indicate that the reduction of infection on pears by "Dithane" and "Zerlate" was highly significant, although there were no significant diferences between the treatments or times of application. Effective control on apples was obtained when spray was applied at the full bloom stage, but when sprayed at the 10-percent bloom and again at full bloom stages results were highly significant.


Results were based on the count of infected fruit spurs, which were classified by an arbitrary index of infection from 1 to 6: (1) no infection; (2) 1 percent to 10 percent infection; (3) 11 percent to 30 percent; (4) 31 percent to 50 percent; (5) 51 percent to 80 percent; and (6) 81 to 100 percent.

Because growers maintain that a 50 percent control of a heavy infection of fire blight in an orchard allows sufficient new growth and set of healthy fruit for a satisfactory crop, and since more than 50 percent control was obtained in these tests, similar tests were conducted in 1948 and 1949.

TABLE 1

Results of spray tests for the control of fire blight on apples and pears in 1948 in Colorado

Treatment	Time of application	Averag		Percent reduction on No. infected spure		
		Pears	Apples	Pears	Apples	
Dithane D-14	10% bloom	10.0	4.8	75,0	60.4	
	Full bloom	16.7	3.3	62.5	72.9	
	10% bloom and					
	full bloom	13.2	3.8	67.5	68.7	
	10% bloom, full					
	bloom, and 90%					
	petal fall	8.5	6.0	79.2	50.5	
Zerlate	10% bloom	15.5		61.3		
	Full bloom	13.0		68.0	*	
	10% bloom and					
	full bloom	13.5		66.8	*	
	10% bloom, full					
	bloom, and 90%					
	petal fall	12.5		69.3		
Check	Unsprayed	40.2	12.0		2	
	LSD 556	19.5	5.5			
	LSD 1%	26.0	7.6	+		



*U.S. Pat. No. 2,529,494

-UNITED STATES RUBBER COMPANY -

Naugatuck Chemical Division, Naugatuck, Conn.

manufacturers of <u>seed protectants</u> — Spergon, Spergon-DDT, Spergon-SL, Spergon-DDT-SL, Phygon Seed Protectant, Phygon Naugets, Phygon-XL-DDT, Thiram Naugets — <u>fungicides</u> — Spergon Wettable, Phygon-XL — <u>insecticides</u> — Synklor-48-E, Synklor-50-W— <u>fungicide-insecticides</u> — Spergon Gladiolus Dust, Phygon Rose Dust — <u>miticides</u> — Aramite.

In 1948, zinc sulfate was used at the rate of two pounds per 100 gallons of water, in order to determine whether or not it was instrumental in securing control of fire blight. "Dithane Z-78" and "Puratized Agricultural Spray" were also used as treatments, applied at the rates and stages indicated in Table 2. "Triton B-1956" (1 oz, per 100 gal.) was added to all spray materials. No DDT was used in 1948. No treatments were used on the checks. The methods and locations in 1948 were similar to those used in 1947.

As shown in Table 2, "Dithane Z-78" gave a significant reduction of fire blight on pears when applied at the full-bloom stage. There was also a highly significant reduction in the percentage of infected spurs on both apples and pears when "Dithane Z-78" was applied at both the 10 percent bloom and fullbloom stages. However, an additional application made in the 90 percent petal fall stage failed to give a significantly better control than an application in the 10 percent bloom followed by an application in the full bloom. This was undoubtedly due to the short-blooming period under Colorado conditions. In states where the blooming period covers a longer time, an application in the 10 percent bloom followed by applications in the full-bloom and 90 percent petal fall would be needed. Neither "Puratized Agricultural Spray" nor zinc sulfate gave satisfactory control of fire blight in these tests.

1949 Tests on Apples

IN 1949, the tests were conducted on apples only in Montrose County. An entire orchard was included in this test and consisted of 118 Jonathan trees and 114 Delicious trees. The varieties alternated in the rows. Two check blocks of nine trees each were left unsprayed in the orchard. All other trees were sprayed with "Dithane Z-78" (2 lbs. per 100 gals. water) to which "Triton B-1956" (½ oz. per 100 gals.) was added. Applications were made at the 10 percent bloom stage with subse-

TABLE 2

Results of spray tests for the control of fire blight on apples and pears in 1947 in Colorado

Treatment	Rate of application when applied and stage of bloom				Percer infected		Percent control over check	
	Per	100 gal.	Bloom Stage		Pears	Apples	Pears	Apples
Dithane Z-78		2 lbs.	Full		5.4	8.2	66.7	37.0
		do.	10% + full		0.8	1.1	95.1	91.0
Puratized Agricultural		1 pt.	Full		17.5	•	-1.3	
spray		do.	10% + full		25.0		-64.8	*
Zinc sulfate		2 lbs.	Full	3	22.5	11.3	-72.0	15.0
		do.	10% + full		12.5	7.8	-228	22.6
Check					16.2	13.3	*	
			LSD 5%		6.4	6.0	*	
			LSD 1%		8.7	8.4		*

quent applications in the full-bloom

For the Jonathan variety results show that 88.1 percent of the sprayed trees and 33.3 percent of the unsprayed trees had from none to a trace of infected spurs; 8.5 percent of the sprayed trees and 39.0 percent of the unsprayed trees had from 1 to 10 percent infection; 1.7 percent of the sprayed trees and 22.3 percent of the unsprayed trees had from 11 to 30 percent infection. and 1.7 percent of the sprayed trees and 5.6 percent of the checks had from 31 to 60 percent infection.

Of the Delicious variety, 97.7 percent of the sprayed trees and none of the unsprayed trees had from zero to a trace of infected spurs; 8.5 percent of the sprayed trees and 39.0 percent of the checks had from 1 to 10 percent infection. None of the sprayed trees had any greater infection. Of the check trees, however, 55.5 percent had from 11 to 30 percent infection.

In 1950, the same procedure was used as in 1949 on Jonathan trees in the same orchard, with the exception that three entire rows were left at random as unsprayed checks. Approximately 50 percent of the trees in the unsprayed check had a medium amount of twig infection and only a trace of fire blight occurred in the sprayed trees.

In these spray tests, conducted on apples and pears at several locations in Colorado in 1947, 1948, 1949, and 1950 for the control of fire blight, it was found that application of "Dithane Z-78" (2 lbs. per 100 gals. water) in the 10 percent bloom and a subsequent application in the full-bloom stage reduced the incidence of current season blossom and twig infection approximately 75 percent.

This spray method, together with sanitary pruning and treatment of hold-over cankers, has become a standard recommendation for fire-blight control in Colorado.

In areas where the blossoming period is considerably longer than that where these investigations were conducted, an additional application should be made about the 90 percent petal-fall stage of blooming.

Sturtevant Fertilizer Equipment

Increases Production...
Cuts Costs of Fertilizer
Manufacture

Announcing the New Continuous Superphosphate Manufacturing Process

For over 50 years, the Sturtevant Mill Company has been a leader in the design and manufacture of fertilizer equipment. Equipment that has cut costs, increased tonnage throughout the industry.

Because of the demand for a continuous process for making superphosphate, Sturtevant proudly announces its Continuous Den to make the same high grade material as has been produced for years in the Batch Den and Excavator. If you are planning to install a continuous superphosphate system, investigate this equipment, today.

Either the Continuous Den or the Batch Den produces super in excellent physical condition with low insoluble and moisture content.

Sturtevant
Den & Excavator

Produces Quality Superphosphate Economically The Sturtevant Den and Excavator Units provide a complete method of producing superior, bulky superphosphate either single, enriched

or triple... quickly and economically. The Batch Den is easily operated by two men to produce 16 to 40 tons per batch and up to 480 tons per day while the Continuous Dens are designed for 12 or more tons per hour. Ruggedly built, they require little if any maintenance.

Sturtevant Granulation Process

Sturtevant fertilizer granulating units provide you with a complete process for manufacturing granular fertilizer. These efficient units can be supplied for various hourly tonnages and certain granule sizes depending on your particular requirements.

Write for complete information about Sturtevant Mill Fertilizer Equipment. Sturtevant Mill

123 CLAYTON STREET - BOSTON 22, MASSACHUSETTS

Insect Pest Reports From Many U. S. Points

This column, reviewing current insect control programs, is a regular feature of AGRICULTURAL CHEMICALS. Mr. Dorward is connected with the department of Insect Peat Survey and Information, Agricultural Research Administration, Bureau of Entomology and Plant Quarantine, U. S. Department of Agriculture, Washington. His observations are based on latest reports from collaborators in the U.S.D.A.'s pest surveys throughout the United States.

1 Ann la

By Kelvin Dorward

OT dry weather of late July and early August influenced European core borer activity considerably. In Kansas, a combination of drought conditions and an abundance of lady beetles aided materially in reducing the second-generation population of the borer. Second-generation borers were expected to be light in southern Illinois and Tennessee, primarily because of drought conditions. Indiana and Ohio also indicate that second-brood borers may be light. Not all states have reported a downward trend, however. Second-brood European corn borer moths were emerging in early August in Wisconsin and indications were that the second-brood population might build up to the level of the 1949 infestation. The same is true for Minnesota where the peak of the second-brood emergence was expected during the last two weeks of August. In Iowa, borer development continued a week ahead of previous years. First-brood infestation was fairly high across northern Iowa with second-brood oviposition spread generally and not concentrated in late planted fields. Peak oviposition in Iowa was anticipated to be about August 8. Maryland and Massachusetts in early August expected increased activity from the borer. Montana, where the borer was first reported in 1951, recorded light infestations in irrigated corn in Dawson and Richard Counties, late in July.

Reports received during early August from several states indicate that grasshoppers were causing damage or their populations were building up. In the Nashville, Tennessee, area,

grasshoppers were abundant and were damaging practically all foliage. The drought condition was thought to be instrumental in the overall feeding of the hoppers. Other states reporting grasshoppers included Georgia where peaches and cotton in Spalding County were damaged as were fescue and orchard grass in Butts County. Control was needed in some Bahia grass plantings in northern Florida. In some northwestern North Carclina counties corn, tobacco, ulfalfa, and grasses were being damaged as wer: red clover and corn in Queen Anne and Kent Counties, Maryland. Grasshopper populations were building up in some Ohio clover and alfalfa fields to a point where control was necessary. Spotted areas of Illinois, mostly in the southern section, had populations high enough to damage clover, soybeans, and corn. Populations in the southern areas of Wisconsin continued to build up. Wide-spread damage to oats occurred in one section of Burnett County which is in the northwestern part of the State. In Cache County, Utah, alfalfa, wheat and range were being damaged. Infestations were high in the cultivated areas of the south central part of Idaho and range species were moving into fields on the edge of irrigated

During early August, secondgeneration chinch bugs caused considerable damage to late thin corn and summer grasses in Illlinois. The highest overwintering population in 20 years is expected in that State unless adverse weather conditions are encountered.

The heaviest lygus bug populations on alfalfa since 1940-1943 were reported from Utah in early August. Reinfestation after bud stage treatment was particularly heavy. These insects were also building up in population in southern Idaho alfalfa and clover seed fields to a point where control measures were being applied. Damage to seed alfalfa was reported from Mesa County, Colorado and heavy populations were on alfalfa in Park County, Wyoming and legumes in Tennessee.

Cotton Insects

URVEYS being conducted in con-Snection with the pink bollworm continue to show an alarming build up. In the July issue of Agricultural Chemicals, it was reported that bloom inspections in the Coastal Bend area of Texas showed a much heavier pink bollworm infestation than in 1951. Gin trash inspection under way in early August of this year in the Rio Grande Valley of Texas presented some interesting figures. The average number of pink bollworms found in gin trash per bale of cotton ginned was as follows: Cameron County, 3,625 pink bollworms in 1952 compared to 23 in 1951: Hidalgo County, 861 to 22; Starr County 1,811 to 79; and Willacy County, 2,942 to 2. During early August, first seasonal infestations were found in three additional west Texas counties: Coke, Glasscock, and Howard. This makes a total of 10 west Texas counties found infested this season by the bloom inspection method as compared with one county of the same group found infested last season using the same inspection method.

In early August, the boll weevil infestation increased in many fields of central, northcentral and northeast Texas to a point where heavy damage resulted. Weevils were migrating in Oklahoma and heavy infestations were expected in fields of succulent cotton. In southwest Oklahoma, an increase was noted during the first week of the month. Weevils were on the increase in Arkansas and sharp increases were experienced in fields that had had infestations throughout the season. Despite the dry weather both North FOR EVERY PROCESS OF

INSECTICIDE PRODUCTION...

Equipment of Proved Performance at Minimum Cost








STANDARD TYPE R.T.R. Uni-Blender Compounding Plant is designed to mix and blend dust concentrates with diluents to produce and package ready to use, fleld strength insecticides of consistently uniform quality. The complete plant requires only 9'x12' of floor space and 13' of head room. Produces up to four 40 cu. ft. batches per hour with only one operator. Users report production of 29,000 pounds in 5 hours with two operators. **DUAL TYPE R.T.R. Uni-Blender Com- pounding Plant** is designed to perform a dual function . . . (1) to formulate dust concentrates . . . (2) to produce field strength insecticides. Users claim this provides
the most economical insecticide formulating equipment
available. Handling 40 cu. ft. batches, the complete plant
requires only 11'x11' of floor space and 16' of head room.

Both Standard and Dual types are available for immediate shipment.



Uni-Blender Liquid Formulating Plants

provide for liquefying and injection of specific technical grade toxicants required for the formulation of numerous concentrates and liquid sprays.

Write for descriptive bulletins

A. E. POULSEN & CO.

3305 East Slauson Avenue, Los Angeles 58, California

Engineers and Manufacturers
Materials processing and materials handling equipment

Established 1939

and South Carolina reported an increase of boll weevils in some areas. In the Baton Rouge area of Louisiana there was a slight increase, but in other areas of the State the trend was slightly downward.

Bollworms were causing damage in some states even though in scattered areas. These insects were on the increase in several North Carolina Counties but damage was not heavy at the time the report was made. The same was true for the Upper Coastal Plains, Sandhills, and Piedmont Counties of South Carolina. In some areas of Georgia, damage was rather high. Injurious infestations of bollworm continued in fields of succulent cotton in Texas but the trend was downward in the central and northern areas. In the Messilla Valley of New Mexico bollworms, eggs and larvae increased sharply during early August. During this period bollworms were the outstanding cotton insects in Graham and Cochise Counties. Arizona.

Spider mites were on the increase in South Carolina, Georgia, Tennessee, Misisippi, Louisiana, Texas, and California. Control measures were being applied in Louisiana and Texas and damage was also reported from South Carolina and Mississippi.

Vegetable Insects

PIDER mites were becoming in-SPIDER mites were considered and services, and services, and services are services and services are services. sugar beet, peppermint and alfalfa in the Prosser vicinity of Washington during the first part of August. Considerable damage in this area is expected before the end of summer. In Orange and Ventura Counties, California, the heavy populations of the two-spotted spider mite were building up along the edge of many lima bean fields. Control measures for mites were necessary in some bean fields of south central Idaho. In eastern Virginia, beans, tomatoes, and strawberries were being injured while on the Eastern Shore of that State, mites were damaging soy-beans.

Leafhoppers, in early August, were damaging vegetables in many sections of the United States. In Mas-

sachusetts they were abundant on beans and lettuce while in Niagara County, New York they were destructive on practically all crops. In Wayne County, New York, leafhoppers were present in "cloud" numbers on lettuce growing on muck soil. Other New York counties reporting injury to various vegetables were Jefferson, Erie, Cayuga and Saratoga. Damage was also reported from Milton and Georgetown areas of Delaware. In the Chillicothe and Columbus areas of Ohio beans were damaged. Leafhoppers were unusually abundant throughout Wisconsin on alfalfa, beans, potatoes, cucumbers and other vegetable crops.

In the eastern, middle and old tobacco belts of North Carolina, horn worms were the heaviest in recent years. Damage to tomatoes from hornworms was reported from South Carolina, California, and Utah.**

"Endrin", "Isodrin" Chosen

The Interdepartmental Committee on Pest Control has recently approved "endrin" and "isodrin" as coined names for insecticides. The names have been cleared by a Patent Office search which makes them available as common names for designating the respective compounds.

Endrin is the insecticidal chemical 1,2,3,4,10,10-hexachloro-6,7-epoxy-1,4,4a,5,6,7,8,8a-octahydro-1,4,5,8-endo-endo-dimethanonaphthalene. The formula is identical to dieldrin, except that dieldrin is the endo-exo isomer.

The material is insoluble in water but is soluble in the usual organic solvents. The solubility of endrin in aromatic hydrocarbon solvents such as benzene and xylene is greater than in paraffinic solvents such as hexane or kerosene.

It is stable to alkalis and unaffected by acidic conditions normally encountered in agriculture. It is compatible with the commonly-used insecticides, fungicides and herbicides, the Committee states. As to phytotoxicity, endrin seems to be tolerated by most plants when used at concentrations necessary for insect control.

Insect pests against which endrin has shown promise, include the boll weevil, the bollworm, tobacco budworm, the variegated cutworm, the fall armyworm and the tarnished plant bug. The material has also "looked promising" against tobacco hornworm, sugar beet webworm, corn earworm, Mexican bean beetle, European corn borer and various aphids.

Tests for toxicity have led to the conclusion that due to its high order of insecticidal effectiveness, coupled with its moderate residual characteristics, the mammalian toxicity of endrin is not expected to preclude the use of effective quantities except on food and forage crops nearing harvest.

Isodrin is the coined name for the insecticidal chemical 1,2,3,4,10,10lexachloro-1,4,4a,5,8,8a-hexahydro-1,-4,5,8-endo-endo-dimethanonaphthalene. The formula is identical to that of aldrin, except that the latter is the endo-exo isomer. Its properties of solubility and compatibility are like these of endrin.

Isodrin seems to be tolerated by most plants when used at concentrations necessary for insect control although corn buds have shown signs of "burn". The chemical shows promise for control of European corn borer, corn earworm, sugar beet webworm, tobacco hornworm, thrips, leafhoppers and various aphids.

As to toxicity, the material must be handled with caution, according to the Committee report, but, like endrin, it is expected to be safe for use, except on food and forage crops nearing harvest time.

Both materials must be handled with extreme care, the Committee report declares. Contaminated clothing must be removed and laundered and products must be washed off skin promptly. Formulating and mixing plants must be well-ventilated to prevent inhalation. "During application, the wearing of respirators should be enforced," the report concludes.

Now-62 new uses make PARATHION PARATHION more effective than ever! This was 62 new disease.

This year, 62 new claims for parathion have been accepted for labeling purposes by the United States Department of Agriculture. These new claims give parathion a total of more than 200 accepted uses...on more than 50 crops!

Here is new proof that parathion is the most widely effective of all insecticides:

ALFALFA Alfalfa caterpillar

APPLE
European sawfly
Leafhopper
Pear borer

APRICOTS

Pandemis moth

BERRIES

Mackberries:
Red-necked cane borer

Boysenberries and dewherries: Aphids Red spider mite

Cranberries: Lecanium scale crawler Currents and gooseberries:

Aphids Red spider mite

Respherries: Aphids Red spider mite CABBAGE, MUSTARD, TURNIP, KALE, RADISH AND CARROT

Vegetable weevil

CHERRY
Casebearer
Fruit fly
Leaf miner
Pandemis moth
Red spider mite
San Jose scale crawler
Shot-hole borer

CITRUS Aphids

CORN

Budworm Corn earworm Leaf aphid Fall armyworm Sap beetle

CUCUMBER, SQUASH AND MELONS Stink bug

Vine borer

GRAPES

Grape berry moth

Aphids Leafhopper

ORNAMENTALS (autdoors)

Bagworm Cottony-cushion scale Juniper webworm Lace bug Lecanium scale Thrips

PEACH

Lecanium scale Red-banded leaf roller Shot-hole borer

Pea weevil

PECAN

Black pecan aphid Nut casebearer Red spider mite Twig girdler Walnut defoliator Webworm

Yellow aphid

PINEAPPLE (before planting)
Mealybug

PINEAPPLE (treatment of beds)
Mealybug
Cricket

POTATO Armyworm

PRUNE AND PLUM San Jose scale crawler

Soybean Velvetbean caterpillar

TOBACCO (seedbed) Flea beetle

Flea beetle Midge larvae Vegetable weevil

TOBACCO (field)
Flea heetle
Hornworm
Vegetable weevil

TOMATO Armywerm

Consult Jocal agricultural authorities for suggestions on desages and application procedures.

AVAILABLE FROM NATIONAL MANUFACTURERS

AMERICAN Cyanamid COMPANY

Handadurer at Tigather Parettien Technics

Agricultural Chamicals Division

Write for new 1952 Parethion Grower's Hamiltonia

ATTENTION DISTRIBUTORS AND DEALERS: Are you taking adventage of the union approximates affered by permitting insurface.

Suppliers' Bulletins

Bag Conveyors Described

Information on "Power Curve" machines for handling bagged materials, has been issued by the Power Curve Conveyor Co., Denver, Colo. Designated as bulletins number 50 and 60, details are given to describe the operation of the devices.

Bulletin #50 contains information on belts for loading trucks and box cars. Completely flexible, the loaders convey bags all the way from packing stations into the truck or car at the rate of 30 bags a minute. The conveyor may be sent to any angle up to 90° while under power and carries bags around corners without interruption, the folder says.

The other bulletin, #60, describes the firm's machines for flattening bags to save room in warehouses and in shipping. The bag is carried between two power-driven conveyors and a pressing and rolling action is applied. The conveying element consists of a series of endless steel spring belts operating over grooved rollers. Spacing between rollers is quickly adjusted by hand, the makers state.

Write for folders #50 and 60 in care of the company, 1235 So. Broadway, Denver, Colo.

Magnetic Trap Announced

A new four-page, two-color brochure, "Magnetic Pipeline Traps for Sanitary and Standard Applications," has been issued by the Eriez Manufacturing Company. The firm's permanent (non-electric) magnetic pipeline traps are designed to remove tramp iron from liquid flow lines. In doing so they afford protection to processing equipment such as pumps, screens, grinders, fillers, etc.

The traps will not corrode and are unaffected by high temperatures, their makers state. Cast of 316 stainless steel with fused magnetic face plates, they will withstand pressures up to 75 psi. The traps have wide use in the chemical industry including fertilizer manufacture.

Eriez' brochure describes with illustrations and data typical magnetic trap installations. Also included are sections on variables affecting magnetic pipeline trap size, and a material classification of products affecting the capacities, in gallons per minute, of the pipeline traps.

 The new brochure (No. B-203) is available from the Eriez Manufacturing Company, Erie, Pa.

Dust Filtering Unit

Filtering and recovery of dust from two or more exhaust fans, by the same filtering unit and without mixing of products, is claimed for the Day "AC" dust filter equipped with plenum splitters. Dust-laden air from each exhaust fan is filtered separately to allow reclamation of product without contamination. (see cut below)

Licensed under Hersey patents, the filter uses felt tubes as the filtering medium. Filter cleaning is said to be continuous and automatic, by means of slow-traveling reverse air jet rings which dislodge collected dust from the filtering area and discharge it to the air stream leading to the bottom of the unit.

Because of this operation, periodic shut-downs of sections for cleaning are eliminated. The Day "AC" filter also permits use of higher air pressures, at air-to-cloth ratios as high as 20 to 1, to handle greater volumes of air per unit. Further details are available from the makers, Day Company, 810 Third Ave., N. E., Minneapolis, Minn.

Stauffer Catalog Revised

Stauffer Chemical Company has just issued a revised 112 page edition of their general catalog covering basic chemicals for industry and agriculture. As in the earlier editions, considerable data are given on all products, and numerous tables on specific gravity, viscosity, specific heat, solubility and other properties give valuable assistance to the user of Stauffer chemicals. Copies available. Write the company, 420 Lexington Ave., New York 17, N. Y.


Pest Control Film Out


Completion of a new 16 mm color and sound motion picture entitled "Livestock Pest Control" has been announced by the California Spray-Chemical Corporation.

Local premiere showings of this new educational movie are now being scheduled.

Pests such as the screw worm, the sheep ked, the heel fly and cattle grub are shown in extreme close-ups, and in "action." Practical, "how-todo-it" scenes show the newest sanitation and control techniques now being

(Turn to Page 137)





for your potash needs

THIS fall, as scheduled, the modern plant and refinery of the Southwest Potash Corporation will begin producing its <u>HIGH-K</u> Brand* Muriate of Potash (60% K₂O min.) in substantial quantities. Rated capacity at the Carlsbad, New Mexico, mine is expected to be about

210,000 tons of K2O annually.

Recognizing the increasing need of both agriculture and industry for high quality potash, Southwest Potash Corporation has designed its facilities to permit rapid expansion to more than twice initial capacity.

Southwest Potash Corporation



AT BROADWAY & NEW TORK & M. S.

© 1952 SPC

Technical Briefs

5 Year DDT Residue

Recent research by the U. S. Department of Agriculture indicates that a single application of DDT to the soil will kill some species of insects for at least five years.

Tests conducted in several locations in the northern states show an average DDT carryover in turf of 92 percent after two years, of 43 percent after four years, 36 percent after six years, and 29 percent after seven years.

Entomologists also found that the rate of application had little influence on the percent of toxic residue in the soil. Soil samples, treated at the rate of 25 and 50 pounds of DDT per acre, show the percentage carryover after eight years of testing was about the same in either case.

Tests with different soil types proved that DDT lasts longer in poor soils and a shorter time in soils high in organic matter. Results of experiments with samples of 85 different soil types from seven states, showed that persistence of DDT was about the same in sandy, silt, or clay loams, but much greater in soils heavy in sand.

After analyzing 223,000 Japanese beetle grubs it was found that DDT began losing its killing power only in the fifth year after application. A single DDT treatment at 25 pounds per acre produced a 98 percent grub kill within four weeks after the grubs were put in the soil, during the first four years. It required slightly more than five weeks for the same kill the fifth year.

Potatoes Need Boron

Should a boron fertilizer be applied to potatoes? Experiments carried out recently on potatoes in Austria have shown that in certain areas the soil shows a marked boron deficiency and that shoots of potato plants stop growing. The upper leaves are dark green with a tough, shiny surface, while the edges of the pinnules often curve upwards and in-

wards. The older leaves gradually turn yellow and the shoot tip withers. Symptoms of boron deficiency in the root vary with the variety. A characteristic symptom is that the vascular ring turns partially or completely brown. Rapid discoloration of the cut surface of the potato is another sign of boron deficiency. Soils derived from weathered rocks have been found to be the most deficient in boron.

—Steineck, O., Die Landwirtschaft, 1951, No. 23-24 (Inst. Pflanzenbau und Pflanzenzuchtung der Hochschule für Bodenkultur, Vienna.

Methyl Bromide Fumigation

A method for fumigating bulk cottonseed in freight cars, developed between 1945 and 1951, is described in this paper.

Efforts were made to attain a satisfactory distribution of methyl bromide throughout a freight-car load of cottonseed by various methods of application, such as applying it in the head space above the load, injecting it into the load, applying half the dosage above and injecting half, and dissolving in carbon tetrachloride and spraying on the surface. None were successful. Other innovations, such as partial loads, additional sealing of floors and doorways with gas proof material, providing risers (vertical and horizontal tunnels) through the load to assist in penetration, and following the application with a blast of carbon dioxide, were also unsuccessful.

Attempts were then made to adapt the method of forced circulation used in fumigation of cottonseed in large steel storage tanks. A gas-distribution pattern was obtained that provided a lethal concentration at all sampling points. In the method finally adopted, a portable blower operated outside the freight car pulls air from beneath the load through a specially-designed duct system and returns it to the space above the load. The blower is run during gas volatilization and for 2 to 10 minutes thereafter,

then disconnected, the car sealed, and the blower moved to the next car to be fuminated.

This method for treating cottonseed was authorized for use on quarantined cottonseed in February, 1950, in a limited area under supervision. The dosage schedule is 7 pounds per 1,000 cubic feet for 24 hours exposure at 60°F or above, and 8 pounds at lower temperatures.

The first trial on a commercial basis was at Lubbock, Tex., where approximately 12,000 tons of cotton-seed in 306 freight cars were fumigated. All but one of the 162 cars that were checked by gas analyses were found to have the required gas concentration.

In July, 1950, the forced-circulation method was authorized as an alternate method for treating cottonseed for the destruction of pink bollworm larvae.

—Summary of paper, "Methyl Bromide Fumigation of Cottonseed in Freight Cars for the Destruction of Pink Bollworms," by G. L. Phillips, U.S.D.A., Bur. of Entomology & Plant Quarantine, Div. of Stored Product Insect Investigations.


Puts Conditioner on Rows

In an effort to make the use of soil conditioners economically feasible for agriculture as well as for the city gardener, American Cyanamid Company is experimenting on its product, "Aerotil" applied in narrow bands to rows where seedlings emerge.

The idea is, according to an industry spokesman, to place the soil conditioner where it can do an optimum amount of good in preventing a crust which hinders emergence of many crops.

By limiting the application to a strip from 1 to 1½ inches wide, a saving of some 90% may be realized, from the cost of treating an entire acreage. It is thought that by employing such a method, or one akin to it, that the cost might be brought as low as from \$15 to \$30 per acre. For certain high-priced crops, enabling seedlings to emerge quickly would be well worth the relatively low cost of applying a good soil conditioner, it is said.

American Cyanamid's experiments are not far enough along at




ATLOX' EMULSIFIERS

Producers of agricultural chemicals use more Atlox emulsifiers than any other kind. Atlox emulsifiers are preferred by formulators for every type of insecticide and herbicide—emulsifiable or oil-diluted concentrates for use in knapsack, lowgallonage, or high-gallonage sprayers.

Atlox emulsifiers are virtually chemically inert—give concentrates excellent shelf life. And toxicant manufacturers appreciate the practical help Atlas gives in developing formulas . . . know they can depend on Atlas to meet delivery commitments.

Send for useful literature giving formulas, testing methods, and other valuable information. Technical assistance always is available.



present to have built up any significant data, the company said.

The appointment of three district sales representatives for Monsanto Chemical Company's merchandising division was announced recently.

Stanley W. Sullivan of Chicago, was assigned the Chicago district, Jack R. Glatthaar of Washington, D. C., was assigned the New England, New York and northern New Jersey district and Robert A. Ehrhardt of St. Louis was assigned the St. Louis district. The newlyformed merchandising division is handling the sales of Monsanto's "Krilium" soil conditioner and other trade-marked products through retail channels.

Radioactive Testing

A new and accurate method of using atomic energy to detect and measure impurities in various materials, has been developed by Union Carbide & Carbon Corp. scientists at Oak Ridge National Laboratory. The analysis technique is now being offered as a service to industry by arrangement of the Atomic Energy Commission through Carbide and Carbon Chemicals Company.

Dr. C. E. Larson, director of the Laboratory, explains that the analysis technique involves placing the test sample in the Oak Ridge graphite reactor, or "atomic furnace." The sample is thus exposed to neutron bombardment, making traces of impurities in the material artificially radioactive. Then, using instruments and detectors developed at the Laboratory, the exact quantities of impurities present may be measured accurately.

According to the experience gained thus far in perfecting the method, neutron-activation analysis can be of much practical value in determining minute quantities of elements in biological substances; including fertilizers and feedstuffs; fine chemicals; food and food additives; insecticides and disinfectants.

The new method has several advantages not common to other methods of analysis. Tests have indicated its extreme sensitivity for detecting and measuring many chemical trace elements, so slight that they cannot be determined by other chemical and physical testing processes. The method can be more specific than other techniques, since elements to be tested, when irradiated, produce artificially radioactive isotopes possessing their own characteristics as to modes of "decay" and types of radiation. The characteristics are never exactly duplicated in any other-radio-isotopes produced.

Contamination, often encountered in conventional analyses, is negligible in neutronactivation analysis, existing only when the irradiated material contains large amounts of elements that strongly absorb neutrons.

The new analysis method permits the examination of larger samples than the amounts usually used in conventional analyses, and it has an unusually high accuracy, according to Dr. Larson.

The Activation Analysis Group of the Laboratory's Analytical Chemistry Divison, which has done much of the work of developing the technique, is conducting the analytical operations of the service, under the supervision of George W. Leddicotte. Its commercial aspects are being handled by the Radioisotope Control Department of the Laboratory's Operations Division, in a manner similar to the raidoisotope plan.

Potato Leafhopper Control

One application of toxaphene spray on second growth alfalfa early



in June protected the crop from the potato leafhopper on the farm of Ivan G. Martin, Ephrata, Pennsylvania. In the picture, the sample on the left was taken from a treated portion of the field, while the stunted clump on the right was taken from the untreated check plot only a few yards away.

Associate County Agent, Harry S. Sloat of Lancaster County, said that the insect, Empoasca fabae (Harr.) had done widespread damage to more than 100 unprotected fields in the area. In addition to stunting the growth, feeding of the insects on the undersides of the leaves produced a yellowing, withered effect. Bloom was also reduced considerably.

Lindane Seed Treatment for Beets

HE progressive sugar beet grower has long recognized the value of seed treatment with a fungicide for the control of damping-off organism which, without such control, would completely destroy many stands of seedlings on thousands of acres throughout many of the best producing areas of the United States. During recent years it has become increasingly apparent that many soil inhabiting pests, particularly certain species of wire worms, have become an additional major factor necessitating control in the establishment of satisfactory stands of sugar beets. Excellent results have been obtained by the use of Isotox Seed Treater (75% Lindane) in controlling limonius canus and other species of wireworms.

Three of the four large processors of sugar beets in California supply their growers with treated beet seed. They have standardized the type of treatment and the materials. The fungicide used is "Phygon" (2,3-dichloro · 1,4naphthoquinone) and the insecticide is "Isotox Seed Treater" (75% Lindane). The former is a paste, the later, wettable powder.

(Turn to page 127)

Black ELIABLE, DEPENDABLE PEST CONTROL PRODUCTS

the famous Black Leaf® trademark is the tried and trusted friend of millions of Americans. For more than 40 years it has been the symbol of quality and dependability to farmers, stockmen, fruit and vegetable growers, florists and home gardeners. Here are some old and some new members of the big family of Black Leaf pest control products:

Addrin: Black Leaf Aldrin Dusts are available in various formulations: 2½% Aldrin, 2½-5-0 (with 5% DDT), 2½-5-0 (with 5% DDT), 2½-5-0 (with 5% DDT and 40% Sulphur), and 2½-10-40 (with 10% DDT and 40% Sulphur). These dusts control boll weevils, boll-worms, leaf worms, thrips and similar insects infesting cotton. Black Leaf Aldrin Emulsifiable Concentrate is produced containing 2 pounds of Aldrin per gallon for spraying cotton to control boll weevils, thrips and grasshoppers.

Arsenate of Lead: Black Leaf Arsenate of Lead is a standard, high-quality Lead Arsenate for control of chewing insects in accordance with state recommendations.

Benzene Hexachloride: Black Leaf 3-5-40, containing 3% gamma isomer of Benzene Hexachloride, 5% DDT and 40% Sulphur is widely used on cotton for control of boll weevils, bollworms, thrips, aphids and similar insects. Also available without Sulphur (3-5-0), and with 10% DDT (3-10-40 or 3-10-0).

Calcium Arsenate: Black Leaf Calcium Arsenate is a standard product, especially suited for dusting cotton to control boll weevils and certain other insects, and for use wherever Calcium Arsenate is recommended for dusting or spraying.

Chlordane: Black Leaf 72% Chlordane Emulsifiable Concentrate is a high-quality product for control of ants, roaches, flies, or similar in sects... and also for control of soil-infesting insects, such as wireworms, Japanese beetle larvae, etc. Other formulations of Chlordane, such as 45% emulsion, 40% dust base and 40% wettable powder, are available.

DDT: Black Leaf 50% DDT Wettable Powder is recommended for control of insects on fruits, vegetables, field crops, etc. Black Leaf 25% DDT Emulsifiable Concentrate is formulated to meet exacting requirements for control of bollworms on cotton, and is generally suitable for spraying crops for which DDT emulsifiable concentrates are recommended. Black Leaf 10% DDT and 5% DDT Dusts are widely used for the control of insects infesting vegetable and field crops and for

the control of cotton bollworms. Formulated on a special tobacco material carrier, Black Leaf 10% DDT Dust makes an excellent product for use on tobacco.

DDT/Parathion; Black Leaf 253 is a combination of 25% of DDT and 3% of Parathion, formulated on a special tobacco material carrier. This highly-effective product controls codling moth, leafhoppers, red-banded leafroller, certain scale insects and mites infesting apples, grapes and similar fruits. This combination offers a high degree of efficiency and economy when used in the second cover spray and subsequent applications as recommended by local authorities.

Dieldrin: Black Leaf Dieldrin Emulsifiable Concentrates are formulated especially for use on cotton, to control boll weevils, thrips and similar insects, and provide long-lasting effectiveness.

Nicotine Products (Nicotine Sulphate, Nicotine Alkaloid, and Fixed Nicotine): The famous Black Leaf 40 and Black Leaf 99 are the recommended forms of Nicotine for use in controlling aphids and similar soft-bodied sucking insects infesting fruits, vegetables and field crops, by spraying or dusting — and Black Leaf 155 (Fixed Nicotine) is the recommended form of Nicotine for spraying fruits to control codling moth and similar insects, its use being particularly desirable during the latter part of the season to avoid undesirable residues at harvest. Made by the world's largest manufacturers of Nicotine insecticides.

Parathion: Black Leaf 15% Parathion Wettable Powder (Dustless) is formulated on a special tobacco material carrier to eliminate the hazardous dustiness of this important new insecticide. The Dustless feature is an extra safeguard.

Sulphur Dusts: Black Leaf Dusts, combining Sulphur, Copper and DDT, are recommended for control of leafspot, leafhoppers and certain other insects and diseases attacking peanuts.

TEPP: Black Leaf Vapo-Fume® 40, containing 40% of tetraethyl pyrophosphate, has special utility in the control of mites and certain other insects. Conveniently packed in 1-gallon containers — and in larger containers for remanufacturing purposes. This product is quick-acting and highly effective. leaving no undesirable residues. Precautions should be observed in handling, as with all phosphatic insecticide materials.

Toxaphene: Black Leaf Toxaphene Emulsifiable Concentrates — widely used to control bollworms, boll weevils and similar insects infesting cotton — are formulated to contain 4, 6, or 8 pounds of Toxaphene per gallon. Also used for controlling spittle bugs, grasshoppers and other insects which require extra killing power. Other Black Leaf Toxaphene products are: 40% wettable powder suitable for spraying; 20% dusts (with and without Sulphur); and 40% concentrate for manufacturing dusts.

Weed Killers: Black Leaf 2,4-D Amine Weed Killer and Black Leaf 2,4-D Ester Weed Killer are available in Midwestern and Northwestern grain-producing areas for control of noxious weeds and certain other plants infesting grain crops.

For House and Garden Use: In addition to the products listed above, the Black Leaf family includes pest control products conveniently packaged for house and garden use, such as: Black Leaf 40%, Black Leaf Acrosol Insect Killer, Black Leaf Arsenate of Lead, Black Leaf 1.00% Rotenone Dust, Black Leaf 50% DDT Wettable Powder, Black Leaf Garden Dust, Black Leaf Warfarin Rat and Mouse Killer Concentrate and Black Leaf Warfarin Rat and Mouse Killer Mixed Bait.

Every Black Leaf pest control product is attractively packaged and carries full directions. If you desire additional information, communicate with the address below. Your inquiry will receive prompt attention.

TOBACCO BY-PRODUCTS & CHEMICAL CORPORATION

Richmond, Va. . Louisville, Ky. . Montgomery, Ala. . Waco, Texas . San Francisco, Calif.

Ohio Pesticide Meeting Brings Information on

CHEMICAL CONTROL of PESTS

HE annual pesticide tour of The Ohio Pesticide Institute, held August 13, 14 at the Ohio Agricultural Experimental Station, Wooster. Ohio, featured a number of experiments dealing with comparisons of various pesticidal materials in different formulations; spraying techniques, including variations in gallenage, nozzle types, pressure variations, etc; and crop lines exhibiting differing degrees of resistance to diseases and insects. More than 100 persons attended the two-day session, which began with an address of welcome by the Dean and Director of the station, L. L. Rummel.

In discussing experiments with thrips control on roses, and insect and mite control on dahlias, R. B. Neiswander, of the experiment station, indicated that insecticides are more effective if applied to foliage instead of the soil, since the pests are then contacted directly in feeding. It was suggested that DDT should be used in all rose formulations, even though it magnifies the mite problem. The latter are effectively controlled with parathion, although this insecticide was not recommended for the home gardener, due to the toxicity hazard. It was reported that DDT seems to encourage the growth of dahlias, and parathion was found particularly good for aphis control.

A series of eleven different fungicide insecticide formulations were applied to cucumbers for the control of insects and diseases. The same formulations used on cucumbers at Marietta earlier in the season controlled anthracnose of cucumber in the following descending order: (1) "Dithane Z-78" and purified DDT: (2) "Manzate" and DDT-P: (3) "Zerlate" and DDT-P: (4) "Vancide 51ZW" and DDT-P:

(5) "T.ibasic" and rotenone; (6) "Crag 658" and methoxychlor; (7) "Crag 658" and DDT-P; (8) "Crag" and "Malathon WP." The control plants were found to have 18 per cent of the foliage remaining at the end of the experiment, whereas "Dihane" (that is treatment 1), treated plants had 65% foliage remaining. Methoxychlor applications on cucum-





Top photo. Officers of Ohio group: lat vice-pres., D. L. Kent, Goodrich Chemical Co., Cleveland; secretary, J. D. Wilson, Ohio Experiment Station; second vice-president, D. Zimmerman, Diamond Fertilizer Co., Sandusky, O.; treasurer, M. G. Farleman, Standard Oil Co. of Ohio; and president, H. E. Bruner, Monsanto Chemical Co.

Center photo: (Kneeling) Dean L. L., Rummel, R. B. Neiswander, C. R. Neiswander, Frank Irona. Standing: H. F. Winter, J. D. Wilson, C. R. Cutright, and H. C. Young.

Lower Photo: Sprayer being demonstrated in Ohio orchard.



NITROGANIC TANKAGE

SOIL & FERTILIZER CONDITIONING

STON

An activated sludge used as a raw material in the manufacture of balanced fertilizers. It aids in mixing and blending and adds organic nitrogen, available phosphoric acid, and humus to the finished fertilizer. Shipped in bulk or bags. Write today for full details.





SOLE DISTRIBUTOR:

H.J. BAKER & BRO.

AGO FIFTH AVENUE . HEW YORK 20, N. Y.

bers did not kill aphis, and it was recommended that parathion be used against this pest.

Results of a series of 16 different soil additives applied to test plots which were then analyzed for disease incidence, are still under anaysis; however, of the compounds tested, "Dithane Z-78" and a complete "Frit" formulation were found to give somewhat better results than the other treatments.

Studies of disease and insect resistance in potatoes indicated that some varieties were resistant to late blight and to insects; other varieties were found resistant to leaf hoppers, flea beetles, and late blight. The Russet Rural variety was found resistant to flea beetles, and this resistance is attributed to the taste of the leaf; still another factor affecting resistance is surface pubescence.

Another study made on potatoes was one dealing with the influence of variations in the spray schedule on maturity date and quality of potatoes. Eight different programs of bordeaux, DDT, and "Dithane D-14" applications, with the crop to be harvested at four different dates, are currently underway in a study of the effect of degree of maturity at digging on resultant chipping and storage quality and on specific gravity. In this experiment, the vines will be killed or eliminated by different methods before digging in a study of the effect of vine killers on vascular staining and keeping quality.

"Systox" applications on potatoes to control leaf hoppers and flea beetles were found inadequate if used for one treatment; however two treatments did give control, although the period of control is still under determination. The major problem in this experiment is the amount of chemical residue in the tubers.

AHIGHLIGHT of the tour was the demonstration of a new experimental sprayer, designed and built by the USDA Engineering Laboratory at Toledo. The unit was mounted on a Massey-Harris Pony Tractor, and featured power take-off at the front. It included quickly adjustable

and removable boom assemblies, and booms equipped with either flat spray nozzles or hollow cone type in 4,3,2, and 1 nozzles for spraying a 51/2-foot row (1 row of tomatoes or 2 rows of potatoes). Of considerable interest is the by-pass arrangement of the spray solution piping, resulting in a siphoning effect when the power is shut off . . . the insecticide formulation is by-passed back to the tank, preventing the clogging of nozzles by the spray compound. The unit permits the use of low gallonage applications and provides accurate control of speed and rate.

Another sprayer demonstrated at the experiment station, was the Myers air-blast sprayer designed to spray row crops or fruit trees. This sprayer could be adjusted to spray a 35' ridge of tomato or potato plants. It has been tested with favor-

able results for the control of most vegetable diseases. In the orchard, the concentrate was observed to penetrate the tree, wetting all branches. The sprayer uses a pressure of about 400 psi, and eight standard hollow cone nozzles for spraying trees.

A program applying nine different insecticide formulations to nine vegetable crops, to determine effectiveness in controlling various insects; the effect on flavor of the harvested crop, and the amount of residue in the edible part of the plant is currently under study at the Ohio Experiment Station. Table 1 summarizes some of the results obtained. (all insecticides were combined with dry "Parzate" at 2/100).

All of the insecticide formulations, with the exception of dieldrin and methoxychlor, gave good control

(Turn to Page 118)

TABLE 1

Insecticide Formulation	lbs./gal.	flea beetle damage/leaf	Bean beetle damage	Cabbage worm
Malathon 25% WP	2-100	20-21	Light	Med.
Malathon 50 3% Em E-20	1-100	20-21	Light	Med.
Malathon 55% Em E-21X	1-100	20-21	Light	Med.
Systox	11/2-100	16	Light	Heavy
NPD 85% Em	1-100	19	Heavy	Heavy
Parathion 15% WP	11/2-100	21	Light	Light
G-2311 25% soln.	1-100	21	Light	Heavy
Dieldrin 25% WP	2-100	4	Heavy	Light
Methoxychlor 50% WP	2-100	22	Light	Med.
Untreated Check		35	Heavy	Heavy

TABLE 2

Fungicide	Amt./100	% Scob on fruit	% Russel
Magnetic — 70 sulfur	8.6 lbs.	5.5	12
Tag fungicide	½ pt.	0	7.5
Puratized ag. spray	1 pt.	1.0	9.5
Coromerc	1/2 lb.	1.0	5.5
Organic mercury A	1 pt.	.5	22.5
Organic mercury C	1 lb.	1.0	23.5
Manzate special	2 lb.	9.5	40.0
Check - no fungicide	_	96.0	-

TABLE 3

% living aphis at various date. Treatment after treatment				s dates
	6/16	6/18	6/27	6/30
	Before treatme	ent		
Ukako	100	101	93	67
Check	100	107	92	50
SP Systox 4 oz./100	100	0	0	-

PRENTOX PEST-TESTED

basic insecticide concentrates

DDT

WETTABLE AND DRY POWDERS OIL CONCENTRATES EMULSIPIABLE CONCENTRATES

CHLORDANE

OIL CONCENTRATES
WETTABLE AND DRY POWDERS
EMULSIFIABLE CONCENTRATES

ROTENONE

DERRIS AND CUBE POWDERED CONCENTRATES

SABADILLA

GROUND SEED DUST CONCENTRATES

TOXAPHENE

WETTABLE POWDERS DUST CONCENTRATES

LINDANE

WETTABLE AND DRY POWDERS OIL CONCENTRATES EMULSIFIABLE CONCENTRATES

RED SQUILL

FORTIFIED POWDER FORTIFIED LIQUID

PYRETHRUM

POWDER NO. 20 EXTRACT

RAX POWDER

A NEW RODENTICIDE CONTAINING WARFARIN



HERE'S A PERFECT PRENTISS RAT KILLING COMBINATION

RAX · PRENTISS WARFARIN

DE-RATTER-MOUSER

PERMANENT BAIT STATION

SEND FOR LITERATURE ON DE-RATTER-MOUSER

PRENTISS DRUG & CHEMICAL CO., Inc.

110 WILLIAM STREET, NEW YORK 7, N. Y. 9 SO. CLINTON STREET, CHICAGO 6, ILL.

National Shade Tree Conference Devotes Much of Meeting to Study of

Chemical Control of Tree Pests

HEMICAL control of insects and plant diseases affecting trees was an important part of the program of the National Shade Tree Conference August 18-22. The 28th meeting of the group was held at the Statler Hotel, Boston, Mass., with one of the largest crowds in the organization's history.

Discussions on the control of birch leaf miner, roadside brush control, chemotherapy, soil conditioners and fungicidal control of diseases were all prominent on the program which also included talks on training arborists, business outlook for the industry and mechanical means of tree surgery.

Dr. W. P. Martin, Department of Agronomy, Ohio State University, Columbus, Ohio, discussed soil conditioners in general and "Krilium" in particular. He said that "Krilium" has been under test at Ohio State since 1950, using different soils, rates of application, varying crops and under other conditions. Results indicate that from 400 to 2,000 pounds per acre (0.02-0.1%) added in powder form to tight and heavytexture (clay) field soils, with subsequent mixing to a depth of 6 inches by disking, have "without exception" changed the clay into soils of good friable crumb structure. Soil granulation often exceeded 80%, after such an application, he said. Moreover, the soil aggregates are water-stable and the conditioning chemicals themselves are resistant to decomposition.

Structural improvement continued into the third year, but not at the original levels.

Dr. Martin emphasized the importance of loosening up the soil and mixing the conditioning chemical in the soil when applying. The soil-aggregating agents themselves cannot loosen up the soil. They bond to the first soil particles they contact and stabilize the condition produced by the mechanical working of the soil. Attempts to improve soil structure with water solutions of conditioning chemicals by sprinkling and without first mixing, have thus far been in-effective, he said.

Good soil tilth has proved beneficial to many plants tested in statistically · controlled experiments, the speaker stated, but where soil structure was not a critical factor in production, no yield inceases were noted. In general, soil aeration improved, favoring growth for plant roots and activity of soil oganisms. Treated soils drain faster in spring rains, and rain water infiltrates rapidly with less runoff and erosion. Cultivation of conditioned soils is also easier and seedlings emerge more quickly because the soil is not crusted, it was pointed out.

For erosion control, the speaker declared, as little as a half pound of "Krilium" dusted on 100 square feet has formed a surface film and has stabilized the soil against raindrop impact and running water.

Mr. Martin emphasized, how-

ever, that soil conditioners do not contain plant nutrients. "They influence plant growth only as they improve soil structure and tilth," he declared.

Birch Leaf Miner Control

Pointing out that the birch leaf miner is a pest found in all parts of the country where birch trees grow, Dr. William W. Cantelo, Bartlett Tree Research Laboratories, Stamford, Conn., told the group of various methods used in controlling the pest.

DDT and parathion have proved effective in laboratory tests in preventing oviposition, but DDT was ineffective in the field. Single sprays of summer oil, DDT oil emulsion and nicotine sulfate were found to be ineffective against the egg state, he

Chlordane (1.6 lbs.); lindane (0.25 lbs.); aldrin (0.25.); dieldrin (0.25 lbs.) and Toxaphene (1.2 lbs.) each in 100 gallons of water, gave excellent control in both egg and larval stages. Nicotine sulfate was effective if applied in 2 C 3 sprays. Addition of sodium salicylate had little effect on the amount of control, he said.

Lindane was effective against all immature stages except the 4th instar larvae and prepupae. Lindane and toxaphene applied to the soil in the fall, killed a quarter of overwintering larvae. Sprays on the lower

(Turn to Page 114)



This is good news in agriculture and in industry. Increased availability of vital plant nutrients in the form of soluble inorganic salts for fertilizer solutions can mean new products and new markets for fertilizer manufacturers.

New emphasis is being placed upon water-soluble fertilizers. Growers and manufacturers are developing new methods... new applications of high-analysis soluble plant foods which combine NTTROGEN...PHOSPHORUS, POTASH and, in many cases, weed killers and insecticides, too.

MONSANTO, to help meet the increasing demands for plant foods, has increased quantities of four basic, soluble fertilizer chemicals... MONO AMMONIUM PHOSPHATE... DI AMMONIUM PHOSPHATE... MONO POTASSIUM PHOSPHATE... PHOSPHORIC ACID 75.0%.

New uses for WATER-SOLUBLE FERTILIZERS are proving profitable for growers... profitable for FERTILIZER MAN-UFACTURERS. Perhaps these products will fit into your production planning. Contact any District Sales Office, or write MONSANTO CHEMICAL COMPANY, Phosphate Division, 1700-B South Second Street, St. Louis 4, Mo.

DISTRICT SALES OFFICES: Birmingham, Boston, Charlotte, Chicago, Cincinnati, Cleveland, Detroit, Los Angeles, New York, Philadelphia, Portland, Ore., San Francisco, Seattle. In Canada, Monsanto Canada Limited, Montreal.

MONSANTO PLANT NU	TRIENT CHE	MICALS	
	N	Pz0s	K20
Mono Potassium Phosphate (Crystals)	-0-	51.6%	34.2%
Di Ammonium Phosphate (Crystals)	21.0%	53.85%	-0-
Mono Ammonium Phosphate (Crystals)	12.2%	61.61%	-0-
Phosphoric Acid (75.0%) (Liquid)	-0-	54.5%	-0-

Here are ways you can help him deliver the goods in the coming election

This worker's future—and yours—depends on how well he understands the importance of his vote and the importance of voting wisely. Too many of us fail to vote, to choose our leaders wisely, to guide them in the direction we want to go. You can help workers in your own business vote effectively by showing them how their votes, intelligently cast, can build a better America.

HERE ARE THE TOOLS-USE ALL OF THEM ...



1. "KERRY DRAKE IN 'THE CASE OF THE SLEEP-ING CITY'"—To help give people a better understanding of their individual vote and what it can mean to community growth, good government and prosperity. This colorful new 16-page booklet follows the time-proven "comic book" technique...dramatizes the relationship between civic responsibility and the ballot.

THIS BOOKLET HAS BEEN TESTED in our own plants by the Psychological Corporation . . . and prozen effective. Because good government is everyone's business, Bemis is making "THE CASE OF THE SLEEPING CITY" available to you at nominal cost for distribution to your people.

2. "YOUR VOTE IS VITAL"—A pictorial booklet showing the factors that help a citizen become

Because we believe these messages, published by Harvey Publications, are growing more urgent every day, Bemis is taking this means to recommend them to every business. Permission to reproduce this advertisement in association magazines, company magazines, etc., will be granted upon request. Plates will be sent without charge. BEMIS BNO. BAG CO., St. Louis 2, Missouri.




an intelligent voter—how to understand political issues; how to compare candidates promises with their records; how to study local issues; how to register; how to vote by machine or by ballot, etc. A useful booklet that will help your employees.

3. "THE MAN WHO WOULDN'T QUIT"—Another tested-and-proved booklet, in comic book technique, which spells out every man's personal responsibility for good government, Announced a few weeks ago, this booklet has already had wide acceptance. It will help YOUR employees.

For samples and complete information, without obligation, mail this coupon now!

HARVEY PUBLICATIONS, INC. 1860 Broadway, New York 18, N. Y.
I'm interested in "The Case of the Steeping City"; "YOUR Vote is VITAL" "The Man Who Wouldn't Quit" Send samples and quantity cost information.
Your Name
Firm
Address
400 T 41.1



LINDANE

The following is a comlete list of products available from KOLKER CHEMICAL WORKS, DIAMOND AL-KALI'S subsidiary, specializing in organic chemicals for agriculture and industry:

DDT. Technical Grade
BHC. Technical Grade
345° and 99°s gammaLINDANE. 99°s gamma Isomer
2.4-D. Weed Killera
2.4.5-T. Brush Killera
K-101. Mitteide


KOLKER CHEMICAL WORKS INC.

80 LISTER AVENUE, NEWARK S. NTW JESSEY Plants: Newark, New Jessey and Houston, Texas



LINDANE, the effective multi-purpose insecticide containing 99% gamma isomer of benzene hexachloride is now available in commercial quantities. LINDANE, together with KOLKER high gamma BHC technical, now offer manufacturers basic BHC materials easily handled and readily formulated into high-quality finished insecticides.

Chemicals you live by



How about you? If you manufacture finished insecticides, there are many advantages for you in buying directly from Pennsalt — a basic producer of chemicals.

PENNSALT offers you a single, dependable source for your technical materials and concentrates. All PENNSALT products are consistently dependable—Pennsalt's own research and testing facilities provide quality control from start to finish.

And PENNSALT agricultural chemicals offer you extra compounding benefits, too!

Here are two good examples:

PENTECH, the unique friable DDT; made by

Pennsalt's patented process for ease of blending and reduced cost of processing.

PENNSALT BHC TECHNICAL 36, with 36% gamma isomer, proportionately less of the other isomers. This means you handle and store up to $\frac{3}{2}$ less of the technical materials.

We'll be glad to send you complete information on prices, delivery and Pennsalt technical assistance; just write to the office nearest you.

Agricultural Chemicals Department

PENNSYLVANIA SALT MANUFACTURING COMPANY

Philadelphia 7, Pa. • Tacoma, Wash Bryan, Texas • Montgomery, Ala. Los Angeles and Berkeley, Calif. • Portland, Orc.

Some Pennsalt Technical Materials & Concentrates:
PENTECH. DDT • PENNSALT BHC TECHNICAL 36 • KRYOCIDE.
(natural cryolite) • HI-GAM. 99 (lindane) • DB-50
(50% DDT dust base) • BHC D-12 (12% gamma dust base)
• BHC-DDT 9:15 (dust concentrate)



PROGRESSIVE CHEMISTRY FOR OVER A CENTURY





EMCOL H-77

the one emulsifier that is still taking that extra load

AFTER FOUR YEARS of success in the field, easy-to-handle, versatile, low-cost EMCOL H-77 is still the one single emulsifier that carries the most weight with pesticide formulators.

Featuring outstanding "carryover" stability with required emulsifiability, EMCOL H-77 does the best job with the greatest number of toxicant materials.

Write for the details in our Technical Bulletin No. 31

VIGILANT SERVICE

THE EMULSOL CORPORATION

59 East Madisan Street

Chicago 3. Illinois

FLEXSPOUT

Helps Sell More of any Product Sold in Steel Pails

BECAUSE CUSTOM ES PREFER THESE FLEXSPOUT FEATURES



LEAKPROOF—One-piece molded plastic, sealed directly on pail metal. Stronger than pail itself.



EASY 10 OPEN—Without special roofs, bail, screwdriver or any share object does the job.



FLEXIBLE POURING SPOUT— Extends for easy pouring. Ne muss or fuss . . . because only inside of spout touches liquid.



RE-SEALS TIGHTLY—With plastic screw cap. Protects, volutile liquids. Sout retracts for easy stantia.

RIEKE METAL PRODUCTS CORPORATION
AUBURN, INDIANA

FLEXSPOUT simplifies using, pouring and storing your liquid products. It's a powerful sales tool for any liquid that's packaged in steel pails.

FLEXSPOUT is odorless, tasteless and nontoxic. It resists attack by almost all chemicals, solvents and oils. Easily applied with one stroke of a simple hand tool, it permits faster filling and sealing, lowers production costs. For a few cents a pail it adds dollars worth of sales appeal to your products.

Almost all pail manufacturers are equipped to furnish pails ready for use with the **FLEXSPOUT** closure. For more details, prices and a free sample, just fill out and mail the handy coupon.

-MAIL THIS COUPON TODAY-

RIEKE METAL PRODUCTS CORPORATION AUBURN, INDIANA

We'd like to know more about the new FLEXSPOUT Closure.

Please send us...

| Free Sample | Brochure |

 Reliability . . . an ingredient of our agricultural chemicals

since 1911

To make certain you buy the best and get the best buy for your particular needs, we maintain our own laboratory located near the source of raw materials. Reliable service must be provided with reliable products and we help to insure this for you with conveniently located branch offices.



TOXAPHENE 40% • BHC • ALDRIN • PARATHION • DDT • SULPHU RDANE . DIELDRIN A Complete Line of Diluents: Talc, Pyrophyllite, Clay, Fuller's Earth

ASHCRAFT-WILKINSON CO.

Home Office ATLANTA, GEORGIA Cable Address ASHCRAFT

. CHARLESTON, S. C. . TAMPA, FLA. . COLUMBUS, OHIO

THINKING ABOUT BHC?

For these and many other insecticides, The Harte System can provide the know-how . . . from planning to profit. Why do it the hard way?


LINDANE?


Write for our booklet describing how the Harte System can help you.

DESIGNERS • ENGINEERS • CONSTRUCTORS

John J. Harte Co.

ATLANTA . NEW YORK HOUSTON . MEXICO, D. F.





PYRAX ABB GIVES YOU DENSER CLOUDS, BETTER COVERAGE

Do you know why Pyrophyllite has become the fastest growing agricultural diluent in the last twelve years? . and why more Pyrax ABB is used than all other Pyrophyllite?

First, Pyrax ABB is extremely well adapted to aircraft dusting because it forms dense clouds that hug the crop . no waste from drifting clouds that settle on adjoining fields.

Second, Pyrax ABB adheres electrostatically, even to the dry underside of foliage you can dust anytime without waiting for proper moisture conditions.

R. T. VANDERBILT CO.

Specialties Department

230 Park Avenue



New York 17, N. Y.

Third, because Pyrax ABB is nonhygroscopic and chemically inert (pH between 6.5 and 7.0), it will not affect the toxic strength of botanicals, DDT, benzene hexachloride, toxaphene, thiocarbamates, or the newer organic toxicants . . . therefore you can use less toxicant and even store these dusts from season to season without affecting their killing power.

If you need a free-flowing, non-absorptive, neutral, and nonhygroscopic agricultural dust diluent that is always in dependable supply, write today for all the reasons why you should be using Pyrax ABB.

	Please	send	technical	information	on
_	PYRA	X Al	BB.		

☐ Please send sample of PYRAX ABB.

NAME

POSITION

(Please attach to, or write on, your company letterhead)

HERE'S THE ANSWER TO YOUR SPREADING PROBLEMS!

Commercial Fertilizer Attachment Hood—Two Sizes

19-foot or 23-feet



with its new
"Motor-Driven Spreader"
offering greater accuracy of spread
with the most positive feed on the market!

SPECIAL ADVANTAGES — Uniformity of spread is not dependent on truck speed. Motor is mounted on catwalk and drives only the twin distributor discs at a constant speed, assuring full width of spread at all times together with uniform distribution.

Conveyor is separately driven from truck drive shaft by a series of V-belts to deliver the correct amount per acre—regardless of truck speed or regardless of whether the truck is driven in low, super-low or any other gear.

Conveyor speed is, therefore, positively syn-

chronized with speed of the rear wheels of the truck and at each revolution of the rear wheels, the conveyor moves a given distance regardless of the truck's speed. Amount of material delivered by conveyor does not vary with hilly or soft field conditions.

Spreader Body Lengths (inside measure) are 9', 11', 13' and 15'. Other body lengths on special order.

Note: When Spreading Attachment is folded up for road-traveling position, width is approximately 7'-5'.



"The NEW LEADER" Self Unloading Bulk Transport

The 20-ton capacity transport above is shown with elevator in place and ready to load a NEW LEADER Spreader truck. These units are proving very profitable; in bad weather they eliminate demurrage on railroad cars; fertilizer gets to the job quickly and spreader trucks can be kept working in the field. The transport, being a self-unloading unit, leaves the tractor truck free to return to pick up another transport load. These

units have four individual compartments of 5 tons each. Each compartment may be unloaded independently of the others. Compartments and rear endgate are removable so that bagged and packaged goods may be hauled instead of bulk loads. Capacity 5 tons to 25 tons, lengths from 11 ft. to 40 ft. Written warranty with all NEW LEADER equipment. Write today for specifications, prices, etc. Fast delivery service sells fertilizer!

FREE! Write for "The Story of a Custom Fertilizer Spreading Service"

HIGHWAY EQUIPMENT COMPANY, INC. CEDAR RAPIDS, IOWA MANUFACTURERS OF THE WORLD'S MOST COMPLETE LINE OF SPREADERS



"fruit salad" has no place in functional packaging

In the merchandising of dry chemicals, as in the design of ladies' hats, it is easily possible to have more packaging than you need.

Over-packaging is expensive any time. Today, more so than ever. As costs rise, it is an increasing drain on profits.

For packages up to twenty-five lbs., many manufacturers of dry chemicals are rediscovering paper bags.

Union CHEM-PAK bags inexpensive-

ly provide the kinds of protection most chemical products require. They help sales too. Eye-catchers, they invite featured shelf display.

Look into sift-proof CHEM-PAK new horizon in chemical packaging.

CHEM-PAK BAGS COST LESS ALL ALONG THE LINE


- Lower First Cost · Freight Savings
- · Less Handling · Fewer Losses
 - · Faster Shelf Turnover

CHEM-PAK

SIFT-PROOF PAPER BAGS FOR CHEMICAL PRODUCTS

UNION BAG






NEW! Mail this coupon for proctical CHEMICAL PACKAGE EVALUATOR. No charge or oblig Union Bag & Paper Corporation Woolworth Building, New York 7 Please send me the new Chemical Package Evalua City

Hudson Falls, N. Y. Billion-bog

CHEM PAR BASS FOR FERTILIZERS—INSECTICIDES, FUNGICIDES, BUSTING POW-BERS—PLANT FOODS—WALL SIZE—PASTE FLOUR—PATCHING PLASTER—PRE-PARED CEMENT-PLASTER OF PARIS-PHARMACEUTICAL CHEMICALS-BYES AND PIGMENTS - SANITARY CHEMICALS - POWDER SHAMPOOS - CHLORINE COMPOUNDS AND ALRALIDES - CAULKING, SEALING, WHITING AND FILLING COMPOUNDS - AND OTHER DRY CHEMICALS WITH COMPARABLE PACKAGING REQUIREMENTS It's a fundamental fact...

SEQUESTRENE

puts the Gille back into crops



and keeps it there for nine months or longer



Mineral deficiencies in crops have been with us like the unsolved crime down at the police precinct — only longer. Now, for the first time, research presents a bona fide lead toward solving the fundamental problem of how to enable crops to assimilate minerals in which they are deficient and thereby promote better growth and bigger yields.

Sequestrene* iron complexes have corrected one of the most difficult mineral deficiencies to cure, iron chlorosis. Field tests employing formulations of Sequestrene and ferrous sulphate on citrus have transformed yellow leaves to hrilliant green and have infused a new rich growth in crops that were dying.

These changes were accomplished within four to six weeks after application and lasted for nine months and longer. The end-result is expected to be bigger yields of better quality.

Many research agencies are continuing to investigate the entire realm of mineral deficiencies including copper, zinc, cobalt, manganese, magnesium and other essential trace elements.

Those interested in the origin of the present work on mineral deficiencies should consult the publications of Stewart and Leonard (1) and Jacobson (2).

Geigy Company, Inc., Insecticide Division, are the exclusive sales agenta for Sequestrene in the agricultural field. Write for technical bulletin, "Metal Complexes of Sequestrene" in Plant Nutrition", and pamphlets on iron chlorosis on citrus and reprints of papers by Stewart and Leonard, and Jacobson.

ORIGINATORS OF

GEIGY COMPANY, INC.



2301211232NI TOO

89 BARCLAY STREET N. Y. 8, N. Y.

Aberdeen, N. C.; Burlington, lawa; Colorado Springs, Colo.; Elkron, Md.; Fresno, Calif.; Houlton, Me.; Leland, Miss.; McGregor, Tex.; Orlando, Fla.; Walla Walla, Wash.

^{*&}quot;Sequestrene" is the registered trademark of the product of Alrose Chemical Co., A Geigy Company.

⁽¹⁾ Stewart, Ivan and Leonard, C. D. Iron Chlorosis — Its possible causes and control. Citres Magazine 14(10): 22:25, 1952. (2) Jacobson, Louis. Maintenance of Iron Supply in Nutrient Solutions by a Single Addition of Ferric Potassium Ethylenediamine Tetraacetate. Plant Physiol. 26(2):411-413, 1951.

INDUSTRY NEWS

Shell Appoints Selden

J. M. Selden has been appointed manager of the Eastern Division of Shell Chemical Corporation,



I. M. SELDEN

it has been announced by J. Oostermeyer, president.

A native of Pittsburgh, Pennsylvania, and a well-known personality in the chemical trade, Mr. Selden has been associated with the marketing of Shell Chemical products since 1933, first as vice-president and director of R. W. Greeff & Co., agents until 1946 for the sale of Shell Chemical products east of the Rockies, and then as sales manager of the chemical firm's eastern division, created in 1946 along with five subsidiary district sales offices to meet growing demand.

In his new position as manager, eastern division, Mr. Selden will continue to administer sales activities as well as assuming a chief part in the introduction to eastern markets, of new Shell Chemical products still under development.

Plans to Market Mulch

The Pacific Lumber Co., San Francisco, is projecting plans for further development of redwood by-products as ingredients of commercial fertilizers and insecticides. Under the trade name of "Palco Pete," the firm has for many years marketed a red-

wood bark mulch, and is currently sponsoring development work with organic chemical by products as fertilizer and insecticide components.

Wm. G. Van Beckum has just been named director of research and development for Pacific Lumber.

Since 1948, Mr. Van Beckum has been manager of technical service and assistant sales manager, Special Products Division, Weyerhaeuser Timber Co. of Longview, Washington.

The Pacific Lumber Company has pioneered research in utilization of bark and other wood raw materials. Mr. Van Beckum will make his headquarters at the firm's San Francisco office.

Atlas Opens S. W. Office

The opening of a new office in Houston, Texas, to provide additional service for users of its industrial chemical products has been announced by the Atlas Powder Company of Wilmington, Del.

George J. King, director of sales of the company's Industrial Chemicals Department, said that the new office will make Atlas technical service more readily available to inindustries in Texas, Louisiana, Colorado, New Mexico and parts of Mississippi

Allen V. Riley, Jr. had been assigned as technical sales representative in this office. He is a graduate of the University of Minnesota and a veteran of World War II, and has been with Atlas since 1948. Prior to this assignment he was attached to the firm's Chicago sales office.

Mr. King explained the move as the latest in a series of steps planned to expand the services of the department, which manufactures and markets surface active agents, plasticizers and other products used by a variety of industries.

Gidney U. S. Potash V-P

Dean R. Gidney has been appointed vice-president in charge of the sales department of United States



DEAN R. GIDNEY

Potash Company, according to an announcement by Horace M. Albright, president and general manager, speaking for the board of directors of the company.

Mr. Gidney, 36, has been in the sales department of the company since 1938, except when on a long leave of absence for war service in the Navy. He was graduated in 1936 from Dartmouth College where he was elected to Phi Beta Kappa. For one year he was employed by the United States Trust Company in New York and in 1937 he joined the United States Potash Company's New York office.

Mr. Gidney, who held a reserve commission in the United States Navy as Ensign, was called to active duty early in 1941. He served in the North and South Atlantic and for two years in Europe with the landing craft staff preparatory to and following the invasion of France. He went on inactive duty as Lieutenant Commander in January, 1946, and rejoined United States Potash. He was advanced to sales manager in 1950. As vice-president he succeeds J. E. Barnes who retired because of ill health several months ago.

Monsanto Opens New Laboratory



Establishment of an agricultural and biological research installation at Creve Coeur, Mo., has been announced by the Organic Chemicals Division of Monsanto Chemical Company.

The facilities, to be known as the "Creve Coeur Laboratories," are in addition to the division's 257-acre Hazelwood Farms established earlier this year to field-test agricultural chemicals. Operations on Hazelwood Farms will be handled by personnel of the Creve Coeur Laboratories. Initially, the staff at Creve Coeur Laboratories will consist of three research groups under the direction of L. E. Weeks, P. P. Wallace and Dr. O. DeGarmo. The groups will engage in research on soil conditioner applications; herbicides, insecticides and fungicides; and mycology, bacteriology and industrial preservation, respectively.

In addition to the laboratories proper, the installation includes 10,-000 square feet of greenhouses for screening and experimental work. Bourbeau, who is presently on assignment with the Economic Cooperation Administration in the Belgian Congo.

Study Peach Canker in N. J.

Appointment of D. Fred Cohoon to a graduate fellowship for research on peach canker has been announced by Dr. W. H. Martin, director of the N. J. Agricultural Experiment Station, New Brunswick.

Mr. Cohoon is a graduate of the University of Western Ontario, London, Ontario, and has studied the use of antibiotics in controlling plant diseases. He will work under the direction of Dr. R. H. Daines, of the Plant Pathology Department.

Davison Appoints Clark

Chester F. Hockley, president of the Davison Chemical Corporation, Baltimore, has announced the appointment of Raymond S. Clark as counsel for the Company. Mr. Clark is a graduate of Groton, Harvard College and Yale Law School and was engaged in private law practice before joining Davison in June of last year.

Cyanamid Names Lapean

Raymond E. Lapean, executive vice president of Chemical Construction Corporation, was named managing director of Cyanamid Products Ltd., according to an announcement by American Cyanamid Co., parent firm. Mr. Lapean has been with Chemical Construction since 1940, when he became special assistant to the president. He was elected vice-president in 1944 and executive vice-president in 1950.

UCC Appoints Ross

J. W. Ross has been appointed Philadelphia district manager for Carbide and Carbon Chemicals Company, a Division of Union Carbide and Carbon Corporation, it has been announced by W. F. Reich, Jr., vice-president. Mr. Ross, who joined UCC in 1936, came to Philadelphia from Atlanta in 1944. He moves up to fill the position formerly held by R. M. Joslin, Jr., now midwestern division manager in Chicago.

New Indian DDT Plant

A new DDT plant is to be built in India with funds from the Indian government, the World Health Organization and the United Nations International Children's Emergency Fund, the WHO has announced. Signing of an agreement for building the plant in Delhi were completed late in July. Capacity of the plant is reported to be 700 tons per pear.

Although American producers of DDT are protesting, three additional DDT plants are being planned for erection in Pakistan, Ceylon and Egypt in addition to the one already agreed upon in India.

According to terms of the agreement, the Delhi plant is to be in production in March, 1954 and the total output will be devoted by the Indian government to malaria control programs. The agreement is said to contain provision for the training of persons from other nations in the manufacture of the insecticide.

Of the \$821,000 cost, the government of India is reported to contribute \$471,000; the U. N., \$250,000 and WHO, \$100,000.

UK Fertilizer Output Cut

A cut in the output of sulfate of ammonia at the Billingham-on-Tees plant of Imperial Chemical Industries in England, has been announced. The 25% reduction is attributed to a seasonal falling off, plus increased competition from abroad. However, the company's manager declares that the production cuts are not "panic measures," but rather a balancing of output with expected sales.

Adds to Soils Staff

Announcement has been made of the appointment of Tsuneo Tamura to the staff of the scils department of The Connecticut Agricultural Experiment Station, New Haven. He will be assistant soil scientist. Mr. Tamura fills the position left vacant by the resignation of Dr. Gerard A.

Becomes Asst. Sales Mgr.



JOHN CROWTHER

T. A. Haschke, sales manager, Stauffer Chemical Co., has announced the appointment of John Crowther as assistant sales manager as of August 1, 1952. Mr. Crowther has been associated with Stauffer for 7 years and was director of eastern division research, at Chauncey, N. Y. Mr. Crowther is a chemical engineer and a member of the American Institute of Chemical Engineers, Commercial Chemical Development Association and Chemists' Club. Before joining Stauffer, he was a Major in the U. S. Air Force, Headquarters, 8th Air Force in Europe.

Emerson Joins Stauffer

Stauffer Chemical Company has announced the appointment of Dr. Frank H. Emerson as technical advisor of agricultural sales. Dr. Emerson received both his Bachelors and Masters degrees in botany from the University of Kansas and was awarded his PhD from Cornell University for work done there in plant pathology. Before joining Stauffer Chemical Company, Dr. Emerson acted as research assistant at the New York Agricultural Experiment Station in Geneva, New York. His home office with Stauffer will be in New York.

New Monsanto Salesmen

Monsanto Chemical Company has announced the appointment of five territory salesmen for its new Merchandising Division.

Gordon Staub of Dearborn, Mich., and Donald D. Reichert of Cudahy, Wis., were assigned to the Chicago district, William R. Bone of Sullivan, Ill., was assigned to the St. Louis district, William P. Sheppard of St. Louis was assigned to the New England territory and Richard T. Bethel of Montclair, N. J., was assigned to the New York territory. The newly formed Merchandising Division is handling the sales of Krilium soil conditioner and other trade-marked products through retail channels.

Riis Succeeds J. H. Friend

The resignation of J. H. Friend, vice-president of International Paper Company and general manager of its Southern Kraft Division, was announced on August 12, by John H. Hinman, president. Mr. Friend will continue to serve as a member of the board of directors and will be available in a consultant capacity on special projects.

Succeeding Mr. Friend as general manager of the Southern Kraft Division will be Erling Riis, vicepresident and assistant general manager, Mr. Hinman said.

Mr. Friend, a native of Wisconsin, became associated in March, 1924, with the late R. J. Cullen and joined International Paper Company in 1927 when the Louisiana Pulp and Paper Company of Bastrop, Louisiana, was purchased.

Mr. Riis was born in Christiania, Norway, and received his education in that country, graduating as a mechanical engineer in 1909. He joined International Paper in 1927.

Stops Sale of "Loamium"

Sale of "Loamium" soil conditioner is illegal in the State of Virginia following a ban on the product by the state Department of Agriculture. Commissioner Parke Brinkley stated that the action was taken because "the product does not meet the requirements for fertilizer in Virginia."

H. J. Baker Exec. Dies

J. Bailey Pratt, Jr., sixty-one, a senior executive of H. J. Baker & Bro., import and export firm, New York, died at Manchester, Vermont on July 11th, 1952.

Mr. Pratt joined the Baker organization in 1923,

To Gen. Dyestuff Post



TOM R. MOORE

Tom R. Moore has been appointed assistant to the vice-president and general sales manager of the General Dyestuff Corporation, New York, it has been announced. Mr. Moore formerly was sales service manager for the Antara Chemicals Division of General Dyestuff.

A native of Roanoke, Va., Mr. Moore was educated at Virginia Polytechnic Institute and started his career in the chemical industry in 1935 with Sherwin-Williams. In 1941 he joined the National Carbon Company as a sales engineer and became associated with the Antara Division as sales engineer five years later. Mr. Moore was appointed sales service manager in 1950.

Gager Joins Kraft Bag

Kraft Bag Corporation, New York, manufacturers of heavy-duty multi-wall shipping sacks, has announced the appointment of James M. Gager, Jr. to its sales staff. He will cover Tennessee, Georgia, South Carolina and part of North Carolina, from headquarters in Chattanooga, Tenn. Mr. Gager has had experience in multi-wall sales and service, and has a practical knowledge of production and operation gained previously as a proprietor of The Gager Lime Company.

Davis Heads Association

H. C. Davis, manager of the Bemis Bro. Bag Co. paper specialty plant in St. Louis, was elected president of the Industrial Bag & Cover Association, and S. G. Yount, president of the Southland Paper & Converting Co. of Los Angeles, vice-president for the year ending in June 1953. our

B.G.C.

is no

fata morgana...



but **DEATH!**



BENZENE HEXACHLORIDE TECHNICAL

14°/_o gamma isomer BENZENE HEXACHLORIDE 12°/_o gamma isomer

74% other isomers

14% inert ingredients

BENZENE HEXACHLORIDE 33% gamma isomer BENZENE HEXACHLORIDE DUST CONCENTRATES

6 and 12 % gamma isomer BENZENE HEXACHLORIDE WETTABLE POWDERS

6 and 12% gamma isomer

other valuable chemicals: WEEDKILLER on base of M.C.P.A. MERCURY SEEDDRESSING

N.V. NOURY & VAN DER LANDE'S EXPLOITATIE MIJ.

P. O. BOX 10, DEVENTER, HOLLAND.

TELEPHONE 4441

Accepts NPA Position



GEORGE J. BRUYN

George J. Bruyn, manager of Niagara sales office of Hooker Electrochemical Company, has been appointed chief of the Alkalies Section, Inorganic and Agricultural Chemicals Branch of the Chemical Division, National Production Author-

th, as of August 1, 1952.

The appointment is temporary, and Mr. Bruyn will be in Washington until April 1st, 1953. He has been granted a leave of absence by the Hooker Company for the tenure of the appointment. During World War II he worked closely with the War Production Board on allocation of Hooker chemicals required for military purposes.

Keen Named to USDA Post

F. P. Keen, Bureau of Entomology and Plant Quarantine, has been appointed special asistant to the leader of the Bureau's Division of Forest Insect Investigations, the U. S. Department of Agriculture has announced.

Mr. Keen, with headquarters at Berkeley, Calif., will be responsible for coordinating more effectively the research, survey, and control activities of the Division in the eastern States. He has been in charge of forest insect investigations and technical supervision of insect control projects in California, and is being replaced by C. B. Eaton.

Am. Potash Buys Eston

American Potash & Chemical Corp. has acquired all of the outstanding capital stock of Eston Chemicals, Inc., of Los Angeles in exchange for 22,837 shares of American Potash Class B stock, Peter Colfax, president, has revealed.

Eston Chemicals, which will

become a division of American Potash, manufactures agricultural and industrial chemicals, refrigerants andaerosols, with sales in the first half of 1952 aggregating \$1,400,000. The company has facilities at Vernon and Turrance.

There will be no change in policies, products or personnel of the Eston division, but the broader facilities of American Potash and Chemical Corporation will enable Eston to expand and strengthen its activities.

Alfred M. Esberg, head of Eston since 1934, has been elected a vice-president of American Potash, and George S. Wheaton, vice-resident, has been named as assistant vice-president.

Union Bag Ups Nelson

Union Bag and Paper Corporation has announced the appointment of E. T. Nelson as assistant western district sales manager for multiwall bags. He will headquarter in the company's Chicago office.

Mr. Nelson joined the Union organization in 1928. Prior to his new appointment he had been a field representative for Multiwall bag sales in the Minneapolis area.

Attends French Meeting

H. L. Woudhuysen, of the New York firm H. L. Woudhuysen & Associates, planned to fly to Paris, France, on September 12th to attend the 3rd International Congress of Crop Protection, to be held from September 15th to 21st in the Sorbonne University. Mr. Woudhuysen was also present when the first meeting of this kind took place at Louvain, Belgium in 1946.

CFA Offices Move

Headquarters of the California Fertilizer Association have been moved to a new location, according to an announcement by Sidney H. Bierly, executive secretary of the group. The new location is suite One, Boothe Building, 475 Huntington Drive, San Marino 9, Calif. The former executive offices were on District Bullevard, Los Angeles. The new address was effective as of August 1, 1952.

Chase Names New S. M.



R. F. NORCOTT

R. N. Conners, vice-president and general sales manager, Chase Bag Co., has announced the promotion of R. F. Norcott to sales manager of the Chase sales organization in Milwaukee. Mr. Norcott, with Chase more than twell-ve years, was formerly a salesman in Buffalo and Chicago. Mr. Norcott assumed his new post August 1st.

New Production Method

The Stauffer Chemical Company has announced that it has obtained exclusive American rights to a new process for the production of ammoniated superphosphate. This process, developed and patented by Rumianca, Societa Per Azioni, Turin, Italy, produces a pelleted material containing nitrogen and phosphorous in amounts which can be varied over a wide range, depending on demand.

About the same proportion of phosphate is water soluble in the material made by this process as in single superphosphate. The material may be bagged or shipped without the aging period usual in superphosphate manufacture. This reduces the cost of the plant by eliminating the need for storage facilities for curing single super. The capacity of existing plants may be enlarged by the use of this process without the need for additional storage. The raw materials are the same as in the standard processess-phosphate rock, sulphuric acid and ammonia.

The process will be made available to other producers under sub-licensing agreements, Stauffer says.

Books and other Aids...

Handbook of Agricultural Pest Control

by S. F. Bailey and L. M. Smith

190 pages, 3.25 in U.S.A.

A practical handbook for the custom spray operator, the pest control operator, farm advisor, agricultural chemical salesman and field worker. This handbook covers the agricultural chemicals (insecticides, fungicides, herbicides, plant hormones and nutrient sprays, defoliants, etc.), their rates of application, useful formulas, as well as chapters on fumigation, spray machines, toxicology, dusts and dusting, aircraft, and mosquito control.

The book is pocket sized (5 x 7½ x ½ inches), bound with a flexible leatheroid cover, for convenient, practical use. It includes more than 100 tables for practical everyday use. Conversion formulas and examples illustrate application to specific uses.

Insect Control by Chemicals

by A. W. A. Brown

817 pages, price \$12.50

817 pages. This text traces the relation between molecular structure and toxicity; classifies the insecticides and gives their chemical, physical properties; discusses the hazards to avoid in formulation, mixing and use of compounds; and illustrates modern application equipment, including nozzles; sprayers, blowers, etc;

The Chemistry and Action of Insecticides

by H. E. Shepard

504 pages, price \$7.00

This new book gives a vast wealth of information on insecticides—their chemical, physical, and toxicological aspects. Helps the chemist determine their important applications and their effects upon insects. Gives history, commercial importance, major uses.

Covers these chemical groups: Arsenical Compounds; Fluorine Compounds, Sulphur Compounds; Copper Compounds; Inorganic Substances; Nicotine; Rotenone; Petrolcum, Soaps, Creosotes; Synthetic Organic Insecticides.

Destructive and Useful Insects Their Habits and Control

by C. L. Metcalf and W. P. Flint

1071 pages, price \$10.00

This authoritative guidebook covers hundreds of both useful and destructive insects—treating the inner and outer structure and form of general species—explaining their eating and breeding patterns, and life cycles—telling you how to recognize them in any stage of development.

Here are descriptions of more than 500 types of insect pests of the U. S. and Southern Canada. The insects are grouped in accordance with the crops, animals, or products they attack. For each insect you are given recognition marks and types of injury it does to man—enabling you to determine exactly what insect is damaging your crop, and supplying you with the most effective means of dealing with it.

Agricultural Chemicals 175 Fifth Ave., New York 10, N. Y. Enclosed is payment. Please send the books checked.	6 9 9 9 9 9
☐ Handbook of Agricultural Pest Control—\$3.25 in U. S.A., \$3.75 elsewhere ☐ Insect Control by Chemicals—\$12.50 ☐ The Chemistry and Action of Insecticides—\$7.00 ☐ Destructive and Useful Insects—\$10.00 (Add 3% sales tax in New York City) Name	Order direct from Agricultural Chemicals 175 Fifth Ave. New York 10, N. Y.
Street	

California Fertilizer Assn. Plans November Meeting



M EETING at the Desert Inn, Palm Springs, Calif., the Califorma Fertilizer Association plans to hold its annual convention November 10-12, according to Sidney H. Bierly, executive secretary and manager of the Association.

This year's meeting will mark the first time the group has met at the Desert Inn (pictured above) and according to information from the Association's new headquarters (at Suite 1, Boothe Building, 475 Huntington Drive, San Marino 9, Calif.) the November gathering will be the largest ever held by the CFA.

Detailed plans for the meeting had not been announced by the Association at press time, but officials declared that speakers representing a wide scope of agriculture would be on the agenda.

CFA president is S. B. Tatem, Swift & Co., Los Angeles; B. H. Jones, Sunland Industries, Inc., Fresno, is vice-president; William E. Snyder, treasurer, and Mr. Bierly, secretary.

The Association has invited all industry people and others interested in agriculture to attend. Hotel reservations are to be made by the individual and Mr. Bierly has advised making reservations as early as possible.

Picture-Story of CFA

The October issue of Agricultural Chemicals will carry a picture story of plants and people who make up the California Fertilizer Association. Watch for it in the next issue. —Ed.

Phillips Personnel Shifted

Five personnel changes in the Bartlesville office of Phillips Chemical Company, effective August 15, 1952, have been announced by K. S. Adams, chairman, and Paul Endacott, president, of Phillips Petroleum Company.

Edward Dolezal, chief mechanical engineer, will be assigned to the newly-created post of chief project coordinator. Austin Morgan, assistant chief mechanical engineer, will succeed Mr. Dolezal as chief mechanical engineer. T. M. Hipp, presently on assignment on an ammonia plant construction near Houston, Texas, will be transferred and assigned to the newly-created post of chief construction engineer.


M. W. Bennett, office manager, will be assigned to the newlycreated position of chief economics engineer, and M. W. Monk, assistant office manager, will succeed Mr. Bennett as office manager.

Cotton Conference to Cal.

Progress in cotton irrigation and harvesting will be highlighted at the 1952 Beltwide Cotton Mechanization Conference to be held October 22-24, at Bakersfield-Shafter, California.

Field demonstrations will be presented at the Cotton Experiment Station at Shafter to supplement the mechanization discussions at Bakersfield.

Sponsor of the event is the National Cotton Council, Memphis, Tennessee.



We're one team now to serve you better

You have probably heard of the recent purchase of Julius Hyman & Company by the Shell Chemical Corporation. Here's additional news about our plans for more efficient service to you:

Julius Hyman & Company Division of Shell Chemical Corporation will market the powerful insecticides, aldrin and dieldrin, as well as the unique soil fumigants, D-D and CBP-55. It will also market on the West Coast Shell Chemical's long-established line of spray oils.

Headquarters for Julius Hyman & Company Division will be in Denver, Colorado, with area sales offices associated with the Shell Chemical offices in New York, Atlanta, Chicago, Houston, St. Louis, San Francisco, Los Angeles and Denver.

The Julius Hyman representative in your area will keep you informed on developments pertaining to our growing line of agricultural chemicals. He can be a valuable ally in today's dynamic agricultural marketing picture and he stands ready to serve you.

Julius Hyman & Company Division SHELL CHEMICAL CORPORATION

Chemical Partner of Industry and Agriculture P. O. Box 2171, Denver 1, Colorado



Hall Forms New Company

J. Newton Hall, formerly vicepresident in charge of sales for Julius Hyman & Co., has just organized a new firm which, operating under the name of Pioneer Chemical Associates, will act as consultants, brokers and marketers, specializing in insecticides and other agricultural chemicals and raw materials therefor. Offices are being opened at 3035 East Colfax Ave., Denver 6, Colorado, telephone FRemont 8839.

As a part of its brokerage and marketing services Pioneer will use its own labels and brand name on some products. This arrangement will allow for maximum flexibility in selecting shipping points, intermingling of supplies from different sources and consolidating regional requirements. Mr. Hall indicates that such flexibility should result in economies and improved service to buyers. Quality standards will be checked and controlled by Pioneer's own technical personnel and by independent laboratories.

Details as to the company's other personnel and branch offices will be announced shortly.

Bagpak Opens K. C. Office

International Paper Co., has opened a new Bagpak Division sales office in Kansas City, Kansas, R. I. LaMarche, division sales manager, announced, W. O. McDuffie has been named to head the new office.

Conn. Field Day Held

The Connecticut Agricultural Experiment Station, New Haven, held its annual field day on August 20, with some 800 persons in attendance. Dr. William L. Slate, director emeritus of the station, as principal speaker, praised the Connecticut farmers for supporting and utilizing agricultural science.

It was the growers themselves, who in 1875, were the prime movers behind the effort to establish the Connecticut Station, first in America-Since that time, he said, Connecticut farmers have been generous with their time, thought and money in its support. In the earlier days, when the Station lacked lands on which to con-

duct its experiments, the farmers were quick to offer their farms for its purposes, often supplying labor and tools as well.

This pattern has continued, the retired Experiment Station Director reported, and today many of the Station's field trials are conducted on the farms of commercial growers.

Mr. Slate was introduced by Dr. James G. Horsfall, present director of the Station, who also introduced to the field day audience, Dr. Paul J. Anderson and members of his staff. Dr. Anderson has, for 27 years, headed the station's tobacco research at its Windsor Tobacco Laboratory. Emphasis at the 1952 field day was on tobacco, with many special exhibits and demonstrations on this crop arranged.

In addition, field day visitors inspected experimental plots where research on plant breeding, disease and insect control, soils and forestry, is in progress.





NOW!

a lower cost
INSECTICIDE
EXTENDER
that's

FOR HIGHEST

at lowest cost always specify uniform Palmetto Clay Natural deposit purity combine with modern production and quality control to bring you the finest low cost extender.

COMPARE!

Compare quality controlled Palmetto Clay with your present diluent. Send for a working sample today.

3 Ways Better!

Finer:

80-85% of Palmetto particles are less than 2 microns in size. Allows most effective active ingredient dispersion.

Less Abrasive:

Less than 0.50% retained on a 325 mesh screen. Virtually eliminates abrasion.

Lasting Suspension:

Make this test, mix 25 grams of Palmetto diluent in 225 c.c. of distilled water by stirring 7 minutes in high speed mixer.

Then immediately fill a 100 c.c. Nessler tube and note height of solids column after 24 hours.

INTERNATIONAL CLAY CORPORATION

P. O. BOX 371

GRANITEVILLE, S. C.

AGRICULTURAL CHEMICALS

Joins Ethyl Corp. Sales



ALBERT E. GRIFFITHS

Dr. Albert E. Griffiths formerly of Socony-Vacuum Oil Co., has joined the chemical cales division of Ethyl Corporation, to supervise technical service work connected with agricultural chemicals, including benzene hexachloride and lindane, according to an announcement by Malcolm P. Murdock, general sales manager.

A native of Center Berlin, N. Y., Dr. Griffiths attended Cornell University, receiving his B.S., M.S., and Ph.D. degrees in 1933, 1937, and 1939, respective-He also studied microchemistry and plant physiology under a fellowship at the Boyce Thompson Institute in Yonkers, N. Y

Dr. Griffiths was associated with G. L. F., Inc., a farm cooperative of New York, from 1933 until 1935. Returning to Cornell, he served as instructor in vegetable crops from 1935 to 1940, while completing his graduate work. From 1940 to 1945, he was professor of horticulture at the University of Arizona, and was also superintendent of the Salt River Valley Vegetable Research Farm, located at Tempe, Ariz., during that period. Prior to joining Ethyl, he was group leader in the Agricultural Chemicals Section of the Technical Service Department of Socony-Vacuum, with which he was associated from 1945 to 1952.

Hold Atomic Energy Meet.

The fourth annual summer symposium on The Role of Atomic Energy in Agricultural Research was to be held August 25-30 at Oak Ridge, Tenn., under the sponsorship of the University of Tennessee. Prominent speakers were scheduled to present papers, including Dr. Nathan S. Hall, professor of agronomy at North Carolina State College: and Dr. Sterling R. Olsen, who is with the regional phosphate laboratory operated at Fort Collins by

Colorado A & M College and the USDA.

Plan \$5 Million Cal. Plant

Filtrol Corp., Los Angeles, has just announced plans for construction, in the immediate future, of a new \$5 million plant at Vernon, Calif. for the manufacture of ammonium sulfate fertilizer and high purity alumina. The plant will further process the alumina to produce alumina catalyst, stabilized alumina ca-

talyst and impregnated alumina catalyst. Marketing plans for these products include service to the chemical industry for various catalytic processes such as oxidation, dehydrogenation and ammoniation. Shell Chemical Corp. will market the ammonium sulfate fertilizer output of the plant. Construction is to begin during the fourth quarter of this year, with completion scheduled for a year later. Wright W. Gary is president of the Filtrol Corp.

FOR BEST RESULTS. depend on Eston agricultural chemicals FOR YOUR 1953 PROGRAM, USE...

ALKRON®

Parathian formulations

BROMOFUME® Ethylene dibromide soil fumigants

ESTONATE DDT dust concentrates and emulsifiable solutions

ESTOMITE Residual type miticide

ESTONOX Toxaphene formulations

METHYL BROMIDE Space fumigant

TETRON® Tetraethyl pyrophosphate formulations

TUMBLEAF Cotton defoliant

You'll find Eston Argicultural Chemicals manufactured to give uniformity, reliability and maximum effectiveness . . . products that will give you the best results...priced to give you every profit advantage. Also... when you buy Eston agricultural products you get the benefit of Eston technical service. Eston technicians can help you with all your insecticide problems.

Write or wire for further information.

FOR THE BEST IN AGRICULTURAL CHEMICALS...IT'S ESTON!



AMERICAN POTASH & CHEMICAL CORPORATION 3100 East 26th Street, Los Angeles, California





NOW! Control ALL Major Fungous Diseases of Tomatoes and Potatoes with NEW Du Pont Manzate

TOMATOES: This single new fungicide now does the job where several different or alternate schedules have been needed to control fungous diseases. It provides such excellent control of early and late blights, gray leaf spot and Septoria that you're sure of higher yields and better-quality tomatoes through a longer picking season.

"Manzate" is available now in a wettable powder formulation. Research indicates that "Manzate" is effective against a wide variety of fungous diseases in addition to those affecting tomatoes and potatoes. For details, write Du Pont, Grasselli Chemicals Dept., Wilmington, Del.

Be Port Chemicals for the Form Include: Fungicides: MANZATE†, PARZATE* (Liquid and Dry), FERMATE, *ZERLATE, *Copper-A (Fixed Copper), *SULFORON* and SULFORON* X Wettable Sulfurs . . Insecticides: DEENATE* DDT, MARLATE* Methoxychlor, LEXONE* Benzene Hexachloride, KRENITE* Dinitro Spray, EPN 300 Insecticide . . Weed and Brush Killers: CMU, AMMATE, *2,4-D, TCA, 2,4,5-T . . . Also: Du Pont Cotton Dusts, Du Pont Spreader-Sticker, PARMONE* Fruit Drop Inhibitor, and many others.

POTATOES: "Manzate" also does the job where several different or alternate schedules have been needed to control potato blights. This new chemical gives excellent results when both early and late blight attack at the same time. It provides a complete program for all season long. Several years' tests of "Manzate" have shown greatly increased yields of No. 1 tubers.

On all chemicals always follow directions for application. Where warning or caution statements on use of the product are given, read them carefully.



BETTER THINGS FOR BETTER LIVING . . . THROUGH CHEMISTRY

1 Trademark of E. I. du Pant de Nemours & Co. (Inc.)

Opens New Chemical Co.



W. LEROY TRAYLOR

W. LeRoy Traylor, who has been associated with the Stauffer Chemical Company for the past fourteen years, — five years of which was spent in the capacity of vice-president and general manager of the Stauffer Chemical Company of Florida, Inc., and the past two years as sales manager of the parent company's South Eastern Division — announces the opening of his own business, the Traylor Chemical and Supply Company of Apopka, Florida. This firm will carry a complete line of agricultural and industrial chemicals — serving customers in Florida, Georgia, Alabama and Latin American countries.

Shell Appoints Boultbee

A. H. Boultbee has been appointed director of manufacturing research for Shell Oil Company, according to F. S. Clulow, vice-president. He will replace D. L. Yabroff, who is returning to Shell Development's research laboratory at Emeryville, Cal., as associate director in charge of chemical products.

Dr. Boultbee was born in Kenora, Ont., Canada, obtained his Ph.D. in chemical engineering from the University of Toronto and then studied at the Imperial College of Science and Technology in London, where he was granted a D.I.C. in chemical technology.

He joined Shell Oil Company as a chemist at Martinez Refinery in 1935.

Marathon Ups Brown

G. Blaine Brown has been appointed advertising manager of the Chemical Division, Marathon Corporation, Rothschild, Wisconsin, the company has announced. Mr. Brown joined the chemical division in 1942 and shortly thereafter entered the Air Force. He was later commissioned and served in the European theatre. On his return from active service he obtained his B.S. degree in chemistry at the University of Michigan and, in 1946, rejoined Marathon in the

chemical research staff. He was soon advanced to the position of chief control chemist. Because of his particular abilities he was later transferred to the sales department in the capacity of sales service supervisor.

The company has also announced the appointment of Frank G. Morman in charge of sales in New England and the Northeastern states.

two PICCO HI-SOLV solvents



INSECTICIDES

AGRICULTURAL SPRAYS

*Increase the Effectiveness

* Decrease the Cost

OF YOUR PRODUCTS

You can improve your sprays and at the same time cut costs, by using high-solvency, aromatic PICCO Hi-Solv Solvents. The analyses given below reveal characteristics that make these two Picco Hi-Solvs ideal for your use. Write for complete data and samples.

Typical Analysis	Hi-Solv 30	Hi-Solv 473
Distillation Range, °F	266-374	400-520
Specific Gravity	0.8300.840	0.900-0.915
Color	Water White	Light Straw
Flash Point	80° F—TCC	180° F—COC



PENNSYLVANIA

INDUSTRIAL CHEMICAL CORPORATION

Clairton, Pennsylvania

Plants at Clairfon, Pa.; West Elizabeth, Pa. and Chester, Pa.

It's a Good Mixer!



Nitrogen is in great demand. Even Phillips tremendous capacity isn't equal to today's requirements. But we're making four different kinds of high-quality nitrogen material for mixers and farmers.

- AMMONIUM NITRATE . . . Phillips 66 Ammonium Nitrate contains 33% N. Small, coated, uniform pellets flow freely and resist caking.
- NITROGEN SOLUTIONS . . . there are three Phillips 66 Nitrogen Solutions for use in the preparation of high-analysis fertilizers and the ammoniation of super-phosphate. These solutions keep handling costs low promote rapid, thorough curing.
- 3. ANHYDROUS AMMONIA . . . Phillips 66 Agricultural Ammonia contains 82% N. Convenient, economical source of nitrogen for fertilizers.
- 4. AMMONIUM SULFATE (see photograph and description above.)

For full information write our nearest district office.



PHILLIPS CHEMICAL COMPANY

A Subsidiary of Phillips Petroleum Company

FERTILIZER SALES DIVISION . BARTLESVILLE, OKLAHOMA

DISTRICT SALES OFFICES:

NORFOLK—610 Royster Bidg. • TAMPA—7 Terrace Office Bidg., 404 Marion St. • HOUSTON—604 City National Bank Bidg.

OMAHA—WOW Bidg. • AMARILLO—First National Bank Bidg. • PASADENA—16 North Marengo Ave. • BARTLESVILLE—Adams Bidg.

Walton Named to Post



CHARLES W. WALTON

Appointment of Charles W. Walton as senior research analyst has been announced by Chemical Construction Corporation. He will specialize in the agricultural chemicals field. Mr. Walton, a retired Colonel of the United States Army, was recently the representative in Italy for Griffith Laboratories, Inc., where he established a sales agency for the firm. He was also a consultant and representative of several other firms.

During his military service, Mr. Walton was chief of operations and Administration of the United States Air Force Service Command, Mediterranean Theatre of Operations, fom 1943 to 1945; Vice President, Economics, Allied Commission Italy in 1945 and 1946; and Comptroller, Air Defense Command, with headquarters at Mitchell Field from 1946 to 1947. He received a B. S. degree in agriculture from the University of Wisconsin and an M. S. degree from the Harvard Graduate School of Business Administration.

Information on Borer

"Battle of the Borer" is the title of an article in the August issue of "Monsanto Magazine", published by Monsanto Chemical Co., St. Louis 4. The illustrated article reviews the life cycle and breeding habits of the borer, and tells of the outlook for holding the pest in check through scientific farming methods and insecticides.

Sprayer, Duster Assn. Elects

R. B. Chapin, president of the R. E. Chapin Mfg. Works, Inc., Batavia, N. Y., was reelected president and chairman of the executive board of the National Sprayer & Duster Assn. at its recent annual meeting. H. F. Brandt, president of the Dobbins Mfg. Company, Elkhart,

Indiana, was reelected vice-president and vice-chairman of the executive board. D. P. Lewis, secretary of the H. D. Hudson Mfg. Company, Chicago, was reelected association treasurer. Frank J. Zink and Earl D. Anderson of Frank J. Zink Associates, Chicago, were reelected respectively counsel and secretary of the association.

Reelected to the executive board in addition to the officers named were: P. L. Hauser, sales manager, Lowell Mfg. Company, Chicago; C. D. Leiter, vice president in charge of sales, F. E. Myers & Bro. Co., Ashland, Ohio; R. W. Merritt, vice-president, Root Mfg. Comapny, Malta, Ohio; and T. M. Burton, vice-president, D. B. Smith & Company, Utica, N. Y.

James R. Hile, manager of the insecticide division, Acme Quality Paints, Detroit, was guest speaker at the annual dinner.

The executive offices of the association are at 4300 Board of Trade Bldg., Chicago 4, Illinois.

COPPER SULPHATE

Crystals - Superfine - Powdered BASIC COPPER SULPHATE

ZEE-N-O

(Neutral Zinc)

The High Test Nutritional Basic Zinc
56% Zinc as Metallic

MANGANO

(Neutral Manganese)

The High Test Nutritional Manganese
55% Mn as Metallic

W. R. E. ANDREWS SALES, INC.

1505 Race Street, Philadelphia 2, Pa. Since 1926

Agricultural Chemicals Specialists



That's right, you can really be out . . . out in the cold if your product is sifting out of its container, especially when Betner has available a variety of bags that positively prevent sifting-out of contents . . . keep contents in the bag.

For instance, the Betner "Duo-Tite" Bag is constructed to hold bulk powdered material, particularly chemicals, with positively "no-sift" of contents. Available in a variety of sizes to hold up to 25 pounds, it's folded, glued and heat-sealed to guarantee "sift-proofness." The Betner "Duo-Tite" bag can be pre-printed up to four colors and can also be made with a combination of materials.

COMPLETE SERVICE!

Betner can supply the special machinery for closing the "Duo-Tite" bog . . . it heat-seals, double folds and pastes the taps in exactly the same manner as the bottom is constructed.



These bags conform to I.C.C. 2-D specifications









• Whatever the packaging need, there's a Betner bag . . . FILL IT!

Benj C Betner Co Devon, Pa.

Plants also located in: Richmond, Virginia; Paris, Texas;
Appleton, Wisconsin; Los Angeles, California

Use of Chemicals Called Essential for Development of Grasslands in U. S.

An estimated 15,000 persons turned out on August 20 for the Grassland Field Day, part of the Sixth International Grassland Congress held at Pennsylvania State College, August 17 to 23. In the neighborhood of 1200 scientists from 49 different countries with 40 official representations already were deep in the problems of grassland improvement.

The visiting scientists had an opportunity to experience an important American way of transferring agricultural information from trial plots and test tubes to practical farming fields. The Field Day proved of especial interest to foreign delegations. On the other hand, the visiting farmers inspected the largest display of implements and agricultural chemicals ever assembled for a specific use, in this case for growing forage crops.

Products from all of the major manufacturers of equipment and supplies in this field were on display in addition to booths prepared by various bureaus in the U.S.D.A. and state agricultural experiment stations. Exhibitors totaled between 60 and 70.

Philip V. Cardon, former administrator for the Agricultural Research Administration and chairman of the official United States delegation, was raised from temporary to permanent president of the Cangress at the opening plenary session-Vice-presidents elected were: C. A. Volio, Minister of Agriculture, Costa Rica; and J. Griffiths Davies, Div. of Plant Industry, C. S. and I. R. O., Canberra, Australia.

During the week, more than 200 papers were read in English, French or Spanish, the official languages of the Congress, in a dozen different sections. Broad topics ranged from the genetics and breeding of forage crops through soil management and the harvesting and preservation of forage. A number of joint sessions, workshops and tours of the campus and college farms were arranged.

Ag. Chemicals Discussed

SINCE world-wide problems were under study, many new and varied uses of agricultural chemicals were discussed. Fertilizers had an important place in the program but supplementary materials also were given consideration. Experiments on

the range lands of peninsular Florida, conducted by J. R. Neller of the Station at Gainesville, show that sulfur must be applied at least once a year for good growth of leguminous forage. The amount of sulfur contained in mixed fertilizer is decreasing, he pointed out, and sulfur-deficient soils in the United States are wide-spread. Leguminous forage requires about 5 pounds of sulfur per ton per acre for good growth, he said.

Deficiencies of copper and zinc

Greetings....

To Members of

Natl. Agricultural Chemicals Assn.

WE are pleased to announce the opening of our new plant at Campo, California, to help meet the increasing demand for "PHYLLITE*" pyrophyllite.

With these new facilities, our production capacity is now 300 tons per day . . . and we still continue to offer the same uniform product to our customers.

As additional information our Campo mill will be the largest producer of feldspar and silica on the West Coast and as the feldspar has a potash content of 16% it is being used extensively in a Stone Meal form for agricultural purposes.

*PHYLLITE (trade name) is absolutely non-abrasive and adheres readily to foliage and all surfaces.

Packed in 50 pound valve bags, 20 ton lots Lowest prices on West Coast, f.o.b. plant Smaller quantities if desired

Write for information and samples, to

PIONEER PYROPHYLLITE PRODUCERS

HAncock 2-2992

P. O. Box 686

Chula Vista, Calif.

It's incredible! ... the commodities that go into a

bag

Polystyrene—Feeds—Calcium Chloride—Salt—Ammonium Nitrate—Cement—Rosin—Fertilizers—Sugar—Flour -Insacticides-Fine Chemicals-Clays-Rock Wool-and over 400 other products costing from less than a cent a pound to over a dollar a pound are being packed in Bagpak Multiwall Paper Bags.



CHRESTA

EB ASSESSES

in the soil which are transmitted to forage crops cause a wasting malady among sheep in South Australia, D. S. Riceman of the University of Adelaide reported. Correction of these deficiencies is brought about by a single dressing of the land with Zn-SO₄. 7H₂O and CuSO₄. 5H₂O₄ each applied at 7 pounds per acre.

With the phenoxyacetic acid derivatives, M.C.P.A. and 2,4-D, it is possible to kill a wide range of pasture weeds, according to J. Strykers, State Agricultural College, Ghent, Belgium. The most favorable time for controlling the widest range of perennial weeds, he has learned, is at the end of the grazing season in September or October. He has obtained good results from the use of small amounts of active substance applied at different times, either as sprays or mixed with sand. Ammonium nitrate creates a problem in mixed fertilizer used as a carrier for these growth regulators, he has found. Ammonia inactivates phenoxyacetic acid derivatives.

These herbicides have two important uses in Britain, W. G. Templeton, Jealotts Hill Research Station, Bracknell, Berks, England, declared. They are: the removal of weeds from cereal crops undersown with a grass-clover mixture in the establishment of short leys, and in the eradication of perennial weeds in permanent pastures that can not be ploughed. While both fall and spring applications have given good control of weeds in his experiments, the autumn applications on the average have shown increased production of useful herbage the year following applica-

Use of weed-free crop seed, prevention of weed seed maturation, adequate fertilization, use of adapted crops, and intelligent management practices were advised by K. P. Buckholtz of the Wisconsin Station. Herbicides show promise of weed control, he said, that can not be easily affected by other methods.

A very small percentage of the compounds tested as herbicides carry through to a satisfactory use stage, W. C. Dutton, Dow Chemical Co., Midland, Mich. stated. It seems reasonable, he commented, that industry should be expected to assume major responsibility in the earlier stages, but industry and public research agencies should cooperate in the advanced stages of the development of a product. Cooperation so far has been excellent, he agreed.

The principal insecticides used on forage crops are aldrin, chlordane, toxaphene and DDT, and in general the dosages required to secure satisfactory results, according to F. C. Bishopp, Bureau of Entomology and Plant Quarantine, U. S. D. A. are not sufficient to adversely affect live-stock. However, the presence of any of these materials, particularly in excessive quantities, may result in their excretion in milk or storage in fat. The relationships of the insecticide, formulation, climatic conditions, date and method of application, kind of crop, curing and storage to the extent of residue, all are important, he said.



BENZENE HEXACHLORIDE

13-15% GAMMA-TECHNICAL-FINELY FLAKED

A prime advantage of insecticides formulated with BHC is their QUICK killing action.

"ETHYL" technical BHC is manufactured in the heart of the cotton belt at Baton Rouge, Louisiana.

ETHYL CORPORATION

100 PARK AVENUE

NEW YORK 17, N. Y.



During the past fifty-five years we have watched many ambitious and enterprising manufacturers endeavor to copy the very extensive Oberdorfer line of bronze rotary gear pumps or present the world with their substitute.

It was not our business to notify these gentlemen that it would be financial suicide to try to duplicate the combination of quality and price so long associated with the name of Oberdorfer thruout the world. History is amply littered with the evidence.

There are available today many spray pumps more intricately designed, more delicately constructed and more costly to purchase and operate in the field.

Why gild the lily! Demand an inexpensive, trouble free, corrosive resistant bronze gear pump that accurately meters your spray formulations, positively, efficiently and at the lowest cost per acre.

Buyberdorfer

Agricultural Pump Div., Oberdorfer Foundries, Inc. 1129 Thompson Rd., Syracuse, N. Y.

BRONZE OBERDORFER
SPRAYING PUMPS

HOW TO combat weeds ...

check their introduction,
spread regrowth . . .
effectively-economically



Here is an authoritative, thorough book that gives you all the data and practical help you need to prepare—and carry out—a tested, efficient, successful method of attack on any weed in any location quickly and effectively. Based on experience, research and experiment, it shows what methods of weed control have been proven most effective for weed of all species—from erab grass to wild muschard—points out what methods can be applied conomically in certain areas—shows how and when to apply a control measure, the erason and rate of application, dusage, etc., and outlines the materials and machinery needed.

Quickly shows you haw to credicate SPECIAL WEEDS Bracken Farm Canada Thistle Horse Nettle Net Grass I Poison Iry, Poison Duk Prickly Pear Queck Grass

Sumac. Poison Out Prickly Pear Quack Grass Russian Thistle Wild Garlic, Wild Onion IN SPECIAL AREAS

pasture range lands lawns golf greens and fairways suchards vineyards along lakes and streams

WEED CONTROL

A Textbook and Manual By WILFRED W. ROBBINS ALDEN C. CRAFTS and RICHARD N. RAYNOR All at the College of Agriculture University of California

543 pages, 6 x 9, 202 illustrations, charts, tables, \$6.00

The book provides the farmer, horticulturist, greenskeeper, highway maintenance engineer, landscape garduner, etc., with a definite ainwer to his particular problem, indicating the methods that are especially adapted to mendow, grainfield, orchard, gold green, rondside, greenhouse, and other specific licetions.

Add 3% Sales Taz in N. Y. City

AGRICULTURAL CHEMICALS

175 Fifth Ave. New York 10, N. Y.

A Compact And Comprehensive Handbook

CHEMISTRY AND USES OF INSECTICIDES

By E. R. de Ong, Consulting Entomologist, Agricultural Technologist.

Written by an outstanding student of entomology and agricultural technology, this book covers all the major insecticidal agents in detail, describing not only their chemical nature and properties, but also their specific action on various types of insects, their methods of application, and their effect on animals and humans.

A valuable source of practical information for agriculturalists, this volume will be essential to all large commercial growers of fruits and vegetables. It will also prove most helpful to progressive farmers, to chemical manufacturers, research chemists, exterminators, students and teachers looking for a modern text for college use.

1948, 445 pages, Illustrated, \$6.75.

Add 3% Sales Tax in New York City

"AGRICULTURAL CHEMICALS"
175 Fifth Avenue New York 10, N. Y.

Use of strong acids, such as sulfuric and hydrochloric, in the preservation of silage is common in Finland, Artturi I. Virtanen, Biochemical Institute, Melsinki, reported. The acidity of forage must be brought to approximately pH4 for proper preservation, he has learned, which is not practicable with protein-rich fodders. Decomposition processes in fodder artificially acidified are slight, he said.

In the field of fungicides for use on special-purpose turf, mercury preparations, long the standard treatment for dollar spot and brown spot, have found worthy rivals in cadmium and thuriam compounds, F. V. Grau, U. S. Golf Association, Greens Section, Beltsville, Md. declared. Development of 2,4-D has advanced knowledge of selective weed control resulting in better turf, he said. Sodium arsenite is gaining in popularity among professional greenskeepers while potassium cyanate is making friends among laymen, he commented.

W. Y. Conference Date Set

Dates for the annual New York State Insecticide-Fungicide Conference have been set for November 11, 12, and 13, according to officials of the group. Program details have not been announced as yet, but the meeting will be held, as usual, at Cornell University, it was indicated.

Speakers Bureau for CFA

The California Fertilizer Association, Los Angeles, Calif., has announced the formation of a speaker bureau to meet the demand for qualified speakers on the use of commercial fertilizers before farm organizations, nursery and floriculture groups, home gardener organizations, service and women's clubs, etc.

Okay Chemical Corp. Terms

Chemical Corp. of Colorado, Denver, which has been threatened with bankruptcy, has worked out an arrangement with its creditors under which the company will continue to operate. Some \$285,000 in unsecured debts will be paid off on the basis of \$33\forall 6 on the dollar, \$11\sqrt{6}\$ of the total claims to be paid immediately in cash and the balance in notes payable

within two years. Claims of four secured creditors did not figure in the hearings. The major creditor was a bank whose claim for \$219,459 due on loans was secured by accounts receivable of \$316,463 and a merchandise inventory of \$55,098.

The company's difficulties resulted in part from some \$92,000 in damage suit judgments obtained by Texas cotton farmers who charged that their crops were damaged by cotton insecticides sold by Chemical Corp. of Colorado. A fire in the company's warehouse last August, causing \$70,000 damages, was said to be another contributing factor.

Phytopaths Meet in Ithaca

Scores of technical papers and the annual fungicide colloquium were on the agenda of the 44th meeting of the American Phytopathological Society, Ithaca, N.Y., September 7-10.


The colloquium, featuring talks by industry leaders and offering manufacturers the opportunity of introducing new materials to the trade,





MINERAL SALTS

One of the country's foremost producers of Agricultural Chemicals and Soluble Mineral Salts.



For MANUFACTURERS, MIXERS And CHEMISTS

We can supply you with most any combination of mineral mixtures mixed to your own specifications. Soluble Manganese, Copper, Zinc, Iron, Boron and Magnesium.

Producers of **ES-MIN-EL** — Essential Mineral Elements.

For further information 'phone, wire ar write

TENNESSEE



CORPORATION

Lockland, Oh

was to be held on Tuesday evening, September 9th, Dr. L. Gordon Utter, Phelps Dodge Refining Corp., New York, was to be in charge of this portion of the program.

Dr. George L. McNew, managing director of Boyce Thompson Institute for Plants Research, APS president, was scheduled to preside at the opening sessions.

NFA "Meter" Offered

A new "Plant Food Meter" has been made available to the fertilizer trade by the National Fertilizer Association, Washington, D. C. The meter consists of a plastic "sleeve" with open ends so that the cardboard inside may be moved up or down. Printed on the card is information regarding the removal of plant food from soil by many different crops, at a certain number of pounds, tons, or bushels per acre.

The meter has space for imprinting a dealer's name, the Association says. This offers the opportunity of an advertising piece that will keep customers aware of the need for replacing plant food as it disappears from the soil with crop removal.

L. A. Kolker Retires

Leon A. Kolker, president and founder of Kolker Chemical Works, Inc., Harrison, N. J., retired August 31. The activities of the Kolker company, which is a division of Diamond Alkali Co., will hereafter be carried on by J. G. Brunton, in charge of sales, and Charles H. Kolker, in charge of operations and engineering, under the supervision of A. L. Geisinger, Diamond Alkali vice-president.

La Forge Rejoins Penick

Harold Noble of S. B. Penick & Co., New York, has announced that Byron La Forge has rejoined the sales staff of the company after an absence of nine months. During this interlude he had been with Chemical Insecticide Corp., Brooklyn. Prior to that time he had been with Penick's insecticide division for thirteen years. He will now be assigned principally to sales work in connection with Penick's essential oil department, with headquarters in New York.

Morrison Joins Summers

Kenneth D. Morrison, recently resigned as president of Naco Fer-



KENNETH D. MORRISON

tilizer Company, subsidary of W. R. Grace & Co., has joined the Summers Fertilizer Company, Baltimore, Md., in the capacity of vice-president in charge of sales.

Mr. Morrison has spent his entire life in the plant food industry. For many years, he was in charge of eastern sales distribution of both foreign and demestic potash producers.

Prior to his association with Naco Fertilizer Company, he was vice-president of The Davison Chemical Corporation.

Oct. Southwide Conference

The Southwide Chemical Conference to be held at Auburn, Alabama, October 23-25, will devote a major portion of its time to fertilizers.

A symposium, "Advances in Fertilizer Technology" is scheduled for Oct. 24. The advance program indicated that the following would appear on the program:

Dr. John R. Taylor, Jr., American Plant Food Council, Washington, D. C.; George V. Taylor, Spencer Chemical Co., Kansas City, Mo.; Dr. Vincent Sauchelli, Davison Chemical Corp., Baltimore; Dr. J. Fielding Reed, American Potash Institute, Atlanta; Dr. R. P. Thomas, International Minerals & Chemical Corp.; Dr. E. C. Kapusta, National Fertilizer Assn.; and Dr. A. M. Smith, Mathieson Chemical Corp., Baltimore.

Chairman of the symposium committee is J. I. Wear, American Potash Institute; and James A. Naftel, Pacific Coast Borax Co., co-chairman.

Sulfur Order Relaxed

With the visible supply of sulfur comparatively plentiful, the National Production authority has dropped distribution controls on sulfur and has increased the ceiling on quantities that may be kept on hand by users. Consumers may now keep a 60-day supply on hand, instead of

the former limit of 25 days.

The relaxing order came after producers' inventories had climbed by 93,000 long tons since the first of the year. Nearly 100 new projects in the U. S. and other countries are expected to add 4,000,000 long tons of sulfur yearly by the end of 1955.

The International Materials Conference estimates that production in the second half of 1952 will be at an annual rate of 6,400,000 long tons, compared with 5,900,000 in 1950 and 6,140,000 in 1951. Despite

For Dependable BLIGHT CONTROL



for your BORDEAUX SPRAYS and COPPER-DUST Mixtures

America's growers have been using Triangle Brand Copper Sulphate for over 60 years because this economical blight control method has been proven effective. Ask your dealer for Triangle Brand Copper Sulphate for Bordeaux Sprays... available in Large Crystals, Small Crystals, Superfine, Snow or Instant (free-flowing powder). Also Triangle Brand Basic Copper Sulphate for sprays and Copper-Dust mixtures.

FREE LITERATURE AVAILABLE ON REQUEST

Write today for these helpful, informative bookiets... "Bordeaux Mixture—Its Efficient Preparation and Use"..."Berdeaux Controls Late Blight on Tematoes"... and "Basic Conper Susinhate."

PHELPS DODGE REFINING CORPORATION

40 Wall St., New York 5, N. Y. * 230 N. Michigan Ave., Chicago 1, III.

the increased production, it was pointed out that this figure is still short of requirements.

The fertilizer industry was noting with pleasure that relaxation of regulations will permit users to accept shipments of sulfur by barge or vessel, as before.

SHADE TREE

(Continued from Page 79)

surface of the leaf brought better control than sprays on the upper surface. The insecticides were readily translocated from one side of the leaf to the other, Dr. Cantelo said.

Before a full auditorium on Wednesday, Dr. Frank E. Egler, consulting vegetationist, Aton Forest, Norfolk, Conn., discussed blanket spraying vs selective spraying for control of roadside brush. In his talk, "Roadside Brush Control — An Application of Plant-Community Management," Dr. Egler declared that "blanket" spraying is obsolete.

He pointed out that several areas of ground are at the side of every highway — the shoulder with no vegetation; the grassy area; and the unmowed grassy area. Before going into large-scale spraying operations, the highway department should be consulted, he said. The unmowed strips are the basic concern, since they are actually abandoned lands. Their growth gots through several stages — from plain soil, to weeds, to grasses, to shrubs to trees, he said. As each new group comes in, it crowds the old out.

Blanket sprays kill all vegetation, he declared, moving the process back to the grass stage. Selective spraying, on the other hand, is a totally different tool, enabling one



NITROGEN is the plant food element that makes grass grow! That's why it pays to make sure that pasture improvement plans include plenty of nitrogen.

When abundant nitrogen is used in a balanced fertilizer program, pastures produce a vigorous growth of good green grazing which supplies low-cost, nutritious, high-protein forage that can be harvested by livestock.

Well-fertilized pastures increase livestock profits with bigger yields of better quality grazing and more grazing days. Dairy cows produce more milk. Beef animals fatten quickly. Less barn feeding is required. Labor and feed costs are greatly reduced.

Here is a recommended pasture fertilization program, whether you are seeding new pastures—or improving established pastures:

- In the late summer or early fall, use a heavy application of high-nitrogen complete fertilizer.
- Follow this with nitrogen top-dressing in the late fall and again very early in the spring. Top-dress with 100 to 200 pounds per acre of AR-CADIAN*, the American Nitrate of Soda, or A-N-L* Nitrogen Fertilizer.

Watch this program make grazing crops get up and grow! Remember, it pays to use plenty of nitrogen on pastures. Extra nitrogen means extra grazing!

*Box II & Pat Off

Nitrogen Division

New York S. N.Y. . Richmond 16, Yo. . Supervell, Yo. . Columbia 1, S.C. . Atlanta 3, Go. . South Point, Ship . San Francisco 3, Go

to bring back vegetation to whatever stage is desired.

Dutch Elm Disease

PLEADING for a "positive approach" to protect healthy trees rather than removing infected ones, A. W. Hurford, secretary, Connecticut Forest & Park Assn., New Haven, told the group that the Dutch Elm disease can be controlled by coordinated control. A combination of DDT spraying, tree sanitation work and chemotherapy will go far toward halting this threat to the nation's elm trees, he said.

Pre-foliar DDT spraying in early spring is a "must" to save many elms, he said. Research is under way at the Connecticut Agricultural Experiment Station, he said, and is expected to result in better means for control of bark beetle through foliage spraying as a chemotherapeutant.

An educational program is needed to encourage communities to take action on the local level to halt further spread of Dutch Elm disease.

A panel under the leadership of Dr. Malcolm A. McKenzie, director, Shade Tree Laboratories, University of Massachusetts, Amherst, discussed "Some New England Insects and Diseases of Shade Trees and Their Control". Appearing with Dr. McKenzie on the panel, were Dr. David H. Marsden; Dr. Philip L. Rusden and Prof. William E. Tomlinson, all of the University of Massachusetts, Amherst.

Some interesting facts were brought out in the discussion. Dr. Marsden, talking on "Some Common Non-Infectious Diseases of Shade Trees in New England", said that plant pests should not be blamed entirely for all diseases on trees. Nature and man, and their influence on the environment, give rise to numerous non-infectious diseases of trees which challenge the skill of arborists in diagnosis and treatment.

These common, non-parasitic diseases include leaf scorch, needle blight, drought injury, frost injury, storm damage, gas injury, salt injury, spray burn and nutitional deficiencies, Dr. Marsden pointed out.

In his report to the NSTC,

secretary-treasurer L. C. Chadwick, Ohio State University, Columbus, pointed out that progress is expected soon by the group's Memorial Research Fund Committee under the joint supervision of the Department of Botany and Plant Pathology of the University of Maryland and the Division of Forest Pathology, Bureau of Plant Industry, Soils and Agricultural Engineering, U. S. Department of Agriculture. The group will study the control of shade tree diseases by

adding chemicals to nutient solutions applied to the roots, he said.

In summing up the work of the organization, he reported that the Shade Tree Conference's 1,203 members represent 45 states, the District of Columbia, Dominion of Canade, England, Guam and South Africa.

Commercial exhibits were on display at the hotel and in addition an educational exhibit attracted much attention. These included displays set

Go the scientific way...go MGK

AEROSOL INSECTICIDE CONCENTRATES

SPRAY INSECTICIDE CONCENTRATES

DUST INSECTICIDE CONCENTRATES

You may want complete formulas . . . ready to put right into your aerosol bombs or your retail packages. You may want combinations of insecticides and synergists that leave you only the minimum of processing to do. You may want to do most of the processing yourself and to you we offer the purest toxicants and synergists in their primary forms. MGK has the best of whatever you want. The emblem "MGK" is satisfying assurance of high efficiency and scientific production in insecticides and insecticide ingredients. Let this single experienced source help you make better products for less money. For complete information about MGK prices write 1703 Southeast Fifth St., Minneapolis, Minn.

> THE PIONEERS OF PYRETHRUM AND ALLETHRIN

Good insecticides protect America's health and harvest.

ING COMPA





Insistence on Integrity!

QUALITY control in pesticides doesn't just happen. It takes sharp, unsatisfied eyes that okay nothing until it has been tested and proved . . . and then okay only that batch. Questioning, testing eyes!

In our laboratories, careful routines are set up for mixing, measuring and packaging. More important, instructions to the chemists go something like this: "Take nothing for granted. Accept no rule of thumb. Approve nothing just because somebody else did." That is our quality control platform.

Tell your customers about this. Sell quality when you sell Thompson-Hayward pesticides. You'll sell more?



Automatic titrator used in titrations.

Control No. 17527 Tested by 3

This product has been carefully examined and found to conform with our specifica-

In case of correspondence regarding this shipment refer to the above control number.

This certifying tag must be on every container of Thompson Hayward chemicals. THOMPSON-HAYWARD CHEMICAL CO. Kansas City 8, Mo.

MINNEAPOLIS . OKLAHOMA CITY

N. LITTLE ROCK . SAN ANTONIO

DES MOINES . DAVENPORT

ST. LOUIS . HOUSTON

THOMPSON-HAYWARD

CHEMICAL COMPANY

KANSAS CITY, MO.

NEW ORLEANS, LA.

DALLAS . WICHITA

MEMPHIS . CHICAGO

OMAHA . DENVER

TULSA

up by the University of Massashsutts, University of New Hampshire, Harvard University; the Massachusetts Audubon Society and the U. S. Department of Agriculture.

NIAGARA

(Continued from Page 49)

ager, supervising sales throughout the midwest and central south. In 1937, he became assistant sales manager and moved to Medina, New York, where his residence has been maintained since. He is president and director of Quimica Sinaloense S.A. e C. V. of Culiacan, Mexico, a subsidiary; also a director of Pine Bluff Chemical Co., Pine Bluff, Arkansas, and Niagara Brand Spray Company, Ltd., Burlington, Ontario, Canada. During the past several years Mr. Vernon has served as president of the board of education of the Medina schools.

B. Earle Vosteen, vice-presient and controller of Niagara Chemical Division and assistant secretary of Food Machinery and Chemical Corporation, joined the Niagara organization in 1928 and moved to Medina, New York from Buffalo, where he was controller of the Beaver Board Companies. He is also an officer and Director of Quimica Sinaloense S. A. de C. V., Pine Bluff Chemical Co., and Niagara Brand Spray Company, Limited In addition to his duties as controller, Mr. Vosteen is financial officer of the division and acts as administrative assistant to the president.

Stuart H. Bear, a graduate of Pennsylvania State College, in entomology and plant pathology, joined the Niagara organization as a salesman in 1931, working in the Spencerport, New York area. He was later transferred to the midwest, where he served as territory manager until 1937, and then became district sales manager, in which capacity he served until called to Middleport as assistant sales manager of the Division in 1947. Mr. Bear will become vice-president and sales manager of the Division. He is also an officer and director of Quimica Sinaloense S. A. de C. V.

Niagara Chemical Division has one of its major plants located at Middleport, New York and there maintains its administrative and research headquarters. Other production plants operated by Niagara are at Aye, Massachusetts; Jacksonville, Florida; Belle Chasse, Louisiana; Greenville, Mississippi; Harlingen, Texas; Pecos, Texas; Richmond, California; Yakima, Washington; Pompano, Fla.; and Tampa, Fla.

In addition to the above, the

following subsidiaries of Food Machinery and Chemical Corp. are operated by the Niagara Division: Quimica Sinaloense S. A. de C. V., Pine Bluff Chemicals Co., Niagara Brand Spray Company, Limited.

The Niagara Chemical Division manufactures, formulates and distributes agricultural chemicals throughout the United States, Canada and Mexico. It also serves the export market in scores of foreign countries.

VELVEX CLAY

In making organic concentrates using benzene hexachloride, chlordane, toxaphene, and other similar materials, it is important to have the concentrates free flowing.

VELVEX Clay can be combined with more costly diluents, such as Fuller's earth, and the result will be a free-flowing concentrate, at a lower cost to the producer.

VELVEX Clay has the following advantages:

NON ABRASIVENESS FINE PARTICLE SIZE ABSORBTIVENESS PROPER BULK

HIGH INSECTICIDAL VALUE OF CLAY ITSELF
LOW PH VALUES
NO PHYTOTOXICITY TO PLANTS
OUTSTANDING ABILITY TO STICK TO THE LEAF

For Further Information or Samples Write to

SOUTHEASTERN CLAY COMPANY

Aiken, South Carolina



S/W Intimate Blending Equipment—9 Ways Better

Sprout-Waldron revolutionized the manufacturing of insecticides by developing the Intimate Blending System. And in a great variety of installations, this Intimate Blending System proves itself 9 ways better:

- 1. Improves product
- 2. Increases efficiency
- 3. Saves on labor cost
- 4. Reduces operating cost
- 5. Cuts maintenance

that are imperfectly blended mean incomplete cover-

age, low field efficiency and lost business.

- 6. Removes dust hazards
- 7. Improves sanitation
- 8. Operates more quietly

9. Improves working conditions

Smaller Streamlined Units-For Localized Operations

To meet the needs of localized operations, Sprout-Waldron has developed smaller, streamlined units, designed to meet the needs of such installations. For all types of insecticide blending operations, ask your SPROUT-WALDRON MAN to show you how you can improve products, safeguard personnel, and increase profits. Write for Bulletin 1-846 today!

Sprout-Waldron & Co., Inc., 7 Logan Street, Muncy, Pennsylvania



SPROUT-WALDRON

The Best in PROCESSING EQUIPMENT Since 1886

Bartlett Executive Dies

Clinton C. Lawrence, 59, executive vice-president and a director of the F. A. Bartlett Tree Expert Co., died August 18 at his home in Stamford, Conn. He was a graduate of Cornell University in plant pathology and during World War I, was a second lieutenant in the Army Air Corps. In 1927, he helped found the Bartlett School of Tree Surgery.

OHIO TOUR

(Continued from Page 77)

of aphis. The methoxychlor-treated plots were the first to show infestation by aphis and the two-spotted mite. None of the materials were phytotoxic, with the exception of NPD, which caused some injury to potatoes. In general, "Malathon" was not quite as good as parathion, and dieldrin was found best for flea beetle

Applications of various fungicidal compounds to potatoes in potato disease control showed SDDC plus tine sulfate at 4-1-100 to give the best results. DDT was used with the formulations at 2/3-100.

Apple scab control studies, using organic mercury fungicides "after infection", were reviewed by H. F. Winter of the experiment station. None of the mercury-treated trees required further fungicide treatments, due to the good initial control mangate plus sulfur treated trees did need extra fungicide treatments. "Coromerc" was found equivalent to "Tag" and "Puratized" in "after infection" applications. "Manzate" treatments seemed to cause some damage by causing a burning of the fruit. Table 2 summarizes some of the preliminary results in apple scab control, on McIntosh variety.

Mite and aphis control on apples indicated that in severe cases of rosy apple aphis, dormant oil sprays are not adequate. Parathion has been most useful against this pest, but it is recommended that application be delayed after the "pink" stage, to allow the parathion to be effective also against other insects. Lead arsenate and DDD are used primarily against red banded leaf rollers. Studies of

the green aphis, indicate that "Systox" is a good aphicide, but does not act as a true systemic on this species.

Among the experiments conducted was a treatment with the "Ukako"* unit . . . against green aphis and other plant pests. The results, seen in table 3, are an obvious indication that the so-called "electronic treatment" is without effect.

In an informal program on dealer problems, M. G. Farleman, Standard Oil of Ohio, reviewed some of the difficulties encountered in selling to the farmer, and particularly some of the inconsistencies in results with products in use. D. L. Kent, Goodrich Chemical Co., Cleveland, pointed out the need of an educational program directed to the actual user, which issue centers on the direction of information to publications reaching the user and dealer. Films on soil and soil conditioners concluded the session.**

OA reprint of an article explaining the use of the Ukako unit, its theory, and experiments involving its use, is available on request from Agricultural Chemicals, 175 Fifth Ave., N.Y.C.

Toxaphene Toxicity Studied

REPORT prepared by the medical director of Hercules Powder Co., Wilmington, reviewing the toxicity hazards in connection with the use of toxaphene, and giving suggestions to physicians on the treatment of persons suffering from accidental or careless misuse of toxaphene-containing insecticides, has been receiving considerable attention from newspapers and radio commentators over recent weeks. And as so often results when a company makes a laudable effort to get needed information of this type into proper channels, the uninformed newsmen and their radio counterparts have picked up only part of the story, - the sensational part of course, - and have tended to magnify the toxicity hazard in connection with use of toxaphene insecticides.

The report itself pointed out that effective insecticides must often necessarily be poisonous in nature "and must have lethal qualities if they are to be used in controlling insect pests. As with other economic poisons used in agriculture, one may expect that increasing use of toxaphene will be accompanied by a few rare instances of human poisoning through accidents and careless misuse of the agent."

The complete absence of poisoning in workers who manufacture toxaphene and in those who are exposed to it, both in the compounding of pesticides and its use in agriculture, are cited, however, in the report to indicate that toxaphene can be handled safely.

In the four most serious case histories described in the report, the insecticide was carelessly left where it was found by small children who ate or drank it. In two other case histories, illness resulted when the insecticide was not properly used. All the persons involved in these cases recovered.

In general, the news commentators have ignored the sections of the Hercules report which caution against careless use of insecticides if such tragedies are to be avoided. They



F. S. ROYSTER GUANO COMPANY Norfolk, Virginia



Nopco's AGRIMUL^e series of emulsifiers for Toxaphene, Chlordane, and other agricultural chemicals cover emulsifier needs for a wide variety of formulations.

Three distinct types are available: nonionic, anionic, and combination—providing the formulator with versatile means for solving problems relative to water conditions, coverage, and adhesion.

Outstanding Advantages of AGRIMUL Emulsifiers:

EASE OF EMULSIFICATION—Emulsions are easily prepared from Toxaphene, Chlordane, BHC, Aldrin, and other polychlor concentrates.

STABILITY—Emulsion-type insecticides adequately stable for agricultural uses are easily and economically prepared.

ANTI-CORROSIVE PROPERTIES—AGRIMUL emulsifiers with anti-corrosive properties minimize insecticide contamination from containers.

Obtain a copy of Nopco's technical bulletin describing AGRIMUL emulsifiers, and giving formulas, by filling out the attached coupon and mailing it today.



NOPCO CHEMICAL COMPANY

Harrison, N. J.

*Reg. U. S. Pat. Of.

-	
	NOPCO CHEMICAL COMPANY
	Harrison New Jersey
	Gentlemen
	Please send me your Bulletin describing Napco AGRIMUL Emulsifiers for Polychlor Insecticides and Herbicides and giving formules.
	NAME
	COMPANY
	ADDRESS
	CITY Zone State

have also failed in many cases to call attention to the fact that the reported poisonings resulted from carelessness and misuse, and have tended merely to stress the idea that toxaphene is pretty dangerous stuff if chomped down diluted only slightly with green vegetables. Where the commentators got their stories is still not completely clear, as they are not normally partial to reading lengthy medical reports. It is indicated however that perhaps a press release issued by the American Medical Association may be the culpable source.

The A. M. A. itself in the July 19th issue of the Journal of the American Medical Association (pgs. 1124-1126) carries the full text of the report under the title "Accidental Poisoning By Toxaphene," the authors being L. C. McGee, M.D., and Howard L. Reed, M.D., medical director and assistant medical director of Hercules Powder Co., and Dr. James P. Fleming of Hearne, Texas. The report is summarized as follows:

 Three deaths in children presumably resulted from accidental ingestion of lethal amounts of an insecticide, toxaptene. A fourth child so poisoned re-

Two instances of group poisoning, that followed ingestion of toxaphene-sprayed collards and chard, respectively, have come to our attention. Among 10 persons known to have eaten the contaminated greens, in 7 symptoms developed consistent with toxaphene poisoning as seen in laboratory animals.

 The onset of poisoning characteristically was abrupt and not associated with abdominal pain, involuntary vomiting, or diarrhea. Illness is manifested by a skeletal muscle response to central nervous system stimulation and by loss of consciousness.

 These instances of poisoning in humans have resulted from accidents or injudicious use of the pesticide.

5. Since similar poisoning may occur in the future, it is essential that physicians recognize the cardinal symptom, a convulsion, and the importance of countering the acute central nervous system stimulation with barbiturates.

6. The complete absence of poisoning in workers who manufacture toxaphene and in those who are exposed to
it, both in the compounding of pesticides
and its use in agriculture, suggests that
it can be handled with safety.
7. This and other effective pesti-

7. This and other effective pesticides are known poisons and must have lethal qualities if there is to be progress in man's war against insects. Such chemicals should be used properly and with precautions against human consumption if tragedies are to be avoided.

To Michigan Chemical Post



ALFRED G. RAUFER

Alfred G. Raufer has been appointed sales director of Michigan Chemical Corporation, Saint Louis, Michigan, effective August 1, 1972, according to an announcement by Roland P. Place, president.

During the past three years, Mr. Raufer has been Michigan Chemical's eastern sales manager. He was for ten years with the Sherwin Williams Company as eastern sales manager of the

household insecticide division, and has a broad background in sales work. Mr. Raufer will make his headquarters at the Corporation's main office at Saint Louis, Michigan.

Joins Development Group

H. A. Hashbarger, development manager of Monsanto Chemical Company's Foreign Department, has joined the development department of the company's Organic Chemicals Division, the company has announced.

Mr. Hashbarger, who has been with the company since 1937, will coordinate Krilium technical information developed by the various departments of the Organic Chemicals Division and will assist A. T. Loeffler,
divisional development department director, in communicating findings
both within and outside the company.

"FIFTH PLATE"

(Continued from Page 59)

an average year would make unmarketable most of the apples grown

in Eastern orchards. Fungicides are widely used to prevent the disease, cherry leaf spot, that completely defoliates trees by mid-summer, reducing the crop that year and sometimes killing the trees. Fungicides have increased the output of anjou pears by 300 million boxes a year. Those, though, are drop-in-the-bucket savings compared with the results yet to be obtained. Fungus diseases are doing an estimated \$4 billion worth of damage a year, and to make a real inroad into this loss will take many such developments as the recently announced use of antibiotics to control halo blight of beans.

We use fumigants to protect our stored grains from destructive and contaminating weevils. We could use a lot more. There has been promising research with fumigants in the control of nematodes and other soil pests. Cost has been a deterring factor in the widespread use of soil fumigants, but if they increase yields of valuable crops as much as the research indicates they will, we can look forward to an

For Export from America's Leading Chemical Manufacturers




- Selective and General Type Weed Killers
- Organic Insecticides
- Herbicides
- Fungicides
- Rodenticides
- Fertilizers, and Allied Chemical Products

WALLACH - GRACER EXPORT CORP. MANUFACTURERS REPRESENTATIVES

55 WEST 42nd STREET, NEW YORK 36, N.Y., U.S.A.

Tel: LO 4-5490

Cable: WALGRAY



BY eliminating slow, costly manual operations, Johnson fertilizer blending plants profitably increase production output . . . and earn important savings in manpower. For example, on one typical Johnson installation, material travels via bucket elevator to clod breaker, where it is pulverized, then carried on belt conveyor to separating screen.

Collecting hopper under screen feeds the pulverized material to a pivoted distributor which charges a sectional 5-compartment bin. Multiple-material batcher accurately weighs 5 (or more) fine-grained materials and discharges batch into mixing unit for final blending operation.

Whether you are interested in complete plant installations, manual or fully-automatic, or need auxiliary equipment to modernize your present facilities, it will pay you to see your C. S. Johnson Co. distributor . . . ar send coupon today for more complete information.

25% for a large midwestern fortilizer manufacturer . . . and, at the same time, greatly reducing menpower requirements in the plant

Photo below, of one of Latin Ameri-

facturing plants, shows happer, plv-

ated distributor and 8-section bin

ca's most modern fortilizer man

#	1	
12		
1 1 1		

CO. CHAMPAIGN, ILL

☐ Send us mor	re date or	Johnson 1	ertilizer	blending	plants.	☐ Heve	Johnson	distributor	coll.
NAME				TITLE					-
COMPANY				DIV.					
STREET									
CITY				STAT	E				AC
Also interested		k phasphate ket alevator							

expanding use of these chemicals. It may be assumed, too, that when markets increase, costs can be reduced -as so often happens in our economy.

Insecticides offer perhaps the greatest potential for saving the crops we produce on our farms. In many cases, we already have the insecticides, and the control methods. We only lack their widespread application on the farm. I believe insecticides might give us the fastest and most far-reaching food production increases of all the pesticides.

What can be done with insecticides in a few, limited fields has already been shown. Simply controlling the codling moth in apple orchards with one of the newer insecticides, for example, reduced damage to fruit from 15 percent to about 3 percent and cut down the number of yearly coverage sprays required to 3 or 4 where once as many as 10 to 12 were required.

Grasshoppers are another pest that we now successfully control. If you can recall the '30's, you'll remember that 'hoppers teamed with drouth to destroy the soil cover and cause the loss of millions of tons of our finest midwestern topsoil. And yet, how many realize that we could have had a similarly heavy outbreak of the pests these past few years, if it hadn't been for research and insecticides?

Last year, Oklahoma farmers saved nearly 2 million acres of small grain from the greenbugs by using chemical sprays. But still the aphids got 1.5 million acres of their crops. Insecticides as residual sprays are being used in 50 million homes around the world to control malaria-carrying mosquitoes. The insecticidal control of horn flies on livestock is currently saving us millions of dollars worth of beef alone, each year.

An extensive control program conducted by the Bureau of Entemology and Plant Quarantine and cooperating states has rid Pennsylvania and western New York of the forestdestroying gypsy moth. The use of new agricultural chemicals that can be put on effectively with airplanes, has reduced control costs to less than a dollar an acre - less than 1/25 of that of a few years ago.

The agricultural chemical industry can point with much pride to these records, but there is so much yet to be done. There are more than 6,000 destructive insects in the U. S. chewing and sucking away, for instance, 15 percent of our cotton and 20 percent of our sugar cane every year. The European corn borer and the corn earworm cause millions of dollars worth of damage in the midwest, ruining feed that we need for our livestock. Stored grain weevils are taking from 5 to 10 percent of our stored grains from us.

Hints of what the future holds for some of these 6,000 pests comes to us from experiment station and USDA reports. Iowa and Kentucky, for example, have found that corn yields can be increased 10 to 40 percent by controlling rootworms with insecticide. Cooperative work between the USDA and Virginia showed that the use of insecticides on the soil to courtrol rootworms in peanuts would increase yields 40 percent.

As we continue to expand our

grasslands program, the place of insecticides — and other agricultural chemicals — gains importance. Some of the new insecticides for example, have raised yields of alfalfa and clover by more than 30 percent, and, at the same time, improved the feeding quality of the forage. This year, farmers have bettered their production on more than 275,000 acres by controlling a single, serious, forage crop pest —the spittlebug. Insecticides saved nearly a half million acres of valuable range grass from grasshoppers in


It's apparent that in the years to come, agricultural chemicals will help the farmers fill the fifth plate. It is equally evident that we're all going to have to pitch in together to get the job done.

Our state colleges and the U-S. Department of Agriculture will help by carrying out fundamental research in the development of pesticides and by determining how those already available can be used best on the farm. Pesticide people will help by supplying the bulk of the

necessary industrial research in the development of new and better pesticides, and by producing them in quantity for distribution throughout America and other countries.

A study group — of which L. S. Hitchner was a member — was set up by Secretary Brannan about a year ago, to analyze and suggest changes in the Department's insect and plant disease control programs. I can do no better than sum up some of their conclusions to indicate to you possible ways we can bulwark our already strong cooperative ties with one another.

This study group recommended that surveys of insect and plant disease infestation be unified and intensified by the federal government, in cooperation with State and local authorities. The group felt that timely surveys that would quickly focus the need for active measures of control or protection in certain areas, could save us much in total production. I can say that we in the Department are working toward the fulfillment of that recommendation. The success of



THE AMERICAN POLYMER CORPORATION

Announces

COMMERCIAL PRODUCTION OF

ACRYLIC SOIL CONDITIONERS

AVAILABLE IN CARLOAD QUANTITIES AS

AGRILON NA 15% AQUEOUS SOLUTION AGRILON N DRY-POWDER

Offered to Manufacturers & Jobbers of Agricultural and Garden Supplies

SAMPLES AVAILABLE ON REQUEST



· AMERICAN · POLYMER CORPORATION

General Offices: 101 FOSTER STREET, PEABODY, MASSACHUSETTS

the cooperative grasshopper control program in Wyoming last year was based on information obtained in such timely, accurate surveys. Today, the USDA is coordinating the efforts of farmers, county agents, and entomologists all over the country to get evidence of local insect and plant disease outbreaks. We believe that in time the value of what is being done now will bring further public support of survey and control activities.

This study group emphasized, too, the need for our agricultural extension services in the various states to take more of the pest control information to the farmers who need and can use it. They believe that losses to insects and diseases were too great and were repeated too often.

It is important that we work together in this educational program not only to inform the farmers how to use pesticides effectively and safely, but also to enlighten the general public about agricultural chemicals—particularly, pesticides. We must inform everyone of the real value of these chemicals and how to use them safely.

A good deal of misinformation about insecticide is being circulated. Some people, on the basis of unfounded fears, are pressing for the establishment of unnecessary limits and even prohibitions against the use of chemicals that are important to farm production and which are safe when properly used. Even automobiles are unsafe when improperly used. The answer is education — not withholding of useful products.

Our objective should be — as the study group recommended — that the farmers and ranchers of this country should so well understand the value of pesticides to them, that they will willingly accept the responsibility of pest control — on their own farms — and in the communities in which they live.

These are some of the tasks before us.

The fifth plate can be filled. We can fill it pretty well through use of insect control alone — and pesticides are a major insect control tool. Practically speaking, we know of course that the job requires the co-

ordinated use of all available tools and that it requires economic conditions favorable to production. Certainly all of us who serve farmers have our work cut out for us.

The pesticide industry's service to agriculture has barely begun. As the nation demands greater production from the farmer, he in turn will demand more agricultural chemicals. Each time he uses agricultural chemicals to build bigger yields and to protect them, he uses with greater efficiency his land, his machinery, his power.

Yes, agricultural chemicals have a great future ahead of them—a future that is wrapped up in the American farmer's ability to produce for the fifth plate.*

HEPTACHLOR

(Continued from Page 37)

garden web worms for which 0.35 lb. actual heptachlor is recommended per acre; and cutworms, for which the recommendation is 0.7 lb. per acre.

Experimental programs are nearing completion regarding the use of heptachlor formulations for the control of insect pests other than those attacking cotton. Promising results have been reported in regard to heptachlor's control of the following pests. The timing of application, dosage and residue levels require further consideration before label registation.

During the past several years, heptachlor formulations have been supplied for experimental evaluation in the control of varied insect pests. Heptachlor formulations are now being tested against the following insects and show effective control with the use of minimal quantities:

Cattle Lice
Cherry Fruit Fly
Corn Billbug
Corn Earworm
Darkling Beetles
Earwigs
Engelman Spruce Beetle
Eye Gnat
Lesser Corn Stalk Borer
Nematodes

Pecan Weevil
Poultry Mite
Squash Bug
Strawberry Crown Moth
Strawberry Root Weevil
Termites
Wheat Stem Sawfly
White Fringed Beetle

Phytotoxicity

IN the course of the heptachlor experimental progress, observations were made to determine evidence of plant injury. No plant injury has been observed or reported using the various heptachlor formulations at different concentrations.

Off-Flavor: Vegetables and fruits treated with heptachlor during the growing season or grown in heptachlor-treated soil were tested for flavor and taste at harvest time. No off-flavor or changed characteristics in taste could be detected.

Analytical Methods

ASPECIFIC method for the determination of heptachlor residues has been developed and is described in basic detail in "Detection and De-

MAGNESIUM SILICATE

ALUMINUM SILICATE

FOR INSECTICIDES USE

Guaranteed Dryness

TALC

Controlled PH.

The Ideal Filler & Diluent

EASY DUSTING

(Softness, Slip, Clingability, Dry)

AIR FLOATED

(Up to 99.5% 325 Mesh)

PACKED IN SO LB. PAPER BAGS - AVAILABLE IN CARLOAD OR TRUCK LOAD LOTS

SAMPLES UPON REQUEST

GEORGIA TALC COMPANY, INC.

CHATSWORTH, GEORGIA

P. O. Drawer 278

Incorporated 1907

Phone 4431

PROMPT SHIPMENT AT ALL TIMES

8 MORE processing problems solved by Beckman Spectrophotometry!

Now saving time and money in 8 operations, the Beckman Model DU Flame Spectrophotometer with accessories used in this true case history was originally bought for only one job. No other single analytical instrument, with the possible exception of the Beckman Model B Spectrophotometer, is capable of carrying out this highly varied list of uses . . .

1. Rejects immature Grapofruit
Real Gold Citrus Products rejects immature
batches and maintains high quality of grapefruit
by determining naringin content. This bitter factor, formerly measured on an inaccurate taste
basis to judge immaturity, is now found accurately
with the Beckman Model DU.

2. Controls Shut-Down Time

2. Ceatrels Shut-Dewn Times Because besperidin content of orange juice varies seasonally, this company was previously unable to predict shut-down time to remove precipitant coatings from heat-exchanger surfaces. Time is now accurately computed and production control is more precise with Beckman Model DU.

is more precise with neckman Model DU.

3. Higher Quality = Lower Cost
Controlling Cu and Fe content of sugar syrup to
few parts per million prevents reduction of Vitamin C in orange base canning. Real Gold maintains long life of Vitamin C in storage by rejecting all raw materials with excessive Fe-Cu levels.
Rapidity of analysis with Peckman Model DU
Flame. Spectrophotometer speeds production at
lower cost.

AOAC standard method used with Beckman Model DU to control citral quantity on every shipment of lemon oil demonstrates increasing versatility of this precision instrument.

6. Gulacturonic Acid Study High galacturonic acid in citrus is believed to be an indication of degree of ripeness. Fruit maturity controls are now being studied using Beckman Model DU.

7. Storage Controls

7. Storage Controls Evaluating ageing by color change in juices in storage is now simplified with Beckman Model DU Spectrophotometer and Reflectance Acces-

6. Determines Sall Needs

Research study is now underway to evaluate or-chard soil needs by determining mineral content of ash from juice. Objective—to imure higher quality at growing points.

FOR MORE COMPLETE DETAILS ASK FOR DATA FILE 33 - 15



BECKMAN INSTRUMENTS used

> BECKMAN INSTRUMENTS control modern industries

Madel DU Flame Spectraphatometer

oviding high resolving power for quantitative determinations more than 40 elements, the Beckman Model DU Flame giver al Gold Citrus Products law initial cost and law operating at (non-technical personnel).

Model DU Reflectance Accessory Set

Because citrus solutions contain too much colloidal foreign matter to allow occurracy with transmitted light—the Beckman Reflectance Accessory Set gives accuracy for measurements of diffuse reflectance from apaque, colored samples in liquid, powder or solid form.

BECKMAN INSTRUMENTS, INC.

SOUTH PASADENA, CALIFORNIA Factory Service Branches: New York - Chicago - Los Angeles

ASCRIMAN INSTRUMENTS INclude: all Meters and Electrodes - Spectraphotomotors - Radioactivity Motors - Special Instruments

termination of 1.4.5.6.7.8.8-heptachloro-3a,4,7, 7a-tetrahydro-4, 7-methanoindene" by Polen, Percy B. and Silverman, Paul, in Analytical Chemistry, 24, 732-735, April, 1952. Heptachlor residues in the order of 0.01 ppm and less are determined by this

Heptachlor as the technical grade and in the commonly-used formulations may be analyzed by a method based upon its reaction with silver nitrate or silver acetate in strong acetic acid solution. Under such conditions the method is specific and may be used even in the presence of other insecticides such as DDT, benzene hexachloride and toxaphene.

Crop Residues: Throughout the 1951-52 season Velsicol heptachlor has been used experimentally in the control of insects on the following crops; cantaloupe, alfalfa, range and forage, beets, cabbage, carrots, celery, cherries, corn, cotton, lima beans, onions, peanuts, peas, potatoes (white) potatoes (sweet), strawberries, yams, turnips and rutabagas. Soil analysis was also included.

As soon as these crops are harvested in the various parts of the country complete residue analyses will be obtained. At the present, residue analysis tests on a portion of the crops already completed, indicate that the residues will be in the range of 0.01 ppm or less. These results are being obtained by both the chemical analysis and bio-assay methods.

Toxicity: Like all chlorinated hydrocarbon insecticides, heptachlor should be handled and applied with normal precautions to avoid any possible contamination. Lehman, A. J. in Association of Food and Drug Off., Quart., Bulletin XV (No. 4) 122-133 (1951) reports the LD 50 for heptachlor to the rat as 90 mg./kg. with symptoms of poisoning similar to aldrin, dieldrin and (technical) chlordane. **

Amendment Proposed

The following notice of proposed amendment to the regulations for the enforcement of the Federal Insecticide, Fungicide, and Rodenticide Act is being published in the

Federal Register. Interested persons may file their views within 30 days after the publication of the notice, according to W. G. Reed, Chief, Insecticide Division, USDA, Washington, 25 D. C.

 Section 162.10 would be amended by adding thereto new paragraphs (h) and (i) to read respectively:

(h) Duration of registration. Unless cancelled in accordance with the provisions of section 4. c. of the act or with the acquiescence of the registrant, or unless continued in effect in accordance with the provisions of paragraph (i) of this section, the registration of an economic poison shall be cancelled at the end of a period of five years following the date of registration of such an economic poison or following the date of any subsequent registered change in formula or labeling or following the date of any continuance of registration pursuant to paragraph (i) of this section: Provided, however, that prior to any such cancellation the Insecticide Division shall send to the registrant a notice of intent to cancel, and, in the event such notice is not sent to the registrant 30 days prior to the expiration of the five year period, the registration shall remain in effect until 30 days following the date such notice has been sent to the registrant at his latest address submitted to the Insecticide Division

(i) Continuance of registration.

If a registrant desires to continue the registration in effect, he shall notify the Insecticide Division in writing and it shall be continued in effect under the same terms as the original registration; Provided, however, that if, on the basis of information available at the time, it appears that the product or its labeling fails to comply with the act the registrant shall be notified and an opportunity given to make the necessary corrections. If the corrections ae not made, continued registration without protest shall be refused but registration under protest as provided in section 4. c. of the act shall be issued if requested in writing by the registrant.

2. Paragraph (2) of section 162,-17 (b) would be amended and new paragraphs (3), (4) and (6) would be added to section 162.17 (b) to read, respectively:

(2) If an economic poison is to be tested for a use which is likely to result in a residue on or in food or feed, a permit for shipment or delivery will be issued only when

(i) The food or feed product will not be used for food or feed except for laboratory animals, or

(ii) Proof satisfactory to the Director is submitted by the applicant that the proposed use will not result in a residue which is hazardous to man or other animals.

(3) A permit for shipment or delivery of any experimental economic poison for testing in any place likely to be frequented by people will be granted only if it is clearly shown in the application for such permit that proper safeguards will be taken to prevent injury to all persons concerned.

(4) All applications for permits covering shipments for experimental use shall be filed in duplicate and must be signed by the shipper or the person making the delivery and must contain the following:

(6) An economic poison intended for experimental use shall not be offered for general sale by a retailer or others, or advertised for general sale.

3. Section 162.17 ((d) would be amended to read:

(d) Cancellation of permits.— Any permit for shipment for experimental use may be cancelled at any time for any violation of the terms thereof or if it shall appear to the Director that the permit should be cancelled for the protection of the rubbic.

TECH. BRIEFS

(Continued from Page 73)

It has been established that greater uniformity of the application of treating materials has been obtained by the use of the spray treater rather than by the slurry or dry dust



JAITE BAGS

OFFER DEPENDABLE PROTECTION FOR YOUR FERTILIZER

Scientifically controlled forest to finished bag.

Sales Offices; Jaite, Ohio. Chicago, Illinois. New York, N. Y. Philadelphia, Penn. An old established concern, centrally located for fast delivery. Almost half a century of paper and paper bag experience stands behind our reputation for production of the highest quality.

THE JAITE COMPANY

"Manufacturers of Paper and Paper Bags Since 1905"

JAITE, OHIO



- SEWN VALVE
- PASTED OPEN MOUTH
- PASTED VALVE
- SEWN OPEN MOUTH

safe-use

safe-use is founded on research

Urge That Label Directions Be Followed Carefully



NATIONAL AGRICULTURAL
CHEMICALS ASSOCIATION

910 SEVENTEENTH STREET, N. W.

WASHINGTON 6, D. C.

AGRICULTURAL CHEMICALS

methods of treating. Since the treating materials used are not water soluble, they are applied as suspensions with the spray treater. Limitations in this method of treating can become serious if due caution is not exercised in the purchase of treating materials. Since dosages of treating materials and the quantity of moisture added to the seed are critical, the orifice size of the spray nozzle is limited. The particle size (5 microns or less) of these materials is, therefore, a definite factor to be considered. Under California conditions, it has been determined that 4% moisture added to processed sugar beet seed in treating is about the maximum allowable quantity. It would be hazardous to apply a greater quantity of moisture in treating because of possible adverse effect on good packaging, on safe tolerance in increased seed size from swelling, and on germination during extended periods of storage,

"The technique of preparing treating materials for spray application to processed sugar beet seed is quite simple. The recommended

dosage of "Phygon" is 0.25% on the seed, by weight, and 0.33% for "Isotox Seed Treater." Expressed in weight quantities this is equivalent to 4 ounces of "Phygon" and 5.3 ounces of "Isotox Seed Treater," respectively, per cwt. of seed. The recommended maximum moisture to be added is 4%. Therefore, the above quantities of treating materials plus any dye solution must be contained in two quarts of water (the slight displacement by treating materials is not considered serious). It is suggested that this maximum allowable amount of water be used per unit of 'mix' so that a larger-size nozzle orifice can be used.

The quantity of dye used is 3 to 5 grams per cwt. of seed. A good 'punchy' color can be obtained with 5 grams of the dye mentioned. Using more than this quantity is not recommended since the dye apparently has some influence in increasing the amount- of foaming which occurs when the 'mix' is being prepared. (Control of foaming will be discussed later.)

The Holly treating plant makes use of a 25 gallon chemical mixing tank equipped with a manually operated agitator. The tank is located so that its contents can be discharged by gravity into the treater solution tank. In making up 'mix', about 10 gallons of water are run into the tank. One gallon of "Phygon" (weight 11.0 pounds) paste is then added and the contents thoroughly agitated. "Isotox Seed Treater" in the amount of 14.65 pounds is weighed up, slurried with a minimum amount of water and added slowly, with agitation to the "Phygon" paste suspension. One pint of concentrated dye solution containing 200 grams of Victoria Green Dye is then added. The volume is completed with water to 22 gallons. This quantity of 'mix' contains the proper quantity of materials to treat 4400 pounds of processed sugar beet seed. One to 2 ccs. of "Ortho Foam Inhibitor" is added and the contents of the tank thoroughly mixed for several minutes. The 'mix' must be filtered through a 60 mesh screen prior to passage into the

HAHN selfpropelled HI-BOY

THE HIGH-CLEARANCE MACHINE Designed Exclusively for spraying

The instant solution to your customers' insect and weed control problems. Here's the ONE unit designed exclusively for spraying. No other unit has the light, yet rigid frame and foolproof power unit construction—coupled with all-weather performance. It's the Hahn Hi-Boy for complete customer satisfaction and increased profits to you. The rigid tubular steel frame is reinforced at all points of strain. The Hi-Boy's 7½ h.p. engine provides ample power for propelling the unit and operating the Hahn heavy-duty pump. The Hi-Boy's boom is adjustable from 24" to 96" in height, ground clearance, 68". Controls, valves, gauge, are located within easy reach of the operator. Easily negotiates field too wet for heavier equipment. For high customer acceptance and for high sales, it's the high clearance Hahn Hi-Boy.

WRITE FOR COMPLETE INFORMATION

HAHN 315 NO. NINTH AVE.

HAHN fractor-SPRAYERS

The low prized, high quality WABASH SPRAYER, equipped with Huben Pump. Frant and free mounted medals.

The HAVY DUIT TRACTOR SPRAYER, complete with Minute Secon Hings, exclusive feature permitting appropriate to raise or lower beam while severee is in semeration



Lightweight, practical, handy, easy to operate! Efficiency and reliability proved by over 4000 users in every part of the world under varying operating conditions. For details on how you can cut YOUR Bag Closing costs with the FISCHBEIN Portable BAG CLOSER . . . write:

DAVE FISCHBEIN CO.

Industrial Sewing Machinery

DEPT. 8 D 38 GLENWOOD AVE., MINNEAPOLIS 3, MINN.



Infusorial Earth • Volcanic Ash

Consult Whittaker on all your Dust Carrier Problems.



AGRICULTURAL DEPARTMENT 260 West Broodway New York 13, N. Y. spray treater solution tank. Some foaming will always result, either with or without dye being added to the 'mix', and will impede seriously its flow through the screen. Therefore, the use of the foam inhibitor is recommended. The skimming-off is slow, sloppy and wasteful. Scrubbing of materials through the screen should not be permitted as this promotes serious operational delays from noz-zle-clogging during the process of treating. Both conditions are apt to occur if the operator does not control foaming.

As a matter of information, concentrated dye solution can be made by dissolving 200 grams of dye in one pint of methanol. Hot water can be used for the purpose, but the dilution will have to be increased so as to get the 200 grams of dye completely into solution. Methanol applied at the rate of one pint per 44 cwt. of seed had no apparent adverse effect on germination.

In any discussion of the use of a spray treater where orifice size of the spray nozzle is a limiting factor, too much stress cannot be placed upon the absolute necessity for a thorough screening of all treating materials, whether used singly or as a composite mix. The same should be said with regard to the cleanliness of the spraying equipment with particular attention being given to preventing the entry of any foreign materials into the seed treating solution tank and the component feed lines.

As a result of thorough investigations by the Divisions of Pathology and Entomology at the University of California, Davis, as well as from commercial data, it has been established that sugar beet seed treated with "Phygon"-Lindane can be stored for a period of at least 40 weeks without any danger to germination. It has also been found that lindane will remain at least 75% effective in the control of wireworms even after the treated seed has been stored a year.

In certain areas in California where processed sugar beet seed has been stored for any length of time, infestations of the Confused Flour Beetle (Tribolium) have been experienced. While the effects on germination from this beetle have not been serious, no sugar beet grower appreciates the presence of a numerous quantity of beetles emerging from a package of sugar beet seed. The treatment of sugar beet seed will prevent such infestations.

PLANNING AHEAD

(Continued from Page 31)

to make the necessary chemicals. Except for a few products, all basic raw materials were in ample supply to meet conditions of an extremely heavy infestation of pests. Distribution facilities, streamlined and expanded following the war, were in full operation and could have handled any eventuality in regard to protection of American crops.

The 1952 season, in all its aspects, is not in the least new to those who have been operating in the pesticides field for the past 50 years or even in the last 25 years. Some who entered the industry during the past 10 years, may not have



Announcing the formation of PIONEER CHEMICAL ASSOCIATES

A NEW ORGANIZATION OFFERING

THE CHEMICAL AND ALLIED INDUSTRIES



ITS SERVICES AS

CONSULTANTS

BROKERS AND

MARKETERS

And the appointment of

MR. J. NEWTON HALL

FORMERLY VICE-PRESIDENT OF

JULIUS HYMAN AND COMPANY

AS GENERAL MANAGER

EXECUTIVE OFFICES

3035 EAST COLFAX DENVER E. COLORADO



PIONEER CHEMICAL ASSOCIATES

WILL SPECIALIZE IN INSECTICIDES, OTHER AGRICULTURAL CHEMICALS AND RAW MATERIALS

anticipated the situation which occurred during the season.

When the sale of an industry's products bears largely on the vicissitudes of nature, there is no recourse, from a practical standpoint, except to build sufficiently large stockpiles of pesticidal materials to satisfy the extreme potential. No matter how carefully a production program is established, there are so many variable factors in the brief seasonal demand, that inevitably, over a period of years, carry-overs are bound to result. Some years, the carry-over stocks will be so large that the gains of the previous season may be submerged by warehousing costs and interest on invested capital.

If we are to remain in this business of production of agricultural chemicals, we must be willing to accept the ups and downs over an extended period. Our top management must realize that in weighing the merits of the pesticides industry as a business, consideration must be given to the variability of each season's operations. There are seldom

long periods of good profits in the pesticides industry. There are longer periods of medium profits graduating into small profits and losses. The mean for all profits, over extended periods, levels on a reasonable return.

Since it is practically impossible to ascertain a season's demand prior to the manufacturing period, only long-range planning by each company through a well-directed program, aimed at product development and varied outlets, can smooth out and stabilize the marketing of agricultural chemicals. One of the safety valves which can be used to cushion the effect of an off-season in domestic consumption is the export market. During the past season, many pesticide companies expanded old plants in foreign countries and built new ones. This does not mean that we should use the foreign markets for dumping excessive production in this country. It is our responsibility to maintain these markets continuously. During the early part of the current season, when the industry realized that stocks were not moving, every

effort was made to obtain relaxation of export controls on some products. Considering our present stocks, export markets are still open and can serve to turn over the present supply of pesticides so that new, fresh stocks will be available for next season's use.

Must Buy Early

BECAUSE our season of distribution is so short, a buy-early program is necessary in order that we move the 1¼ billion pounds of pesticidal chemicals in 30 to 90 days. Simply because weather conditions in 1952 did not produce an insect infestation which would rapidly drain off pesticides in the distribution chanton will occur in 1953. Our buyearly program must be rebuilt.

There is reason to believe that many of the goals set by the Defense Production Administration for the production of food, fiber and feed have been reached for the present. As our population increases, however, production goals will have to be set higher and higher. This will









NO SEWING





AUGER-M

Quickly fills self-closing bags to the desired weight. Saves time in filling and eliminates need for sewing or other expensive bag closing equipment.

The AUGER-MATIC Bag Packer has a multi-twist auger which can quickly be changed to pack any material from powder to pellets.

AUGER-MATIC Packing reduces dirt and waste . . . a time and labor saver. Packer can easily be moved from one filling point to another.

Send for descriptive literature.

E. D. CODDINGTON MFG. CO. 5036 NORTH 37TH STREET . MILWAUKEE P. WISCONSIN



THE LAMINATED BAG

Has All These Desirable Features:

Water-resistant

Weather-safe

Siftproof

Odorproof

Puncture-resistant

Acid-resisting

Grease-repellent

Contamination-proof

Burlap (or cotton) on the outside, paper on the inside-the two laminated together with a smooth, even film of asphalt (or other special adhesives) to make a strong, protective bag guaranteed to preserve the original fine quality of your product.

Write

Wire

Phone



Our

Nearest

Office

MENTE & CO., INC.

Isaac T. Rhea, Pres.

Box 1098 Savannah

Box 690 New Orleans Box 204 Houston

SPECIALISTS BAG SINCE take up our present productive capacity of pest control chemicals and cause further necessary expansion of the industry.

Conservation programs such as the grasslands improvement program now in force throughout the world present vast new markets for pesticides. There are some 800,000,000 acres of grasslands throughout the United States which can be improved through the use of chemicals. The prices of meat, milk and animal fiber are such that improvement of grasslands by use of chemicals is highly profitable.

Forest insect and disease control has been adopted as a national program and offers outlets for pesticidal materials. Soil conservation involving the use of grasses and legumes increases the production of seed - a high-value crop requiring the use of agricultural chemicals for profitable production.

The economy of our country is such that a farm-labor shortage now exists and will continue to worsen.

This problem has been alleviated to a large extent through the use of chemicals and mechanization. More and more reliance will be placed on chemicals to increase agricultural production on the limited number of acres available.

We need a program directed by top management which believes in merchandising. By merchandising, I do not mean solely the packaging of our products in an attractive manner, but merchandising in its broadest sense. New markets and uses must be developed; old markets should be expanded. Advertising should be directed toward education. Credit surveys permit an examination of the industry's credit practices and position in relation to other industries, and point the way to sound financial policies.

Acceptability of our products by the public is essential to continued operation of the industry. Educational programs familiarizing the public with the importance and necessity of these products in the production of food

and fiber are essential. The use of these products in the right way must be a part of this program.

A comparatively poor year in sales of agricultural chemicals does not signify a collapse of the industry. We are in a period of transition where our expansion, at the express request of government agencies, has overreached the current season's demand. The only limit for the expanion of sales of agricultural chemicals lies in the degree of the industry's will to search out new products, markets, expand old ones and educate the agriculturist in profitable use of pesticides through merchandising programs. There are many undeveloped markets available to the industry which can be tapped only through sound merchandising to create the demand for our materials.

As a source of information and coordination, we already have the machinery from which to produce such an industry-wide program the National Agricultural Chemicals Association.**



COMPLETE WEST COAST CHEMICAL SERVICE

ALDRIN, DDT, BHC, TEPP, DDD, DIELDRIN, CHLORDANE, TOX-APHENE, PARATHION, MALATHON, ARAMITE, K-6451, IPC, SODIUM CHLORATE, PENTACHLOROPHENOL, 2,4-D, 2,4,5-T, WARFARIN, POLYACRYLATES

(Micro-fine airmilled)

(Liquid concentrates and emulsifiables)

Custom Grinding • Mixing and Filling of Liquids • Grinding and Packaging of Powders (Wetteble powders and dust concentrates)

> 641 S. Fourth Street Richmond, California

HECKATHORN & CO.




WEST COAST DISTRIBUTION GRICULTURAL CHEMICALS

MANUFACTURERS AND DISTRIBUTORS

Complete line of Insecticides, Miticides, Fumigants, Fungicides Cotton Defoliants, Systemics, Soil Conditioners, Weed Killers, Fertilizers, Rodenticides

UNITED CHEMICAL COMPANY


Head Office: 641 South Fourth St., Richmond, Calif. Branches: Sacramento, Fresno, Bakersfield, Pomon



INTERCHANGEABLE ORIFICE TIP SPRAY NOZZLES

You have your choice of hundreds of tip sizes for any type of spraying . . . when you use TeeJet Spray Nozzles. Easy to change orifice tips give you completely different nozzles at very low cost. All TeeJet Spray Nozzles are precision machined throughout, Monel metal screen assures free





- Drilled and Milled arifice tips for super-precision . . , for exact valume control, uniform distribution, and long
- Interchangachle orifica tips . . . every spray type in complete capac-ity ranga from one gallen per hour and up.
- Eleven different spray angles from 0° to 150° in all capacities.



SPRAYING SYSTEMS CO. 3230 RANDOLPH STREET



Yes, MARIETTA has BIG ideas. and BIG concrete storage siles that offer large capacity, labor saving storage and materials handling facilities to meet your present and future needs.

MARIETTA concrete stove silos can be erected in any arrangement-line, cluster, or combined with existing units for added storage. They will protect your agricultural chemicals from fire, acid. air, maisture and the elements and reduce insurance rates. Require little or no unkeep expense. Can be equipped with conveyors and happers to facilitate handling, increase plant value.

Let a MARIETTA engineer put his ideas to work to increase your storage capacity and handling efficiency. Write, wire or phone for an appointment . . . TODAY.

The MARIETTA concrete corporation

- · MARIETTA, OHIO
- 509 Fifth Avenue New York 17, N. Y.
- Pulaski Hwy. at Race Rd.
 Baltimore 21, Md.



LINDANE

R. W. GREEFF & CO., INC.
10 ROCKEFELLER PLAZA, NEW YORK, N. Y.
1818UNE TOWER, CHICAGO, ILLINOIS

Berkshire

..... longtime specialists in magnesia for agriculture

Berk's EMJEO (80/82% Magnesium Sulphate) Calcined Brucite (fertilizer grade) 70/74% MgO Calcined Magnesite 90/95% MgO Other Fertilizer Materials.

INSECTICIDES - FUNGICIDES

Mercury Compounds for Agricultural Use

DITHIOCARBAMATES

Ferric - Zinc

Export

Berkshire Chemicals, Inc.

420 Lexington Avenue, New York 17, N. Y. Cable Address-"Berkskem" New York

Sales Agents for F. W. Berk & Company, Inc.

Import

Phillips Office Moved

The district sales office of Phillips Chemical Company has been transferred from Los Angeles to Pasadena, California, where it will be located in the Citizens Bank Building, 16 North Marengo Avenue. Robert B. Prock is in charge.

SUPPLIERS BULLETINS

(Continued from Page 69)

practiced.

"Livestock Pest Control" is now available for showings to livestock asociations, agricultural groups, schools, 4-H clubs and similar groups. Requests for showings will be filled in order of receipt and should be directed to the California Spray-Chemical Corporation, Richmond, California.

Miticide Described

A new extremely fine wettable powder is outstandingly destructive to mites in the egg stage. In fact, it is more effective as an ovicide than as an adulticide. It also has a long residual effect.

Called "Ovotron Wettable", it contains 50 percent p-chlorophenylp-chlorobenzenesulfonate.

Its toxicity to warm-blooded animals as well as to parasites and predators of crop pests is said to be low. It is also reported to have low toxicity to insect pollenizers.

Growers now have the opportunity to combine chemical and biological control in one operation. At the same time "Ovotran Wettable" is not hazardous to the operator.

It is chemically compatible with other insecticides and fungicides. For example, it is compatible with such chlorinated hydrocarbons as DDT, toxaphene, chlordane, methoxychlor and lindane, with phosphates, arsenicals and lime-sulphur. "Ovotran Wettable" is one of few materials compatible with alkaline mixtures.

To date, it has shown unusual miticidal effectiveness on citrus and deciduous fruit and nut trees, shade trees, ornamental shrubbery, cotton, melons, grapes and other crops. Widespread field tests have shown that this chemical is not injurious to these same plants and trees when it is used as suggested. With a few exceptions, it appears relatively nontoxic to plant foliage.

Dow Chemical Co. Bulletin, Midland, Mich.

ECONOMICS

(Continued from Page 34)

ceed. It is obvious that the earlier it can be recognized that a compound will fail, the less money will have been wasted on it, and the smaller will be the cost which a successful compound will bear. In Table III, a scheme has been chosen arbitrarily, whereby compounds are dropped out at various stages so that about one product in 1,800 eventually succeeds. This falls in line with our conclusion that only one compound in 2,000 screened is sold in any quantity. Remember that this is the

HLW EMULGATES

. . . THE ORGANIC METALS WHICH EMULSIFY

QUESTION: What is the Industrial position of H. L. Woudhuysen & Associates?

ANSWER: H. L. Woudhuysen & Associates is engaged in the successful and constant development of organo-metallic fungicide and nutrient emulsion concentrates through the application of a basic and unique process.

THESE PRODUCTS ARE EXCLUSIVE IN NATURE AND REPRESENT A NEW APPROACH TO A COMBINATION OF BASIC MATERIALS FOR PLANT DISEASE CONTROL.

QUESTION: What is the Commercial position of H. L. Woudhuysen & Associates?

ANSWER: H. I. WOUDHUYSEN & ASSOCI-ATES DOES NOT PLAN TO REACH THE RETAIL AND CONSUMER MARKET DIRECTLY.

H. L. Woudhuysen & Associates, therefore, is prepared to discuss distribution agreements with reputable processors, wholesalers, distributors and exporters, while free label arrangements could also be considered.

for complete information, write to



H. L. WOUDHUYSEN & ASSOCIATES

DIGBY 4-1857

factory at Long Branch, N. J

17 BATTERY PLACE - NEW YORK 4, N. Y.



This extra breathing protection against toxic insecticides is yours in ONE Respirator at no increase in cost! Recently accepted new type filters (Cat. No. CR-73488.

Carton of 100) do the job and they are interchangeable with your present M.S.A. Farm Spray Respirator,

DEALERS WANTED—Extra profits and business. Your present customers are your best prospects. Cartridges and filters mean repeat business. Write now for details.



MINE SAFETY APPLIANCES CO.

Braddock, Thomas and Meade Sts.
Pittsburgh 8, Pa.
AF YOUR SERVICE: 74 Branch Offices in the
United States and Conada



Potassium Nitrate

A quick soluble source of Potash and Nitrogen for use in the preparation of concentrated water soluble fertilizers or starter solutions.

Our private formula service is available for the preparation of water soluble fertilizers under your own label.

Davies Nitrate Co., Inc.

114 Liberty St

New York 6, N. Y.

Plant at Metuchen, N. J.



.... that give your chemicals full killing power . . . and safety

Accessories Mfg. Co.

720 McGee St.

Kansas City, Mo.

average experience and that any individual company may have an appreciably better or worse cost history on the chemicals it introduces.

Some saving over the cost presented is achieved by the dropping of compounds midway through field testing or development so that they do not use the full charge for that particular section. Of course, any accounting department would point out that it is necessary to assess against this expense an interest rate on the money expended until it was returned via profit. Another such cost is the not uncommon case of excess of cost price over sales price at the outset of production. There are other omissions but let us then use the figure of \$1,000,000 as the cost of research and development of an agricultural chemical, admitting that it is a guess. For the 50 new chemicals introduced in the last decade the cost would be \$5,000,000 per year. This is about 4% of the total present dollar volume sold. (High, but not out of line in an expanding field). There are at least

40 companies doing work in this field and this would figure to an average expenditure of \$125,000 per year.

There can be no question but that this expenditure has been and will be continued to be justified as far as the farmer is concerned. Cases of his increased yields per acre and improved quality of product due to these new chemicals are well known. I believe also that this research can be profitable to companies provided that : (1) They can afford to invest a total of \$1,000,000 or more over a five to ten year period before expecting a return on their investment: (2) They closely scrutinize and direct their work so that unsuccessful compounds are discarded as quickly as possible and that they do not work in fields where, if successful, the volume of the market is inadequate to return their investment; (3) They are prepared at the end of the five or ten years to invest the 2 to 10 million dollars necessary in plant equipment to take advantage of their research.

Fred Lodge, NFA, Retires

Fred S. Lodge, a recognized authority in the fertilizer industry and for the past 18 years connected



with the National Fertilizer Association, retired August 1 from active participation in the industry.

Mr. Lodge held the position of secretary-treasurer of the NFA and was at one time its acting president. His thorough knowledge of the in-

Announcement


We are proud to announce that "A.P.E." S.A. Applicazione Processi Elettrochimici, Genova, Italy has joined the many outstanding and internationally well known Chemical Manufacturers in appointing us their Export Agents for the sale and distribution of:

Triple Mineral Superphosphate Granular Phosphoric Anhydride 42-45%

WALLACH - GRACER EXPORT CORP. MANUFACTURERS

55 WEST 42nd STREET, NEW YORK 36, N. Y., U. S. A. Tel: LO 4-5490 Cable: WALGRAY

Depend on RODGERS BLENDERS and BATCH MIXERS



- EASY TO CLEAN . BALL BEARINGS SPECIAL DESIGN PACKING GLANDS
- DUST-TIGHT . LEAKPROOF . WATERTIGHT

We are also manufacturers of Agitators, Kettles, Pawder and Paste Fillers . . . We can build to specifications to fit your special requirements!

REQUEST CATALOG NO. 43

ONARCH WEED SPRAYS



Exclusive Western Distributor: W. A. Westgate, Davis, Calif.

PYROPHYLLITE

Ideal As A

DILUENT

CARRIER

INSECTICIDES

GLENDON PYROPHYLLITE COMPANY


10 EAST 40th ST.

NEW YORK 16, N. Y.

Plant and Mines Located at GLENDON, N. C.

Ask for Our Pamphlet

Formerly Glendon Division of Carolina Pyrophyllite Co. 0-------



TOXISOL "A"

the new, low-cost DDT solvent. Useful for other insecticidal requirements, too. For prices, specifications and typical tests, write:

Richfield Oil Corporation

555 South Flower St. - Los Angeles 17 MANUFACTURERS OF RICHFIELD WEEDKILLER "A" (CONTACT HERBICIDE) and RICHFIELD AQUATIC WEEDKILLER dustry and its problems made him an authority not only in the United States, but abroad as well. He has for many years contributed technical essays on the subject of fertilizers to the Encyclopoedia Britannica, and is currently working on a new manuscript for it.

In all, Mr. Lodge has spent the past 44 years in the fertilizer industry. He was graduated in 1908 from the University of Illinois, and began working that year for Armour & Co. as a chemist. Through the years he advanced into more responsible positions and in 1933, was made assistant director of manufacturing.

Mr. Lodge expects to reside on his farm near Forkville, Pennsylvania, but will continue to maintain a desk at the NFA offices in Washington. He will be available to the Association for consultation at various times.

Locusts in Philippines

With swarms of locusts attacking many agricultural areas in the Philippines, including central Luzon, where the major portion of the nation's rice is grown, reports from the islands say that rice fields, corn fields and cocoanut trees have been stripped by the insects.

Late in August, the government had mobilized manpower and equipment to combat the infestation, and all available insecticides had been requisitioned for the task. Planes equipped with booms for spraying were being pressed into service.

According to reports from the Islands, the government had been slow to heed warnings of the impending attack, so that hasty preparations had to be made after the arrival of the locusts.

S. H. Doggett, Mfgr., Dies

Sidney H. Doggett, 54, president of Doggett-Pfeil Co., Springfield, N. J., died August 16 at a Morristown (N. J.) hospital. Mr. Doggett was one of the founders of the firm, established in 1921, and prominent in the manufacture of agricultural and horticultual chemicals. He was also a director of Doggett-Pfeil Co. of New York, Inc. His widow, 2 sons and a brother survive.

FERTILIZER OUTLOOK

(Continued from Page 44)

phosphate rock, mostly for fertilizer purposes.

Since 1940 the annual imports of potash have been running at less than 30,000 short tons (except in 1950 when imports jumped to 200,000 tons) compared with average exports of approximately 65,000 tons.

A major concern to both farmers and fertilizer manufacturers is the problem of transportation costs. Between 1939 and 1950 transportation costs accounted for 10 to 14 percent of the value of fertilizer at its destination. This does not include transportation costs of raw materials to fertilizer manufacturers. During the same period, transportation costs of phosphate rock accounted for 35 to 50 percent of its value at destination.

One of the most promising opportunities for lowering the cost of plant nutrients to the farmer is by increasing the concentration of mixed fertilizers, thereby reducing transportation, handling, storage, and bagging costs per unit of nutrients. It is estimated that a saving of at least \$20 a ton of plant nutrients could be effected by increasing the concentration of the average mixture from 23 percent (the present level) to 30%.

The fertilizer industry anticipates a steady, substantial growth in production and use of the higher concentrations. In 1900, the average plant-nutrient content of mixed fertilizers consumed in the United States was 13.9 percent. By 1949, it had risen to 22.5 percent. The proportion of plant nutrients in mixed fertilizer is expected to rise to 27 or 28 percent by 1975 — provided the trend continues at the same rate of increase.

The transportation advantages of higher concentrations spring from the fact that — with certain exceptions — the same scale of rail and motor rates usually applies to all types of fertilizer and fertilizer materials.

FRY BAG SEALER



Makes Sift-Proof Seals in Heavy Weight Paper Bags

Fry Model CSG automatically makes a double folded sift-proof heat seal in the top of any heavy weight paper bag. The first fold is securely heat scaled; the second is glued for extra safety.

Bags handled include polyethylene and pliafilm lined, polyethylene coated and those with thermoplastic top sealing bands. Simple adjustments for bags of various heights. This model also handles bags which are not heatsealable by gluing the folds.

Machine above is perfect for granular or fine products such as insecticides, chemicals, powdered points, fertilizers, dog foods, etc.

Other models available . . . when writing, please submit a sample of your bag and your product.

GEORGE H. FRY COMPANY

167 Front Street N. Y. 7, N. Y.

ATTENTION DUST MIXERS!

We are now offering complete, integrated blending and impregnation systems for handling practically all basic chemicals in formulating concentrates and finished dusts. Contact our engineers for details.

THE YOUNG MACHINERY COMPANY

Florida Field Trials

Field screening of agricultural chemicals. Thanks to the agricultural chemical industry for the many trials we have been privileged to run during this season.

New trials will be started in September. May we have your reservations of space at an early date. This will insure you a place in the trials and more efficient handling of the work.

DR. G. R. TOWNSEND

Box 356

Belle Glade, Fla.

BORDEAUX MIXTURE

ANSBACHER - OXO BRAND

12 x 4 lb. bags per carton

BELOW MARKET

40,000 lbs. 9¢ lb. FOB, N. Y.

WRITE OR WIRE

Barclay Chemical Co., Inc. NEW YORK 13, N. Y. WORTH 4-5120

75 VARICK STREET

"COHUTTA"

POWDERED TALC

An excellent carrier for insecticides and fungicides. Produced by

Cohutta Talc (

Georgia

DOLOMITIC

ANALYSIS GUARANTEED



LIMESTONE

. 60% Calcium Carbonate 39% Magnesium Carbonate

NILLINGHAM-LITTLE STONE COMPANY

1201-4 Healey Building, Atlanta 3, Ga.

"38 Years Service to the Fertilizer Industry in the Southeast"

How to Increase Use

A SSUMING sufficient supplies, there are several important ways that greater use of fertilizer for increased food and fiber production can be encouraged. In general, these are: (1) public policy, (2) technological progress, (3) education, and (4) economic feasibility.

Public policy, in the field of fertilizer use, favors efficient utilization of our soil resources to contribute continually to the highest possible standard of living for the people of the United States.

Technological progress-development of improved materials — both from the standpoint of plant-nutrient content and ease of handling will have a marked effect on future use of fertilizer. Much progress has been made with respect to certain crops in determining the combinations of practices which give far greater results than practices applied separately.

Education, through the dissemination of knowledge and skills in the use of fertilizer and demonstration of its practical worth in farming, has also been supported by public policy. Educational activities on the part of Government and industry through radio, television, publications, meetings, tours, and demonstrations —will provide the farmer information to help get the most from fertilizer.

Finally, there is economic feasibility — a real test for the introduction or continuation of specific farming practices. Whether it "pays," in the final analysis, will largely determine how much fertilizer is used.

The favorable influence of coordination of the research, educational, and economic factors under the guidance of public policy is becoming increasingly evident since the close of the Second World War. The progress made indicates much greater opportunity in the future.

Fertilizer in new forms and formulas and the application of fertilizers in combination with other field operations, such as seeding, tillage, and the use of herbicides and irrigation water, point to areas of present research which may become more diverse and more extensive.

TECHNICAL BRIEFS

(Continued from Page 73)

lasts longer in poor soils; a shorter time in soils high in organic matter. Results of experiments with samples of 85 different soil types gathered from seven eastern states showed that persistence of DDT was about the same in sandy, silt, or clay loam, but much greater in soils made up primarily of sand.

In terms of dead Japanese beetle grubs, laboratory tests involving more than 223,000 of these soil-dwelling pests showed DDT beginning to lose its killing power only in the fifth year after application. A single treatment with DDT at the 25-pound-per-acre rate resulted in 98 percent grub kills within about four weeks after the grubs were put in the soil, during the first four years. It required slightly more than five weeks to get the same kill during the fifth year.

Insect Resistance in Crop Plants

by Dr. Reginald H. Painter

520 pages, price \$9.80

Here is a complete analysis of the relationships between crops and phytophagous insects together with a full analysis of the insect-resistant varieties of important crops thus far developed.

CONTENTS

Mechanisms of Resistance

Factors Affecting the Expression or the Permanence of Resistance

Resistance to Insects in Wheat

Resistance to Insects in Corn

Resistance to Insects in Cotton

Resistance to Insects in Sorghums

Resistance to Insects in Potato

Methods and Problems in Breeding for Resistance to Insects in Crop Plants

Add 3% Sales Tax in N. Y. City

AGRICULTURAL CHEMICALS

175 Fifth Ave.

New York 10, N. Y.

LOOKING FOR AN EASTERN DISTRIBUTOR FOR YOUR FARM CHEMICALS?

SEE F&B*

MIXERS, PACKERS and DISTRIBUTORS of:

- · INSECTICIDES
- · FUNGICIDES
- · FERTILIZERS

*FAESY & BESTHOFF, INC.

325 SPRING ST., NEW YORK 13, N. Y.

Tel. Algonquin 5-7300

Plant and Warehouse: Hicksville, L. I.

Classified Advertising

Rates for classified advertisements are ten cents per word, \$2.00 minimum, except those of individuals seeking employment, where the rate in five cents per word, \$1.00 minimum. Address all replies to Classified Advectisements with Box Number, care of AGRICULTURAL CHEMICALS, 175 Fifth Ave., New York 10. Clessing date: 25th of preceding month.

Positions Wanted:

Ph.D. PLANT PATHOLOGY, M. A. organic chemistry. Considerable experience field crop disease control, herbicides, bio-assay and development of agricultural chemicals field in general. Wishes responsible position, research or development. Address Box 667, c/o Agricultural Chemicals.

BIOLOGIST, Ph.D., 20 years biological research experience, desires position as head of biological research program in agricultural chemicals. Has proved capabilities as research leader. Excellent knowledge of chemistry. Enthusiastic, hard worker. Married. Good health. Address Box No. 668, c/o Agricultural Chemicals.

Positions Available

CHEMICAL ENGINEER: Real opportunity for young man not over 30 years. No engineer so employed; therefore, individual initiative and developments will not be lost through maze of personnel. Variety of engineering duties. Agricultural chemical background desirable, but not essential. Philadelphia area. Address Box No. 669, c/o Agricultural Chemicals.

SALESMAN WANTED: Established eastern agricultural chemical firm desires aggressive salesman for west coast with headquarters in Los Angeles area. Excellent opportunity for sincere individual. Address Box No. 670, c/o Agricultural Chemicals.

OPPORTUNITY—Technical-minded salesmen with chemical, agronomy or related background and proven sales record to represent leading manufacturer of agricultural chemicals in midwestern and southeastern states. Challenging combination of sales and development. Qualified men will find permanent positions with attractive growth opportunity. Reply in confidence to Box No. 671 c/o Agricultural Chemicals.

AGRICULTURAL CHEMIST with experience in liquid insecticide manuacturing, position in the far West. References required. Replies held in confidence if desired. Address Box 672, c/o Agricultural Chemicals.

AGRICULTURAL & INDUSTRIAL CHEMICAL SALESMAN required by expanding firm. Territory—southern states, east of the Mississippi River. Prefer southern resident centrally located. Please state background and experience. Excllent opportunity. All replies confidential. Address Box 673, c/o Agrcultural Chemicals.

Miscellaneous

Buy-Trade-Sell: Phenol, Naphthalene, Para, DDT, BHC, Pyrethrum, Glycols, Cellosolves, Ethanolamines. Other Chemicals, Solvents, etc. Chemical Service Corporation, 86-02 Beaver St., New York 5, N. Y.

FOR SALE: 20,000 lbs. 10 Gamma BHC dust concentrate; 30,000 lbs. 50% DDT dust concentrate, 50 lb. 4 ply paper bags. Nationally known brand Excellent condition. Will sacrifice. Send us your offers. Chemical Formulators, Inc., Box 35, North Charleston, W. Va.

FOR SALE: Brush control business. Netted \$10,000 or \$40,000 gross in '51. Engaged in brush control service along hwys. and right of ways in N. Y. State. Farms and pvt. property offer excellent potential for expansion. Sale incls. 4 acres of land, equip. (valued at \$30,000) and good will. A sound investment opportunity at \$90,000. No. 2H-5412 Affiliated Brokers of Charles Ford & Assoc., Inc., Dept. 2312, 10 N. Clark, Chicago, Illinois.

ACS to Alantic City Meet

Three days of the week-long fall meeting of the American Chemical Society at Atlantic City, N. J., were to be largely devoted to the section on fertilizer technology. These days, September 17, 18 & 19, were to feature scores of papers on fertilizer manufacturing, application, and chemistry.

Dr. S. F. Thornton, F. S. Royster Guano Co., Norfolk, Va., is chairman of the section and J. D. Romaine, American Potash Institute, Washington, D. C., secretary.

The Division of Agricultural and Food Chemistry was to conduct a number of symposia, including one on the formulation and action of herbicides. Dr. A. S. Crafts, University of California, was to preside at this symposium.

Presentation of the ACS Priestly Medal, highest honor in American chemistry, was to be presented to Dr. Samuel Colville Lind, Carbide & Carbon Chemicals Corp., Oak Ridge, Tenn., as one of the highlights of the meeting.

ALVIN J. COX, Ph.D.

Chemical Engineer and Chemist (Formerly Director of Science, Government of the Philippine Islands. Retired Chief, Bursau of Chemistry, State of California, Department of Agriculture.)

ADVISER ON AGRICULTURAL CHEMICAL PROBLEMS AND INVESTIGATIONS

Consultant in reference to apray injury and damage, claims, including imports of fruits and nuts, formulas, labeling, advertising and compliance with law.

8 1118 Emerson Street
9 Palo Alto, California

Theodore Riedeburg Associates

Sales Consultants

and

Manufacturers' Representatives

on

Agricultural Chemicals

415 Lexington Ave., Rm. 808 New York 17, New York MU rray Hill 7-1488-89

Carl N. Andersen, Ph.D.

Consulting Chemist

Phenyl Mercury Compounds Insecticides, Fungicides-

> 342 Madison Ave. New York 17, N. Y. Phone: VA 6-0492

Laboratory: Briarcliff Manor, New York
Phone: Briarcliff 6-1550

CONSULTING ENTOMOLOGIST

Insecticides — Formulation Plant Pathology—Research Entomology—Legal Service

Author of "Chemistry & Usos of Insecticides"

DR. E. R. de ONG 926 Stannage Ave. Albany 6, Calif.

Advertisers' Index

Accessories Mfg. Co.	138	Fulton Bag & Cotton Mills	. 10
Agricultural Chemicals, Inc.	July	Geigy Co.	
Alrose Chemical Co.	June	General Chemical Division, Allied	. ,
Aluminum Co. of America		Chemical & Dye Corp3rd	Cover
American Agricultural Chemical Co	Aug.	Georgio Tolc Co.	
American Cyanamid Co.	68	Glendon Pyrophyllite Co.	
American Polymer Corp		Greeff & Co., R. W.	
American Potash & Chem. Carp.	Aug.		
Andrews, W. R. E. Sales, Inc.	. 105	Hohn, Inc.	129
Antara Chemicals, Division of General		Hammond Bag & Paper Co	. 27
Dyestuffs Corp.	April	Harte, John J.	
Arkell & Smiths	Aug.	Heckathorn & Co.	
Armour & Co.	21	Highway Equipment Co.	
Ashcraft-Wilkinson Co.	13,86	Hercules Powder Co	
Atlas Powder Co.	. 72	Hercules Steel Products Carp	
Attapulgus Clay Co.		Hough Co., Frank G.	
		Huber, J. M. Corp	
Barclay Chemical Co	142	Hudson Pulp & Paper Carp	. July
Baker H. J. & Bro.		International Clay Com	100
Bagpak Division, International Paper C		International Clay Carp.	121
Baughman Mfg. Co.		Irvington Smelting & Refining Works	. 131
Beckman Instrument Inc.		Jaite Co.	127
Bemis Bro. Bag Co.		Johns-Manville Co.	
Berkshire Chemicals, Inc.		Johnson, C. S. Co.	
Betner, Benj. C. Co.	106	Kolker Chemical Works, Inc.	
California Spray Chem. Co.	56	Koppers Co.	
Chase Bag Carp.		Kroft Bag Co.	
Chemical Construction Corp.	. Aug.	Kron say Co	Aug.
Coddington Mfg. Co., E. D.	. 134	Lion Oil Co	. 23
Cohutta Talc Co.			
Combustion Engineering-Superheater,		Marietta Concrete Corp.	
Inc. Raymond Pulverizer Div.	17	McLaughlin Gormley King Co.	
Columbia Southern Chemical Corp		Mente & Co.	
Cooper, Wm. & Nephews, Inc.		Mercantile Agencies Export Corp	
Cox, Dr. Alvin J.		Michigan Chemicals Corp.	
		Mine Safety Appliances Co.	
Davies Nitrate Co., Inc.	138	Monarch Manufacturing Works, Inc	
Davison Chemical Corp.	. Aug.	Monsanto Chemical Co	16, 50
de Ong, Dr. E. R.		National Aniline Division, Allied	
Diamond Alkali Co.		Chemical & Dye Corp	. 11
Dow Chemical Co.		National Agricultural Chemical Ass'n.	
du Pont de Nemours & Co., E. i2		Naugatuck Chemical Division, U. S. Rubber Co.	62
Edco Corp.	123	Niggara Chem. Div. Food Machinery &	
		Chem. Corp.	
Entoma		Ninol Laboratories, Inc.	7
Emulsol Corp.		Nitrogen Division, Allied Chemical &	
Ethyl Corp.		Dye Corp.	
Eston Chemicals, Inc.	101	Nopco Chemical Corp.	
Faesy & Besthoff, Inc.	143	Noury & Van der Lande	
Fischbein Co., Dave	130	Object to the Samueline Inc.	110
Flag Sulphur & Chemical Co.		Oberdorfer Foundries, Inc.	110
Floridin Co.		Pacific Coast Borax Co.	August
Fry Co., Geo. H.	141	Penick, S. B. & Co.	
Pry Co., Geo. H.	A	een checked carefully but no responsibility	

Fulton Bag & Cotton Mills	10
Geigy Co.	91
General Chemical Division, Allied	
Chemical & Dye Corp3rd	Cove
Georgia Tak Co.	12:
Glendon Pyrophyllite Co.	
Greeff & Co., R. W.	13
Hahn, Inc.	12
Hammond Bag & Paper Co.	
Harte, John J.	
Heckathorn & Co.	
Highway Equipment Co.	8
Hercules Powder Co	21
Hercules Steel Products Carp	
Hough Co., Frank G.	
Huber, J. M. Corp.	. 3
Hudson Pulp & Paper Carp	Jul
International Clay Carp.	10
Irvington Smelting & Refining Works	12
irvingion smelling a kenning works	1.3
Jaite Co.	12
Johns-Manville Co.	-
Johnson, C. S. Co.	12
Kolker Chemical Works, Inc.	
Kraft Bag Co.	Aug
Krom Bag Co	Mug
Lion Oil Co	
Marietta Concrete Corp.	13
McLaughlin Gormley King Co.	11
Mente & Co.	13
Mercantile Agencies Export Corp	Aus
Michigan Chemicals Corp.	2
Mine Safety Appliances Co.	13
Michigan Chemicals Corp. Mine Safety Appliances Co. Monarch Manufacturing Warks, Inc.	14
Monsanto Chemical Co1	8, 8
National Aniline Division, Allied	
Chemical & Dye Corp	1
National Agricultural Chemical Ass'n.	
Naugatuck Chemical Division, U. S.	
Rubber Co.	6
Niggara Chem. Div. Food Machinery &	-
Chem. Corp.	1
Ninol Laboratories, Inc.	
Nitrogen Division, Allied Chemical &	
Dye Corp	11
Nopco Chemical Corp.	
Noury & Van der Lande	9
Oberdorfer Foundries, Inc.	
Pacific Coast Borax Co	ugu
Penick, S. B. & Co	1

Pennsylvania Industrial Chemical	
Corp.	103
Pennsylvania Salt Manufacturing	
Co.	83
	113
Phelps Dodge Refining Corp.	
Phillips Chemical Co	104
Pioneer Chemical Associates	132
Pioneer Pyrophyllite Corp.	107
Pittsburgh Agricultural Chemical Co.	
a Division of Pittsburgh Coke and	
Chemical Co.	26
Chemical Co. Pittsburgh Plate Glass Co., Corona	
Chem. Div.	June
Polash Company of America	3
Prentiss Chemical Co.	78
Private Brands, Inc.	138
Poulsen, A. E. & Co. Powell, John & Co	0.0
Powell, John & CoZnd C	.gvei
Pulverizing Machinery Co	Aug.
Raymond Pulverizer Division, Combus-	
tion Engineering-Superheater, Inc	17
from Engineering-Superneurer, Inc	
Remsen Chemicals, Inc.	
Richfield Oil Corp.	140
Rieke Metal Prod. Corp	8.5
Riedeburg, Theodore Associates	144
Rodgers, George G. Co. Royster Guano Co.	140
Royster Guano Co.	119
Seacoast Laboratories, Inc	60
Shell Chemical Co	98
Southeastern Clay Co.	117
Southwest Potash Carp	70
Spencer Chemical Co	Aug.
Spraying Systems Co.	136
Stauffer Chemical Co.	111
Sprout, Waldron & Co	118
Sturtevant Mill Co.	64
	04
Tennessee Corp. Tennessee Products & Chemical Carp Texas Gulf Sulphur Ce. Thompson-Hayward Chemical Co. Titlastad, Nicolay Carp	112
Tennessee Products & Chemical Carp	Aug.
Texas Gulf Sulphur Co.	Aug.
Thompson-Hayward Chemical Co.	116
Titlestad, Nicolay Carp. Tobacco By-Products & Chemical Carp. Townsend, Dr. G. R.	July
Tobacco By-Products & Chemical Corn.	7.4
Townsend Dr G 9	143
Townsend, Dr. Q. R.	144
Union Bag & Paper Corp.	89
Union Special Machine Co. United Chemical Co. U. S. Industrial Chemicals, Inc4th (Aug
United Chemical Ca.	135
U. S. Industrial Chemicals, Inc4th (Cover
U. S. Potash Co.	6
U. S. Steel Corp.	25
v	87
Vanderbill Co., R. T.	87
Veluical Cara.	
virginia-Carolina Chemical Corp.	/4
Warren Div., Amer. Steel Dredge Co.	July
Warren Div., Amer. Steel Dredge Co Whittaker Clark & Daniels, Inc.	130
Williams Potent Crusher & Pulsarines Co	54
Williagham Little Stone Co	145
Williams Patent Crusher & Pulverizer Co Willingham-Little Stone Co. Wallach-Gracer Export Corp	120
Wallach-Gracer Export Corp	, 139
Wisconsin Alumni Research	
Foundation	
Woodward & Dickerson, Inc.	
Woudhuysen, H. L. & Associates	
Wyandatte Chemicals Co	
Young Machinery Co	
roung mochinery co	1.44

MEETING CALENDAR

National Agricultural Chemicals Assn., Easex & Sussex Hotel. Spring Lake. N. J., Sept. 3, 4, 5. American Phytopathological Society, Cornell University. Ithaca. N. Y., September 9-12.

Fertilizer Section. American Chemical Society. Convention Hall. Atlantic City. N. J., Sept. 14-13.

Association of Official Agricultural Chemists. Shoreham Hotel. Wash-ington. D. C., September 29 to October 1.

Association of American Feed Control Officials, Shoreham Hotel, Washington, D. C., October 1 & 2.

Association of American Fertilizer Control Officials, Shoreham Hotel, Washington, D. C., October 3, Association of Economic Poisons Control Officials. Shoreham Hotel.
Washington, D C., October 4.
National Pest Control Association.
Rice Hotel, Houston, Texas.

Rice Hotel. Houston. Texas.
October 20-22.
Meeting of Fertilizer Section of National Safety Congress. Chicago.
Ill. October 22 & 23.
Scuthwide Chemical Conference.
Auburn Alabama. October 23-25.
Sixth Annual Beltwide Cotton
Mechanization Conference. Bakersfield California Oct. 22-24. ersfield, California, Oct. 22-24.

California Fertilizer Association. Desert Inn. Palm Springs, Calif.. Nov. 10-12.

New York State Insecticide & Fun-gicide Conference, Ithaca, N. Y., November 11-13.

November 11-13.
National Fertilizer Association Fall
Meeting, Boney Plaza Hotel,
Miami, Fla., November 19-21.
Joint meeting, North Central Weed
Control Conference and Western
Canadian Weed Conference,
Royal Alexandra Hotel, Winnipeg, Canada, December 8, 1952.
American Association of Economic
Entomologists. Hotel Bellevue
Stratford, Phila., Pa., Dec. 13-18.

TALE ENDS

AROLD Noble, vice-president of S. B. Penick & Co., New York, and Prof. Carl Paul Link of the University of Wisconsin, presented a round-table discussion on rodent control which was broadcast by "Voice of America" on July 31. The discussion, which covered chiefly large scale rodent control with warfarin, lasted for 15 minutes and was beamed to the British Isles, Germany, other parts of Europe and North Africa.

Prof. Link, discoverer of the anti-coagulant rodenticide powers of

warfarin, handled scientific phases of the broadcast while Mr. Noble outlined the history of the use of the material in rodent control in the United States.

The original round-the-world "Voice" broacast has been rearranged for a rebroadcast in native languages to all countries of the Far East, Middle East, and throughout Latin America. When the rebroadcast program is completed it will cover the full scope of the "Voice of America" broadcasting range.

Cornell Veg-News recently carried a brief item lauding the merits of a "new fertilizer," called "ERUNAM". It termed the article an "inert soil conditioner," a "selective pesticide," killing harmful weeds, bugs and diseases while fraternizing with "approved" ones . . . contains no "nasty chemicals; it's purely organic" and "one pound of this concentrated product is equivalent to 16 ounces." The product? Erunam, spelled backwards, spells "MANURE."

We still think that someone should come forth with a product, "Drawrof," which spelled backwards, spells "forward".

Crop duster planes seem to have a number of functions aside from applying various kinds of toxicants to various agricultural crops.

The latest such role is that of a rescue taxi to hustle people from communist-dominated lands.

Miss Hana Pavlickova, 24, upon arriving in the U. S. recently as a refugee, said she was one of four persons who squeezed into a single-scater crop-dusting plane in Czechoslovakia and was hustled over the border via this improvised airlift. So crowded was the plane, she said, that one of her legs was dangling outside as the pilot took off.

The plane had been dusting fungicides on potatoes that Czech propagandists claimed had been infected with a "plague" by American airmen.

J. F. Prochaska retired recently after 42 years of service in the agricultural chemicals sales department of E. I. du Pont de Nemours & Co., most recently in the Cleveland office. Mr. Prochaska was originally with the Grasselli Company and joined Du Pont when Grasselli was incorporated as a division in 1928. Mr. Prochaska was active in forming the Ohio Pesticide Institute in 1947.



Drill!

A GAIN, and again, and again. Just like the kind of advertising that pays off in the long run, — regular advertising month in and month out, again and again and again.

And the kind, for exmple, that can pay off well in the field of insecticides, fertilizers, fungicides, herbicides and other chemical specialties, are the ads in

AGRICULTURAL CHEMICALS

175 FIFTH AVE.

NEW YORK 10, N. Y.

Formulators! Dust Mixers! Use "GENERAL CHEMICAL"

BASE MATERIALS and TOXICANTS and be sure of the best!

Check Your Needs Here...Then Call General Chemical!

BASIC CHEMICALS FOR AGRICULTURE

DDT

Dust Base, 50% & 75%

LEAD ARSENATE

Standard Astringent

LINDANE

Technical Dust Base, 25% Emulsifiable Concentrate, 20%

Technical (15%, 36% & 90% Gamma) Dust Base, 12% Gamma

DDD

Dust Base (50% DDD)

PARATHION

Dust Base Emulsifiable 284 lbs. per gal.

CALCIUM **ARSENATE**

Standard Low Lime

ZIRAM

(76% Zinc Dimethyl Dithiocarbamate)

FERBAM

(76% Ferric Dimethyl Dithiocarbamate)

Acid, Amine, Esters **Formulations**

2.4.5-T

Acid, Amine, Esters Formulations

TCA SODIUM SALT

90% Dry Powder 50% Liquid Concentrate

POTASSIUM CYANATE

Technical Cotton Defoliant

DINITRO

Pre-Emergence Herbicides

ORGANIC MITICIDES

"Aramite" Genite¹ 883 (p-chlorophenyl p-chlorobenzene sulfonate)



SPRAYS & DUSTS

Agricultural Chemical Department

GENERAL CHEMICAL DIVISION

ALLIED CHEMICAL & DYE CORPORATION
40 Rector Street, New York 6, N. Y.

Serving Agriculture from Coast to Coast

Stored Corn<u>can</u> be kept Weevil-Free



handling or precautions.

It takes only one pound of Grain Protectant to treat 10 bushels of corn, and it is packaged in 50, 25 and 10 pound bags to supply the convenient amount for any size crop.

This new method for control of stored grain insects gives the agricultural chemical industry a whole new field of service for helping the American farmer fulfill the nation's food requirements.

But for the farmer to have this Protectant when he needs it, his dealers must get their stocks on hand right away!